arXiv:0802.1088v1 [math.GR] 8 Feb 2008

MSC2000 20D25, 20D06, 20E07, 20G40

CARTER SUBGROUPS OF FINITE GROUPS

E.P.Vdovin

Contents

1 Introduction **1**
- 1.1 General characteristic of the results of the paper . . . 1
- 1.2 Notation and results from group theory . . . 3
- 1.3 Linear algebraic groups . . . 4
- 1.4 Structure of finite groups of Lie type . . . 7
- 1.5 Known results . . . 10

2 Conjugacy criterion for Carter subgroups **11**
- 2.1 Main results of this section . . . 11
- 2.2 Preliminary results . . . 12
- 2.3 Proof of Theorem 2.1.4 . . . 14
- 2.4 Some properties of Carter subgroups . . . 14

3 Conjugacy in simple groups **16**
- 3.1 Brief review of results of the section . . . 16
- 3.2 Preliminary results . . . 17
- 3.3 Almost simple groups . . . 24

4 Semilinear groups of Lie type **28**
- 4.1 Basic definitions . . . 28
- 4.2 Translation of basic results . . . 30
- 4.3 Carter subgroups of special type . . . 34

5 Carter subgroups of semilinear groups **38**
- 5.1 Brief review of results of this section . . . 38
- 5.2 Carter subgroups of symplectic groups . . . 39
- 5.3 Группы с автоморфизмом тройственности . . . 42
- 5.4 Classification theorem . . . 43
- 5.5 Carter subgroups with unipotent radical . . . 44
- 5.6 Carter subgroups without unipotent radical . . . 47
- 5.7 Carter subgroups of finite groups are conjugate . . . 50

6 Existency criterion **50**
- 6.1 Brief review Краткий обзор результатов параграфа . . . 50
- 6.2 Criterion . . . 51
- 6.3 Example . . . 53
- 6.4 Classification of Carter subgroups . . . 54

List of Tables **56**

References **57**

ABSTRACT: In the paper it is proven that Carter subgroups of a finite group are conjugate. A complete classification of Carter subgroups in finite almost simple groups is also obtained.

KEY WORDS: Carter subgroup, finite simple group, group of Lie type, linear algebraic group, semilinear group of Lie type, semilinear algebraic group, conjugated powers of an element

1 Introduction

The present paper is a slightly shorten version of doctoral thesis "Carter subgroups of finite groups". The results of the thesis were published in [3–6], [37], and [43]. Although, in the thesis all papers were rewritten in order to obtain a coherent text. This rewritten text is given in the present paper.

1.1 General characteristic of the results of the paper

We recall that a subgroup of a finite group is called a *Carter subgroup* if it is nilpotent and self-normalizing. By a well-known result, any finite solvable group contains exactly one conjugacy class of Carter subgroups (see [14]). If a group is not assumed to be finite, then Carter subgroups can be even nonisomorhic. Indeed, if N_1, N_2 are two nonisomorhic nilpotent groups, then they are Carter subgroups in their free product. On the other hand, a finite nonsolvable group may fail to contain Carter subgroups, the minimal counter example is the alternating group of degree 5. Although there is not known a finite group containing nonconjugate Carter subgroups, and the following problem, due to R.Carter, is known.

Problem 1.1.1. (**Conjugacy Problem**) Are Carter subgroups of a finite group conjugate?

This problem for several classes of finite groups closed to be simple was investigated by many authors. In symmetric and alternating groups Carter subgroups were classified by L.Di Martino and M.C.Tamburini (see [25]). In every group G such that $\mathrm{SL}_n(q) \leq G \leq \mathrm{GL}_n(q)$ Carter subgroups were classified by L.Di Martino and M.C.Tamburini, and in the case $G = \mathrm{GL}_n(q)$, by N.A.Vavilov (see [26] and [2] respectively). For symplectic groups $\mathrm{Sp}_{2n}(q)$, general unitary groups $\mathrm{GU}_n(q)$, and, for q odd, general orthogonal groups $\mathrm{GO}_n^{\pm}(q)$ the classification of Carter subgroups was obtained by L.Di Martino, A.E.Zalessky, and M.C.Tamburini (see [27]). For some sporadic simple groups Carter subgroups were found in [22]. In the mentioned above nonsolvable groups Carter subgroups coinside with the normalizers of Sylow 2-subgroups, and hence are conjugate.

A finite group G is called a *minimal counter example to Conjugacy Problem* or simply a *minimal counter example*, if G contains nonconjugate Carter subgroups, but in every group H, with $|H| < |G|$, Carter subgroups are conjugate. In [23] F.Dalla Volta, A.Lucchini, and M.C.Tamburini have proven that a minimal counter example should be almost simple. This result allows to use the classification of finite simple groups to solve Conjugacy Problem.

Note that the using of F.Dalla Volta, A.Lucchini, and M.C.Tamburini result to the classification of Carter subgroups in almost simple groups essentially depends on the classification of finite simple groups. Indeed, in order to use the inductive hypothesis that Carter subgroups in every proper subgroup of a minimal counter example are conjugate, one needs to know that all almost simple groups of order less than the order of a minimal counter example are found. To avoid using the classification of finite simple groups we strengthen the result of F.Dalla Volta, A.Lucchini, and M.C.Tamburini, proving that if Carter subgroups are conjugate in the group of induced automorphisms of every non-Abelian composition factor, then they are conjugate in the group.

To inductive description of Carter subgroups in almost simple groups one needs to know homomorphic images of Carter subgroups and intersections of Carter subgroups with normal subgroups, i. e., the answers to the following problems.

Problem 1.1.2. Is a homomorphic image of a Carter subgroup again a Carter subgroup?

Problem 1.1.3. Is the intersection of a Carter subgroup with a normal subgroup again a Carter subgroup (of the normal subgroup)?

The first problem is closely connected with Conjugacy Problem, namely, if Conjugacy Problem has an affirmative answer, then the first problem also has an affirmative answer. So we shall solve both of these problems by considering Carter subgroups in almost simple groups. It is easy to see that the second problem has a negative answer. Indeed, consider a solvable group Sym_3 and its normal subgroup of index 2, the alternating group Alt_3. Then a Carter subgroup of Sym_3 is a Sylow 2-subgroup, while a Carter subgroup of Alt_3 is a Sylow 3-subgroup. Thus in the paper some properties of Carter subgroups in a group and some of its normal subgroups are found.

The present paper is divided into six sections, including Introduction. In the introduction we give general results of the paper, and also necessary definitions and results.

In the second section we prove that Carter subgroups of a finite group are conjugate, if they are conjugate in the group of induced automorphisms of every its non-Abelian composition factor, thereby strengthen the results of F.Dalla Volta, A.Lucchini, and M.C.Tamburini. In the second section we also obtain some properties of Carter subgroups.

In the third section we consider the problem of conjugacy for elements of prime order in finite groups of Lie type. At the end of the third section, by using the results on conjugacy, we obtain the classification of Carter subgroups in a wide class of almost simple groups.

In the fourth section we introduce the notion of semilinear groups of Lie type and corresponding semilinear algebraic groups, and transfer the results on the normalizers of p-subgroups and the centralizers of semisimple elements in groups of Lie type. We also obtain some additional results on the conjugacy of elements of prime order in these groups.

In the fifth section we complete the classification of Carter subgroups in almost simple groups and prove that Carter subgroups of almost simple groups are conjugate. As a corollary we obtain an affirmative answer to Conjugacy Problem and prove that a homomorphic image of a Carter subgroup is a Carter subgroup.

In the sixth section we study the problem of existence of a Carter subgroup in a finite group, give a criterion of existence, and construct an example showing that the property of containing a Carter subgroup is not preserved under extensions. More over in the last subsection of sixth section we give tables with classification of Carter subgroups in almost simple groups.

I am grateful to my scientific adviser, a corresponding member of RAS, V.D.Mazurov. В. Д. Мазурову. His contribution to my development as a mathematician, an his continuous support are inestimable. I am also sincerely thankful to Professor M.C.Tamburini, who has initiated my work in these problems, and has provided a help during the work. I especially thank Dr. A.V.Vasiliev, Dr. M.A.Grechkoseeva, Dr. A.V.Zavarnitsine, and Dr. D.O.Revin for very useful discussion of the paper, that allows to simplify some proofs and to improve inaccuracies and mistakes. I am also pleased to Professor A.S.Kondratiev for valuable comments, that improving the final text. I am grateful and wish to honor a serene memory of Professor Yu.I.Merzlyakov, who awaken my interest to algebra and group theory.

The work is supported by Russian fond of basic research (projects numbers 99–01–00550, 01–01–06184, 02–01–00495, 02–01–06226 and 05–01–00797), grants of President RF for young scientists (MK–1455.2005.1 and MK–3036.2007.1), SB RAS (gran N 29 for young scientists and Integration project 2006.1.2), and Program "Universities of Russia" (project number UR.04.01.202). A part of the work has been made during my post-doctoral fellowship in the university of Padua (Italy), and I am grateful to this university, to all members of algebra chair, and, especially to Professor F.Menegazzo for support.

1.2 Notation and results from group theory

Out notation is standard. If G is a group, then $H \leq G$ and $H \trianglelefteq G$ mean that H is a subgroup and a normal subgroup of G respectively. By $|G : H|$ we denote the index of H in G, $N_G(H)$ is the normalizer of H in G. If H is normal in G, then by G/H we denote the factor group of G by H. If M is a subset of G, then $\langle M\rangle$ denotes the subgroup generated by M, $|M|$ denotes the cardinality of M (or the order of an element, if there is an element instead of a set). By $C_G(M)$ we denote the centralizer of M in G, by $Z(G)$ we denote the center of G. The conjugate of x by an element y in G is written as $x^y = y^{-1}xy$ (${}^yx = yxy^{-1}$), by $[x, y] = x^{-1}x^y$ we denote the commutator of x, y. The symbol $[A, B]$ means the mutual commutant of subgroups A and B of G. For groups A and B the expressions $A \times B$, $A \circ B$, and $A \leftthreetimes B$ mean direct, central, and semidirect products respectively of A and B with a normal subgroup B. If A and B are subgroups of G such that $A \trianglelefteq B$, then the factor group B/A is called a *section* of G. The Fitting subgroup of G is denoted by $F(G)$, the generalized Fitting subgroup is denoted by $F^*(G)$.

The set of Sylow p-subgroups of a finite group G we shall denote by $Syl_p(G)$. Ifφ is a homomorphism of G, g is an element of G, then G^φ, g^φ are the images of G and g under φ respectively Gphup@G^φ, g^φ. By G_φ we denote the set of stable points of G under an endomorphism φ. By $\mathrm{Aut}(G)$, $\mathrm{Out}(G)$ and $\mathrm{Inn}(G)$ we denote the group of all automorphisms, the group of outer automorphisms, and the group of inner automorphisms of G respectively. If G is a group, we denote by $\mathbf{P}G$ the factor group $G/Z(G)$. An isomorphism $\mathbf{P}G \simeq \mathrm{Inn}(G)$ is known, in particular, if $Z(G)$ is trivial, then $G \simeq \mathrm{Inn}(G)$, and we may assume that $G \leq \mathrm{Aut}(G)$. A finite group G is said to be *almost simple*, if there exists a finite group S with $S \leq G \leq \mathrm{Aut}(S)$, i. e., $F^*(G) = S$ is a simple group. For every positive integer t by $\mathbb{Z}_t$ we denote a cyclic group of order t.

If π is a set of primes, then by π' we denote its complement in the set of all primes. For every positive integer n by $\pi(n)$ we denote the set of prime divisors of n, and by n_π we denote the maximal divisor of n such that $\pi(n_\pi) \subseteq \pi$. As usual we denote by $O_\pi(G)$ the maximal normal π-subgroup of G, and we denote by $O^{\pi'}(G)$ the subgroup generated by all π-elements of G. If $\pi = \{2\}'$ is a set of all odd primes, then $O_\pi(G) = O_{2'}(G)$ is denoted by $O(G)$. If $g \in G$, then by g_π we denote the π-part of g, i. e., $g_\pi = g^{|g|_{\pi'}}$.

Let G be a group, A, B, H be subgroups of G and B is normal in A. Then $N_H(A/B) = N_H(A) \cap N_H(B)$. If $x \in N_H(A/B)$, then x induces an automorphism $Ba \mapsto Bx^{-1}ax$ of A/B. Thus there exists a homomorphism of $N_H(A/B)$ into $\mathrm{Aut}(A/B)$. The image of this homomorphism is denoted by $\mathrm{Aut}_H(A/B)$ and is called a *group of induced automorphisms* of H on the section A/B. In particular, if $S = A/B$ is a composition factor of G, then for each subgroup $H \leq G$ the group $\mathrm{Aut}_H(S) = \mathrm{Aut}_H(A/B)$ is defined. Note that the structure of $\mathrm{Aut}_H(S)$ depends on the choice of a composition series. If A, H are subgroups of G, then $\mathrm{Aut}_H(A) = \mathrm{Aut}_H(A/\{e\})$ by definition.

1.3 Linear algebraic groups

Necessary information about the structure and properties of linear algebraic groups can be found in [12]. Since we consider linear algebraic groups only, we shall omit the word "linear" for brevity.

If $\overline{G}$ is an algebraic group, then by $\overline{G}^0$ we denote the unit component of $\overline{G}$. An algebraic group is called *semisimple*, if its radical $R(\overline{G})$ is trivial, and an algebraic group is called *reductive*, if its unipotent radical $R_u(\overline{G})$ is trivial (in both cases an algebraic group is assumed to be finite). A connected semisimple algebraic group is known (for example, see [12, Theorem 27.5]) to be a central product of connected simple algebraic groups, while a connected reductive algebraic group $\overline{G}$ is known to be a central product of a torus $\overline{S}$ and a semisimple group $\overline{M}$ with $\overline{S} = Z(\overline{G})^0$, $\overline{M} = \left[\overline{G}, \overline{G}\right]$, and $\overline{S} \cap \overline{M}$ is finite.

If $\overline{G}$ is a connected reductive algebraic group, then let $\overline{T}$ be its maximal torus (by a *torus* we always mean a connected diagonalizable (d-) group). The dimension of a maximal torus is called a *rank* of an algebraic group. By $\Phi(\overline{G})$ the root system of of $\overline{G}$ with respect to a maximal torus $\overline{T}$ (it does not depend on the choice of a maximal torus) is denoted, and $W(\overline{G}) \simeq N_{\overline{G}}(\overline{T})/\overline{T}$ is the *Weyl group* of G. If $\overline{G}$ is a reductive group of rank n, then the dimension of the centralizer of any its element is not less than n. An element is called *regular* if the dimension of its centralizer is equal to n. In particular, a semisimple element s is regular, if $C_{\overline{G}}(s)^0$ is a maximal torus of $\overline{G}$.

Recall that or every root system Φ there exists a set of roots $r_1, \ldots, r_n$ such that each root of Φ can be uniquely written as $\sum_{i=1}^n \alpha_i r_i$, where all coefficients α_i are integers, and either nonnegative, or nonpositive. Such a set of roots is called a *fundamental set* of Φ, and its elements are called *fundamental roots.* At that a fundamental set is a basis of $\mathbb{Z}\Phi \otimes_{\mathbb{Z}} \mathbb{R}$. The dimension of $\mathbb{Z}\Phi\otimes_{\mathbb{Z}} \mathbb{R}$ is called a *rank* of Φ. Note that the ranks of $\overline{G}$ and of its root system $\Phi(\overline{G})$ are equal. Below we assume that all fundamental roots are positive. Then a root r is *positive* if and only if it is a linear combination of fundamental roots with nonnegative coefficients. For a root system Φ by Φ^+ (Φ^-) the set of all positive (negative) roots is denoted. The number $h(r) = \sum_{i=1}^n \alpha_i$ is called a *height* of $r = \sum_{i=1}^n \alpha_i r_i$. In every irreducible root system Φ there exists a unique root of maximal height, which is denoted by r_0 below. Note that the Weyl group $W(\Phi)$ of a root system Φ is generated by reflections in fundamental roots, which are called *fundamental reflections.* If we denote by $l(w)$ the minimal number of multipliers in a decomposition of w into the product of fundamental reflections, a (*length*), then there exists a unique element of maximal length, denoted by w_0 below, that is a unique element of the Weyl group mapping all positive roots into negative roots. In general, $l(w)$ is equal to $|\Phi^- \cap (\Phi^+)^w|$, i. e., to the number of positive roots, that w maps into negative roots.

Let $\overline{G}$ be a connected simple algebraic group, π be its exact rational representation, Γ_π be a lattice generated by weights of the representation π. By Γ_{ad} we denote the lattice generated by the roots of Φ, by Γ_{sc} we denote the lattice generated by the fundamental weights. The lattices Γ_{sc}, Γ_π, and Γ_{ad} do not depend on the representation of $\overline{G}$, and the following inclusions $\Gamma_{ad} \leq \Gamma_\pi \leq \Gamma_{sc}$ (see [12, 31.1]) hold. Several distinct algebraic groups, which are called *isogenies*, is known to exist for a given root system. They differs by the structure of Γ_π and the order of the finite center. If Γ_π coinsides with Γ_{sc}, a group $\overline{G}$ is said to be *simply connected*, it is denoted by $\overline{G}_{sc}$. If Γ_π coinsides with Γ_{ad}, a group $\overline{G}$ is said to have an *adjoint type*, it is denoted by через $\overline{G}_{ad}$. Every linear algebraic group with a root system Φ can be obtained as a factor group of $\overline{G}_{sc}$ by a subgroup of its center. The center of $\overline{G}_{ad}$ is trivial, and this group is simple as an abstract group. The factor group Γ_{sc}/Γ_π is denoted by $\Delta(\overline{G})$ and is called a *fundamental group* of $\overline{G}$. The factor group Γ_{sc}/Γ_{ad} depends on the root system Φ only and is denoted by $\Delta(\Phi)$. Хорошо известно, The group $\Delta(\Phi)$ is known to be cyclic, except the root system $\Phi = D_{2n}$, when $\Delta(D_{2n}) = \mathbb{Z}_2 \times \mathbb{Z}_2$ is elementary Abelian of order 4.

Let $\overline{B}$ be a Borel subgroup, $\overline{T} \leq \overline{B}$ be a maximal torus, and $\overline{U} = R_u(\overline{B})$ be a maximal connected unipotent subgroup of $\overline{G}$. There exists a unique Borel subgroup $\overline{B}^-$ such that $\overline{B} \cap \overline{B}^- = \overline{T}$, denote by $\overline{U}^- = R_u(\overline{B}^-)$. If we fix an order on $\Phi(\overline{G})$, then each element $u \in \overline{U}$ (respectively $u \in \overline{U}^-$) can be uniquely written in the form

$$u = \prod_{r\in\Phi^+} x_r(t_r) \tag{1}$$

(respectively $u = \prod_{r\in\Phi^-} x_r(t_r)$), where roots are taken in given order, elements t_r-s are in the definition field of $\overline{G}$ and $\{X_r, r \in \Phi\}$ is a set of *1-dimensional $\overline{T}$-invariant subgroups* (a set of *root subgroups*). The multiplication of elements from distinct root subgroups is defined by *Chevalley commutator formulae.*

Lemma 1.3.1. [15, 5.2.2], (Chevalley commutator formulae) *Let $x_r(t)$, $x_s(u)$ be elements from distinct root subgroups X_r and X_s respectively, $r \neq -s$. Then*

$$[x_r(t), x_s(u)] = \prod_{ir+js\in\Phi;i,j>0} x_{ir+js}(C_{ijrs}(-t)^i u^j),$$

where the constants C_{ijrs} do not depend on t and u.

Substantially this formulae means that the mutual commutant of X_r and X_s is in the group generated by subgroups X_{ir+js}, where $i, j > 0$ and $ir + js \in \Phi$.

Let c_i be the coefficient of a fundamental root r_i in the decomposition of r_0. Primes, dividing c_i-s, are called *bad* primes. The diagram, obtained from the Dynkin diagram by addition of $-r_0$ and its connection with other fundamental roots by usual rule, is called an *extended Dynkin diagram.* Let $\overline{R}$ be a (connected) reductive subgroup of maximal rank of a connected simple algebraic group $\overline{G}$. As we already noted, in this case $\overline{R} = \overline{G}_1 \circ \ldots \circ \overline{G}_k \circ Z(\overline{R})^0$, where $\overline{G}_i$-s are connected simple algebraic groups of rank less, than the rank of $\overline{G}$. More over, if $\Phi_1, \ldots, \Phi_k$ are root systems of $\overline{G}_1, \ldots, \overline{G}_k$ respectively, then $\Phi_1 \cup \ldots \cup \Phi_k$ is a subsystem of Φ. There exists a nice algorithm, due to Borel and de Siebental [13] and, independently, Dynkin [7], of determining subsystems of a root system. One needs to extend the Dynkin diagram to the extended Dynkin diagram, remove some vertices from it, and repeat the procedure for obtained connected components. Diagrams obtained in this way are subsystem diagrams and diagram of any subsystem can be obtained in this way.

In Table 1 we give extended Dynkin diagrams of all irreducible root systems and coefficients of fundamental roots in the decomposition of r_0. The numberring in Table 1 is chosen as in [24].

Table 1: Root systems and extended Dynkin diagrams

Φ	Extended Dynkin diagram
A_n	 $-r_0$ -1 r_1 r_2 r_3 r_n 1 1 1 1
B_n	 r_1 1 r_2 r_3 r_{n-1} r_n $-r_0$ 2 2 2 2 -1
C_n	 $-r_0$ r_1 r_2 r_{n-1} r_n -1 2 2 2 1
D_n	 r_1 r_{n-1} 1 r_2 r_3 r_{n-2} 1 $-r_0$ 2 2 2 r_n -1 1
E_6	 r_1 r_3 r_4 r_5 r_6 1 2 3 2 1 2 r_2 -1 $-r_0$
E_7	 $-r_0$ r_1 r_3 r_4 r_5 r_6 r_7 -1 2 3 4 3 2 1 2 r_2


E_8
r_1 r_3 r_4 r_5 r_6 r_7 r_8 $-r_0$
2 4 6 5 4 3 2 -1
3 r_2
F_4
$-r_0$ r_1 r_2 r_3 r_4
-1 2 3 4 2
G_2
$-r_0$ r_1 r_2
-1 2 3

For every semisimple element $s \in \overline{G}$, where $\overline{G}$ is a connected reductive group, the unit component $C_{\overline{G}}(s)^0$ is a reductive subgroup of maximal rank and $C_{\overline{G}}(s)/C_{\overline{G}}(s)^0 \simeq D \leq \Delta(\overline{G})$ (see Lemma 1.5.2 below).

1.4 Structure of finite groups of Lie type

Our notation and definitions for finite groups of Lie type mainly agree with that of [15] (except the definition of finite groups of Lie type, see below). If G is a finite group of Lie type with the trivial center (we do not exclude non-simple groups of Lie type, such as $A_1(2)$, all exceptions are given in [15, Theorems 11.1.2 and 14.4.1] and cited i Table 2 below), then $\widehat{G}$ denoted the group of inner-diagonal automorphisms of G. In view of [41, 3.2] we have that $\mathrm{Aut}(G)$ is generated by inner-diagonal, field, and graph automorphisms. Note that the definition of a field and a graph automorphisms in the present paper is slightly different from the definitions given in [41], precise definitions are given in subsection 4.1. Since we are assuming that $Z(G)$ is trivial, then G is isomorphic to the group of its inner automorphisms, and so we may assume that $G \leq \widehat{G} \leq \mathrm{Aut}(G)$.

Table 2: Groups of Lie type which are not simple

Group	Properties
$A_1(2)$	Group is solvable
$A_1(3)$	Group is solvable
$B_2(2)$	$B_2(2) \simeq \mathrm{Sym}_6$
$G_2(2)$	$[G_2(2), G_2(2)] \simeq {}^2A_2(3)$
${}^2A_2(2)$	Group is solvable
${}^2B_2(2)$	Group is solvable
${}^2G_2(3)$	$[{}^2G_2(3), {}^2G_2(3)] \simeq A_1(8)$
${}^2F_4(2)$	$[{}^2F_4(2), {}^2F_4(2)]$ is the simple Tits group

Let $\overline{G}$ be a simple connected algebraic group over an algebraic closure $\overline{\mathbb{F}}_p$ of a finite field of positive characteristic p. Here $Z(\overline{G})$ can be nontrivial. An endomorphism σ of $\overline{G}$ is called a *Frobenius map*, if $\overline{G}_\sigma$ is finite, while the kernel of σ is trivial (i. e., σ is an automorphism of G as an abstract group). Groups $O^{p'}(\overline{G}_\sigma)$ are called *canonical finite groups of Lie type*, and every group G, with $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$, is called a *finite group of Lie type*. If $\overline{G}$ is a simple algebraic group of adjoint type, then we shall say that G also has an *adjoint type*. Note that in [15] only groups $O^{p'}(\overline{G})$ are called finite groups of Lie type. But later in [19] R.Carter says that every group $\overline{G}_\sigma$ is a finite group of Lie type, for every connected reductive group $\overline{G}$. Moreover, in [18] and [24], without any comment, every group G with $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$ is called a finite group of Lie type. Thus giving the definition of finite groups of Lie type and of finite canonical groups of Lie type we intend to clarify the situation here. For example, $\mathbf{P}\mathrm{SL}_2(3)$ is a canonical finite group of Lie type and $\mathbf{P}\mathrm{GL}_2(3)$ is a finite group of Lie type. Note that an element of order 3 is not conjugate to its inverse in $\mathbf{P}\mathrm{SL}_2(3)$ and is conjugate to its inverse in $\mathbf{P}\mathrm{GL}_2(3)$. Since such information about the conjugation is important in many cases (and is very important and useful in the paper), we find it reasonable to use such notation.

In general, for a given group of Lie type G (if we consider it as an abstract group) a corresponding algebraic group is not uniquely defined. For example, if $G = \mathbf{P}\mathrm{SL}_2(5) \simeq \mathrm{SL}_2(4)$, then G can be obtained either as $(\mathrm{SL}_2(\overline{\mathbb{F}}_2))_\sigma$, or as $O^{5'}((\mathbf{P}\mathrm{GL}_2(\overline{\mathbb{F}}_5))_\sigma)$ (for suitable σ-s). Hence, for every finite group of Lie type G we fix (in some way) a corresponding algebraic group $\overline{G}$ and a Frobenius map σ such that $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$.

We say that groups ${}^2A_n(q)$, ${}^2D_n(q)$, ${}^2E_6(q)$ are defined over $\mathbb{F}_{q^2}$, groups ${}^3D_4(q)$ are defined over $\mathbb{F}_{q^3}$, and the remaining groups are defined over $\mathbb{F}_q$. The field $\mathbb{F}_q$ in all cases is called a *base field*. In view of [31, Lemma 2.5.8], if $\overline{G}$ is of adjoint type, then $\overline{G}_\sigma$ is the group of inner-diagonal automorphisms of $O^{p'}(\overline{G}_\sigma)$. If $\overline{G}$ is simply connected, then $\overline{G}_\sigma = O^{p'}(\overline{G}_\sigma)$ (see [42, 12.4]). In any case, in view of [31, Theorem 2.2.6(g)] $\overline{G}_\sigma = \overline{T}_\sigma O^{p'}(\overline{G}_\sigma)$ for every σ-stable maximal torus $\overline{T}$ of $\overline{G}$. Let $U \leq \langle X_r | r \in \Phi^+\rangle = \overline{U}$ be a maximal unipotent subgroup of G (at that $\overline{U}$ is a maximal connected σ-stable unipotent subgroup of $\overline{G}$). Then each $u \in U$ can be uniquely written in form (1), where elements t_r-s are in the definition field of G. If $O^{p'}(G)$ coinsides with one of the groups ${}^2A_n(q)$, ${}^2B_2(2^{2n+1})$, ${}^2D_n(q)$, ${}^3D_4(q)$, ${}^2E_6(q)$, ${}^2G_2(3^{2n+1})$, or ${}^2F_4(2^{2n+1})$, then we shall say that G is *twisted*, in the remaining cases G is called *split*. If $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$ is a twisted group of Lie type and $r \in \Phi(\overline{G})$, then by $\bar{r}$ we always denote the image of a root r under the symmetry of the root system, corresponding to the graph automorphism used during the construction of G. Sometimes we shall use the notation $\Phi^\varepsilon(q)$, where $\varepsilon \in \{+,-\}$, and $\Phi^+(q) = \Phi(q)$ is a split group of Lie type with the base field $\mathbb{F}_q$, $\Phi^-(q) = {}^2\Phi(q)$ is a twisted group of Lie type defined over the field $\mathbb{F}_{q^2}$ (with the base field $\mathbb{F}_q$).

Let $\overline{R}$ be a connected σ-stable subgroup of $\overline{G}$. Then we may consider $R = G \cap \overline{R}$ and $N(G,R) = G \cap N_{\overline{G}}(\overline{R})$. Note that $N(G,R) \neq N_G(R)$ in general, and $N(G,R)$ is called the *algebraic normalizer* of R. For example, if we consider $G = \mathrm{SL}_n(2)$, then the subgroup of diagonal matrices H of G is trivial, hence $N_G(H) = G$. But $G = (\mathrm{SL}_n(\overline{\mathbb{F}}_2))_\sigma$, where σ is a Frobenius map $\sigma : (a_{i,j}) \mapsto (a_{i,j}^2)$. Then $H = \overline{H}_\sigma$, where $\overline{H}$ is the subgroup of diagonal matrices in $\mathrm{SL}_n(\overline{\mathbb{F}}_2)$. Thus $N(G,H)$ is the group of monomial matrices in G. We use the term "algebraic normalizer" in order to avoid such difficulties and to make our proofs to be universal. A group R is called a *torus* (respectively a *reductive subgroup*, a *parabolic subgroup*, a *maximal torus*, a *reductive subgroup of maximal rank*) if $\overline{R}$ is a torus (respectively a reductive subgroup, a parabolic subgroup, a maximal torus, a reductive subgroup of maximal rank) of $\overline{G}$. A maximal σ-stable torus $\overline{T}$ of $\overline{G}$ such that $\overline{T}_\sigma$ is a Cartan subgroup of $\overline{G}_\sigma$ is called a *maximal split torus* of $\overline{G}$.

Assume that a reductive subgroup $\overline{R}$ is σ-stable. In view of [42, 10.10] there exists a σ-stable maximal torus $\overline{T}$ of $\overline{R}$. Let $\overline{G}_{i_1}, \ldots, \overline{G}_{i_{j_i}}$ be a σ-orbit of $\overline{G}_{i_1}$. Consider the induced action of σ on the factor group

$$(\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}})/Z(\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}}) \simeq \mathbf{P}\overline{G}_{i_1} \times \ldots \times \mathbf{P}\overline{G}_{i_{j_i}}.$$

Since $\mathbf{P}\overline{G}_{i_1} \simeq \ldots \simeq \mathbf{P}\overline{G}_{i_{j_i}}$ are simple (as abstract groups), then σ induces a cyclic permutation of the set $\{\mathbf{P}\overline{G}_{i_1}, \ldots, \mathbf{P}\overline{G}_{i_{j_i}}\}$, and we may assume that the numberring is chosen so that $\mathbf{P}\overline{G}_{i_1}^{\sigma} =$

$\mathbf{P}\overline{G}_{i_2}, \ldots, \mathbf{P}\overline{G}^{\sigma}_{i_{j_i}} = \mathbf{P}\overline{G}_{i_1}$. Thus the equality

$$\begin{aligned}&(\mathbf{P}\overline{G}_{i_1} \times \ldots \times \mathbf{P}\overline{G}_{i_{j_i}})_\sigma = \\ &\qquad \{x \mid x = g \cdot g^{\sigma} \cdot \ldots \cdot g^{\sigma^{j_i-1}} \text{ for some } g \in \mathbf{P}\overline{G}_{i_1}\}_\sigma \simeq (\mathbf{P}\overline{G}_{i_1})_{\sigma^{j_i}}\end{aligned}$$

holds. In view of [42, 10.15] the group $\mathbf{P}\overline{G}_{\sigma^{j_i}}$ is finite, hence $O^{p'}((\mathbf{P}\overline{G}_{i_1})_{\sigma^{j_i}})$ is a canonical finite group of Lie type, probably with the base field larger than the base field of $O^{p'}(\overline{G}_\sigma)$.

Let $\overline{B}_{i_1}$ be the preimage of a σ^{j_i}-stable Borel subgroup of $\mathbf{P}\overline{G}_{i_1}$ in $\overline{G}_{i_1}$ under the natural epimorphism, and $\overline{T}_{i_1}$ be a σ^{j_i}-stable maximal torus of $\overline{G}_{i_1}$, contained in $\overline{B}_{i_1}$ (their existence follows from [42, 10.10]). Then from the note at the beginning of section 11 from [42], subgroups $\overline{U}_{i_1}$ and $\overline{U}^-_{i_1}$, generated by $\overline{T}_{i_1}$-invariant root subgroups, taken over all positive and negative roots respectively, are also σ^{j_i}-stable. Since $\overline{G}_{i_1}$ is a simple algebraic group, then $\overline{G}_{i_1}$ is generated by subgroups $\overline{U}_{i_1}$ and $\overline{U}^-_{i_1}$. Now $Z(\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}})$ consists of semisimple elements, so the restriction of the natural epimorphism $\overline{G}_{i_1} \rightarrow \mathbf{P}\overline{G}_{i_1}$ on $\overline{U}_{i_1}$ and $\overline{U}^-_{i_1}$ is an isomorphism. Therefore, for each k the subgroups $(\overline{U}_{i_1})^{\sigma^k}$ and $(\overline{U}^-_{i_1})^{\sigma^k}$ are maximal σ^{j_i}-stab-le connected unipotent subgroups of $\overline{G}_{i_k}$ and they generate $\overline{G}_{i_k}$.

Thus, $\overline{U}_{i_1} \times (\overline{U}_{i_1})^\sigma \times \ldots \times (\overline{U}_{i_1})^{\sigma^{j_i-1}}$ and $\overline{U}^-_{i_1} \times (\overline{U}^-_{i_1})^\sigma \times \ldots \times (\overline{U}^-_{i_1})^{\sigma^{j_i-1}}$ are maximal σ-stable connected unipotent subgroups of $\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}}$ and they generate $\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}}$. By [42, Corollary 12.3(a)], we have

$$\begin{aligned}&O^{p'}((\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}})_\sigma) = \\ &\qquad \langle (\overline{U}_{i_1} \times (\overline{U}_{i_1})^\sigma \times \ldots \times (\overline{U}_{i_1})^{\sigma^{j_i-1}})_\sigma, (\overline{U}^-_{i_1} \times (\overline{U}^-_{i_1})^\sigma \times \ldots \times (\overline{U}^-_{i_1})^{\sigma^{j_i-1}})_\sigma \rangle \simeq \\ &\qquad\qquad \langle (\overline{U}_{i_1})_{\sigma^{j_i}}, (\overline{U}^-_{i_1})_{\sigma^{j_i}} \rangle = O^{p'}((\overline{G}_{i_1})_{\sigma^{j_i}}).\end{aligned}$$

By [42, 11.6 and Corollary 12.3], the group $\langle (\overline{U}_{i_1})_{\sigma^{j_i}}, (\overline{U}^-_{i_1})_{\sigma^{j_i}} \rangle$ is a canonical finite group of Lie type. More over, from the above arguments it follows that the groups $\langle (\overline{U}_{i_1})_{\sigma^{j_i}}, (\overline{U}^-_{i_1})_{\sigma^{j_i}} \rangle / Z(\langle (\overline{U}_{i_1})_{\sigma^{j_i}}, (\overline{U}^-_{i_1})_{\sigma^{j_i}} \rangle)$ and $O^{p'}((\mathbf{P}\overline{G}_{i_1})_{\sigma^{j_i}})$ are isomorphic. Denoting $O^{p'}((\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}})_\sigma)$ by G_i, we obtain that G_i is a canonical finite group of Lie type for all i. Subgroups G_i-s of $O^{p'}(\overline{G}_\sigma)$, appearing in this way, are called *subsystem subgroups* of $O^{p'}(\overline{G}_\sigma)$.

Since $\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}}$ is a σ-stable subgroup, then $\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}} \cap \overline{T}$ is a σ-stable maximal torus of $\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}}$. Therefore we may assume that for each σ-orbit $\{\overline{G}_{i_1}, \ldots, \overline{G}_{i_{j_i}}\}$, the intersection $\overline{T} \cap \overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}}$ is a maximal σ-stable torus of $\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}}$. Then $\overline{R}_\sigma = \overline{T}_\sigma (G_1 \circ \ldots \circ G_m)$ and $\overline{T}_\sigma$ normalizes each of G_i-s.

For a σ-orbit $\{\overline{G}_{i_1}, \ldots, \overline{G}_{i_{j_i}}\}$ of $\overline{G}_{i_1}$, where $G_i = O^{p'}((\overline{G}_{i_1} \circ \ldots \circ \overline{G}_{i_{j_i}})_\sigma)$, consider $\mathrm{Aut}_{\overline{R}_\sigma}(G_i)$. Since $G_1 \circ \ldots \circ G_{i-1} \circ G_{i+1} \circ \ldots \circ G_k \circ \overline{Z}_\sigma \leq C_{\overline{R}_\sigma}(G_i)$, we have that $\mathrm{Aut}_{\overline{R}_\sigma}(G_i) \simeq \left(\overline{T}_\sigma G_i\right) / Z\left(\overline{T}_\sigma G_i\right)$. From [31, Proposition 2.6.2] it follows that automorphisms induced by $\overline{T}_\sigma$ on G_i, are diagonal. Therefore, the inclusions $\mathbf{P}G_i \leq \mathrm{Aut}_{\overline{R}_\sigma}(G_i) \leq \widehat{\mathbf{P}G_i}$ hold, in particular, $\mathrm{Aut}_{\overline{R}_\sigma}(G_i)$ is a finite group of Lie type.

Now consider the case, when $\overline{L} \trianglelefteq \overline{H} \leq \overline{G}$, where $\overline{L}$ and $\overline{H}$ are σ-stable and closed. Clearly σ induces an action on $\overline{H}/\overline{L}$ and, if $\overline{L}$ is connected, then Lang-Steinberg Theorem (Lemma 1.5.3) implies $(\overline{H}/\overline{L})_\sigma = \overline{H}_\sigma / \overline{L}_\sigma$. Let $\overline{R}$ be a σ-stable connected reductive subgroup of maximal rank (in particular, $\overline{R}$ can be a maximal torus) of G. Since groups $N_{\overline{G}}(\overline{R})/\overline{R}$ and $N_W(W_{\overline{R}})/W_{\overline{R}}$ are isomorphic, where W is the Weyl group of $\overline{G}$, $W_{\overline{R}}$ is the Weyl group of $\overline{R}$ (and it is a subgroup of W), we obtain an induced action of σ on $N_W(W_{\overline{R}})/W_{\overline{R}}$, and we say that $w_1 \equiv w_2$, for

$w_1, w_2 \in N_W(W_{\overline{R}})/W_{\overline{R}}$, if there exists an element $w \in N_W(W_{\overline{R}})/W_{\overline{R}}$, satisfying to the equality $w_1 = w^{-1}w_2w^{\sigma}$. Let $Cl(\overline{G}_\sigma, \overline{R})$ be the set of $\overline{G}_\sigma$-conjugated classes of σ-stable subgroups $\overline{R}^g$, where $g \in \overline{G}$. Then $Cl(\overline{G}_\sigma, \overline{R})$ is in 1-1 correspondence with the set of σ-conjugate classes $Cl(N_W(W_{\overline{R}})/W_{\overline{R}}, \sigma)$. If w is an element of $N_W(W_{\overline{R}})/W_{\overline{R}}$, and $(\overline{R}^g)_\sigma$ corresponds to the σ-conjugate class of w, then $(\overline{R}^g)_\sigma$ is said to be obtained by *twisting* of the group $\overline{R}$ by the element $w\sigma$. Further $(\overline{R}^g)_\sigma \simeq \overline{R}_{\sigma w}$. The construction of twisting is known and is given, for example, in [17] with all necessary results. When $\overline{H} = \overline{T}$ is a σ-stable maximal torus and $W = N_G(T)/T$, then by [19, Proposition 3.3.6],

$$\left(\frac{N_G(T_w)}{T_w}\right)_\sigma = \frac{(N_G(T_w))_\sigma}{(T_w)_\sigma} \simeq C_{W,\sigma}(w) = \{x \in W \mid \sigma(x)wx^{-1} = w\}. \tag{2}$$

Now assume that the group $\overline{R}$ is a σ-stable parabolic subgroup of $\overline{G}$ and $\overline{U}$ is its unipotent radical. Then it contains a connected reductive subgroup $\overline{L}$ such that $\overline{R}/\overline{U} \simeq \overline{L}$. A subgroup $\overline{L}$ is called a *Levi factor* of $\overline{R}$. More over, if $\overline{Z} = Z(\overline{L})^0$, then $\overline{L} = C_{\overline{G}}(\overline{Z})$ (see [12, 30.2]). Let $R(\overline{R})$ be the radical of $\overline{R}$. Then it is a σ-stable connected solvable subgroup, hence by [42, 10.10], it contains a σ-stable maximal torus $\overline{Z}$. Now $C_{\overline{G}}(\overline{Z}) = C_{\overline{R}}(\overline{Z})$ is a σ-stable Levi factor of $\overline{R}$. Thus each σ-stable parabolic subgroup of $\overline{G}$ contains a σ-stable Levi factor $\overline{L}$ and $\overline{L}$ is a connected reductive subgroup of maximal rank of $\overline{G}$.

1.5 Known results

In this section we recall some structure results that will be often used below.

Lemma 1.5.1. [34, Theorem 2.2] *Let $\overline{G}$ be a connected reductive algebraic group, $s \in \overline{G}$ be a semisimple element of $\overline{G}$ and $\overline{T}$ be a maximal torus of $\overline{G}$, containing s.*
Then $C_{\overline{G}}(s)^0$ is a reductive subgroup of maximal rank of $\overline{G}$. The centralizer $C_G(s)$ is generated by a torus $\overline{T}$, those $\overline{T}$-root subgroups X_r, for which $s^r = e$ and representatives n_w of elements $w \in W$, which commute with s. Further $C_{\overline{G}}(s)^0$ is generated by the torus $\overline{T}$, those $\overline{T}$-root subgroups X_r, for which $s^r = e$, and each unipotent element centralizing s, is in $C_{\overline{G}}(s)^0$.

Lemma 1.5.2. [34, Proposition 2.10] *Let $\overline{G}$ be a simple algebraic group and s be its semisimple element of finite order.*
Then the factor group $C_{\overline{G}}(s)/C_{\overline{G}}(s)^0$ is isomorphic to a subgroup of the fundamental group $\Delta(\overline{G})$. In particular, if $\overline{G}$ is simply connected, then $C_{\overline{G}}(s)$ is connected.

Lemma 1.5.3. [42, Theorem 10.1] *Let $\overline{G}$ be a connected algebraic group and σ be a Frobenius map.*
Then the map $x \mapsto x^{-1}x^{\sigma}$ is surjective.

The following lemma is known as Borel-Tits theorem.

Lemma 1.5.4. *Let X be a subgroup of a finite group of Lie type G such that $O_p(X)$ is nontrivial. Then there exists a σ-stable parabolic subgroup $\overline{P}$ of $\overline{G}$ such that $X \leq \overline{P}$ and $O_p(X) \leq R_u(\overline{P})$.*

Proof. Define $U_0 = O_p(X)$, $N_0 = N_{\overline{G}}(U_0)$. Then $U_i = U \cdot R_u(N_{i-1})$ and $N_i = N_{\overline{G}}(U_i)$. Clearly U_i, N_i are σ-stable for all i. In view of [12, Proposition 30.3], the chain of subgroups $N_0 \leq N_1 \leq \ldots \leq N_k \leq \ldots$ is finite and $\overline{P} = \cup_i N_i$ is a proper parabolic subgroup. Clearly $\overline{P}$ is σ-stable. □

Lemma 1.5.5. (Hartley-Shute lemma [33, Lemma 2.2]) *Let G be a finite canonical adjoint group of Lie type with the definition field $\mathbb{F}_q$. Let H be a Cartan subgroup of G and $s \in \mathbb{F}_q$. If G is twisted and $r = \bar{r}$, then assume also that s is in the base field of G.*
Then there exists an element $h(\chi) \in H$ such that $\chi(r) = s$, except the following cases, when $h(\chi)$ can be chosen so that $\chi(r)$ would have indicated values:

(a) *$G = A_1(q)$, $\chi(r) = s^2$;*

(b) *$G = C_n(q)$, r is a long root, $\chi(r) = s^2$;*

(c) *$G = {}^2A_2(q)$, $r \neq \bar{r}$, $\chi(r) = s^3$;*

(d) *$G = {}^2A_3(q)$, $r \neq \bar{r}$, $\chi(r) = s^2$;*

(e) *$G = {}^2D_n(q)$, $r \neq \bar{r}$, $\chi(r) = s^2$;*

(f) *$G = {}^2G_2(3^{2n+1})$, $r = a$ or $r = 3a + b$, where a is a short, b is a long fundamental roots, $\chi(r) = s^2$.*

Theorem 1.5.6. [37, Theorem 1.1] *Let $q = p^\alpha$, where p is a prime, and assume $G = \mathrm{Sp}_n(q)$, or $\mathrm{SO}^\varepsilon_n(q) \leq G \leq \mathrm{GO}^\varepsilon_n(q)$, where q is odd, or $\mathrm{SU}_n(q) \leq G \leq \mathrm{GU}_n(q)$. If G admits a a Carter subgroup K, then either K is the normalizer of a Sylow 2-subgroup of G, or one of the following holds:*

(a) *$G \in \{\mathrm{Sp}_2(3),\ \mathrm{SL}_2(3),\ 2.\mathrm{SU}_2(3)\}$ and K is the normalizer of a Sylow 3-subgroup of G;*

(b) *$G = \mathrm{GU}_3(2)$ has order $2^3 \cdot 3^4$, and K has order $2 \cdot 3^2$.*

Moreover, if G is orthogonal, K is a 2-group, except possibly when $G = \mathrm{SO}^\varepsilon_2(q)$.

2 Conjugacy criterion for Carter subgroups

2.1 Main results of this section

Definition 2.1.1. A finite group G is said to satisfy condition **(C)**, if, for *every* non-Abelian composition factor S of *every* composition series of G and for every its nilpotent subgroup N, Carter subgroups of $\langle \mathrm{Aut}_N(S), S\rangle$ are conjugate (in particular, they may not exist).

Lemma 2.1.2. *Let H be a normal subgroup of a finite group G, $B \lhd A \leq G$ and $S = (A/H)/(B/H)$ is a composition factor of G/H, and $L \leq G$.*
Then $\mathrm{Aut}_L(A/B) \simeq \mathrm{Aut}_{LH/H}((A/H)/(B/H))$.

Proof. Since $H \leq B$, then $H \leq C_G(A/B)$, so we may assume that $L = LH$. Further more we may assume that $L \leq N_G(A) \cap N_G(B)$ and $G = LA$. Then the action on A/B given by $x : Ba \mapsto Bx^{-1}ax$ coincides with the action on $(A/H)/(B/H)$ given by $xH : BaH \mapsto Bx^{-1}axH$, and the lemma follows. □

The following lemma is known.

Lemma 2.1.3. *Let G be a finite group, H be a normal subgroup of G and $\overline{N}$ be a nilpotent subgroup of $\overline{G} = G/H$.*
Then there exists a nilpotent subgroup N of G such that $NH/H = \overline{N}$.

Proof. Clearly we may assume that $G/H = \overline{N}$. There exists a subgroup U of G such that $UH = G$. Choose a subgroup of minimal order with this property. Then $U \cap H$ is contained in the Frattini subgroup F of U. Indeed, if there exists a maximal subgroup M of U, not containing $U \cap H$, then clearly $MH = G$, which contradicts the minimality of U. Thus the group U/F is nilpotent, hence U is nilpotent and $N = U$. □

By Lemmas 2.1.2 and 2.1.3 it follows that, if a finite group G satisfies $(\mathbf{C})$, then for every its normal subgroup N and solvable subgroup H, groups G/N and HN satisfy $(\mathbf{C})$.

In this section we prove that if G satisfies $(\mathbf{C})$, then its Carter subgroups are conjugate. More precisely, the following theorem will be proven.

Theorem 2.1.4. *If a finite group G satisfies $(\mathbf{C})$, then Carter subgroups of G are conjugate.*

Below in subsections 2.2, 2.3 we are assuming that X is a counter example to Theorem 2.1.4 of minimal order, i. e., that X is a finite group satisfying $(\mathbf{C})$, and X contains nonconjugate Carter subgroups, but Carter subgroups in every group M of order less than $|X|$, satisfying $(\mathbf{C})$, are conjugate.

2.2 Preliminary results

Recall that X is a counter example to Theorem 2.1.4 of minimal order.

Lemma 2.2.1. *Let G be a finite group satisfying $(\mathbf{C})$, $|G| \leqslant |X|$, and H be a Carter subgroup of G.*
If N is a normal subgroup of G, then HN/N is a Carter subgroup of G/N.

Proof. Since HN/N is nilpotent, we have just to prove that it is self-normalizing in G/N. Clearly, this is true if $G = HN$. So, assume $M = HN < G$ (note that by Lemmas 2.1.2 and 2.1.3 the group M satisfies $(\mathbf{C})$). By the minimality of X, $M^x = M$, $x \in G$, implies $H^x = H^m$ for some $m \in M$. It follows $xm^{-1} \in N_G(H) = H$ and $x \in M$. This proves that HN/N is nilpotent and self-normalizing in G/N. □

Lemma 2.2.2. *Let B be a minimal normal subgroup of X and H, K be non-conjugate Carter subgroups of X.*

(1) *B is non-soluble.*

(2) *$X = BH = BK$.*

(3) *B is the unique minimal normal subgroup of X.*

Proof. (1) We give a proof by contradiction. Assume that B is soluble and let $\pi : X \to X/B$ be the canonical homomorphism. Then H^π and K^π are Carter subgroups of X/B, by Lemma 2.2.1. By the minimality of X, there exists $\bar{x} = Bx$ such that $(K^\pi)^{\bar{x}} = H^\pi$. It follows $K^x \leq BH$. Since BH is soluble, K^x is conjugate to H in BH, hence K is conjugate to H in X, a contradiction.

(2) Assume that $BH < X$. By Lemma 2.2.1 and the minimality of X, BH/B and BK/B are conjugate in X/B: so there exists $x \in X$ such that $K^x \leq BH$. It follows that K^x is conjugate to H in BH, hence K is conjugate to H in X, a contradiction.

(3) Suppose that M is a minimal normal subgroup of X different from B. By (1), M is non-soluble. On the other hand, $MB/B \simeq M$ is a subgroup of the nilpotent group $X/B \simeq H/H \cap B$, a contradiction. □

Lemma 2.2.3. *Let K be a Carter subgroup of a finite group G. Assume that there exists a normal subgroup $B = T_1 \times \ldots \times T_k$ of G such that $G = KB$, $Z(T_i) = \{e\}$, and T_i is not decomposable into direct product of its proper subgroups for all i.*
Then $\mathrm{Aut}_K(T_i)$ is a Carter subgroup of $\langle \mathrm{Aut}_K(T_i), T_i \rangle$.

Proof. Assume that our statement is false and G is a counterexample with k minimal, then $k > 1$. Since each group T_i has trivial center and is not decomposable into direct product of proper subgroups, a corollary of Krull-Remak-Shmidt theorem [38, 3.3.10] implies that the action by conjugation of G on the set $\{T_1, \ldots, T_k\}$ induces permutations of this set. Clearly, G acts transitively, by conjugation, on the set $\Omega := \{T_1, \ldots, T_k\}$. We may assume that the T_j-s are indexed so that G acts primitively on the set $\{\Delta_1, \ldots, \Delta_p\}$, $p > 1$, where for each i:

$$\Delta_i := \{T_{1+(i-1)l}, \ldots, T_{il}\}, \qquad k = pl.$$

Denote by $\varphi : G \to \mathrm{Sym}_p$ the induced permutation representation. Clearly, $B \leq \ker \varphi$, so that $G^{\varphi} = (BK)^{\varphi} = K^{\varphi}$ is a primitive nilpotent subgroup of Sym_p. Hence p is prime and G^{φ} is a cyclic group of order p. In particular, $Y := \mathit{ker}\ \varphi$ coincides with the stabilizer of any Δ_i, so that φ is permutationally equivalent to the representation of G on the right cosets of Y. For each $i = 1, \ldots, p$, let $S_i = T_{1+(i-1)l} \times \ldots \times T_{il}$. Then $Y = N_G(S_i)$ and $B = S_1 \times \ldots \times S_p$. Consider $\xi : Y \to \mathrm{Aut}_Y(S_1)$, let $A = Y^{\xi}$, $S = S_1^{\xi}$. Clearly S is a normal subgroup of A; moreover, S is isomorphic to S_1, since S_1 has trivial center. On the other hand, for each $i \neq 1$, $S_i \leq \ker \xi$, since S_i centralizes S_1.

Denote by $A \wr \mathbb{Z}_p$ the wreath product of A and a cyclic group $\mathbb{Z}_p$, and let $\{x_1 = e, \ldots, x_p\}$ be a right transversal of Y. Then the map $\eta : G \to A \wr \mathbb{Z}_p$ such that, for each $x \in G$:

$$x \mapsto \left(\left(x_1 x x_{1^{x^{\varphi}}}^{-1}\right)^{\xi}, \ldots, \left(x_p x x_{p^{x^{\varphi}}}^{-1}\right)^{\xi} \right) x^{\varphi}$$

is a homomorphism. Clearly Y^{η} is a subdirect product of the base subgroup A^p and

$$S_1^{\eta} = \{(s, 1, \ldots, 1) | s \in S\}, B^{\eta} = \{(s_1, \ldots, s_p) | s_i \in S\} \leq Y^{\eta}.$$

Moreover, $\ker \eta = C_G(B) = \{e\}$, so we may identify G with G^{η}. We choose $h \in K \setminus Y$. Then

$$G = \langle Y, h \rangle,\ h^p \in Y,\ K = (Y \cap K)\langle h \rangle$$

and we may assume

$$h = (a_1, a_2, \ldots, a_p)\pi,\ a_i \in A, \quad \pi = (1, 2, \ldots, p) \in \mathbb{Z}_p.$$

For each i, $1 \leqslant i \leqslant p$, let $\psi_i : A^p \to A$ be the canonical projection and let $K_i := (K \cap Y)^{\psi_i}$. Clearly, $Y^{\psi_i} = A$. Moreover, for each $i \geqslant 2$, $K_i = K_1^{h^{i-1}} = K_1^{a_1 \ldots a_{i-1}}$ since h normalizes $Y \cap K$. Let $N := (K_1 \times \ldots \times K_p) \cap Y$. N is normalized by K, since $K = (N \cap K)\langle h \rangle$ and $K_i^h = K_{i+1 \pmod p}$. We claim that K_1 is a Carter subgroup of A. Assume $n_1 \in N_A(K_1) \setminus K_1$. From $Y = (Y \cap K)B$, it follows $n_1 = h_1 s$, $h_1 \in K_1$, $s \in N_S(K_1) \setminus K_1$. Let $b := (s, s^{a_1}, \ldots, s^{a_1 \ldots a_{p-1}}) \in B$. Then b normalizes N, for:

$$K_i^b = K_i^{s^{a_1 \ldots a_{i-1}}} = K_1^{a_1 \ldots a_{i-1} s^{a_1 \ldots a_{i-1}}} = K_1^{s a_1 \ldots a_{i-1}} = K_1^{a_1 \ldots a_{i-1}} = K_i.$$

Now $[b, h^{-1}] := b^{-1} h b h^{-1} \in Y$ is such that:

$$[b, h^{-1}]^{\psi_i} = 1 \text{ if } i \neq p, [b, h^{-1}]^{\psi_p} = [s, (a_1 \cdot \ldots \cdot a_p)^{-1}]^{a_1 \cdot \ldots \cdot a_{p-1}},$$

where $a_1 \cdot \ldots \cdot a_p = (h^p)^{\psi_1} \in K_1$. Since $s \in N_S(K_1)$, it follows

$$[s, (a_1 \cdot \ldots \cdot a_p)^{-1}] \in K_1, \ [s, (a_1 \cdot \ldots \cdot a_p)^{-1}]^{a_1 \cdot \ldots \cdot a_{p-1}} \in K_p.$$

So $[b, h^{-1}] \in N$ and $b \in N_G(N\langle h\rangle)$. But $K \leq N\langle h\rangle$, implies $N_G(N\langle h\rangle) = N\langle h\rangle$. Indeed, if $g \in N_G(N\langle h\rangle)$, then K^g is a Carter subgroup of $N\langle h\rangle$. But $N\langle h\rangle$ is soluble, hence there exists $y \in N\langle h\rangle$ with $K^g = K^y$. Now K is a Carter subgroup of G, thus $gy^{-1} \in K$ and $g \in N\langle h\rangle$. Therefore $b \in N, s \in K_1$, i. e., $n_1 \in K_1$, a contradiction.

Now $A = K_1(T_1 \times \ldots \times T_l)$ and $l < k$. By induction we have that $\mathrm{Aut}_{K_1}(T_1)$ is a Carter subgroup of $\langle \mathrm{Aut}_{K_1}(T_1), T_1\rangle$. In view of our construction, $\mathrm{Aut}_K(T_1) = \mathrm{Aut}_{K_1}(T_1)$ and the lemma follows. □

2.3 Proof of Theorem 2.1.4

Recall that $B = T_1 \times \cdots \times T_k$, where $T_i \simeq T$ is a non-Abelian simple group. What remains to prove is $k = 1$. Осталось доказать, что $k = 1$. In the notations of the proof of Lemma 2.2.3 we have shown that H_1 is a Carter subgroup of A. If $k > 1$, then $|A| < |X|$ and A satisfies **(C)**. So each K_i is conjugate with K_1 in A and $N_A(K_i) = K_i$, $i = 1, \ldots, p$. It follows easily that N is a Carter subgroup of Y. Let $y := (y_1, \ldots, y_p) \in N_Y(N)$. From $N^{\psi_i} = K_i$ we have $y_i \in N_A(K_i) = K_i$ for each i, hence $y \in N$.

We have seen that, to each Carter subgroup K of X we can associate a Carter subgroup $N = N_K$ of Y, such that K normalizes N_K. Clearly, $N_K \neq \{e\}$, otherwise X would have order p. So let H be a Carter subgroup of X, not conjugate to K, and let N_H be the Carter subgroup of Y corresponding to H. If $k > 1$, then Y is a proper subgroup of X and Y satisfies **(C)**. By the minimality of X we obtain that N_H and N_K are conjugate in Y, and we may assume that $N_H = N_K$. Then $HN_H = KN_H$ is solvable, hence, the subgroups H and K are conjugate. This contradiction completes the proof of Theorem 2.1.4.

2.4 Some properties of Carter subgroups

Here we shall prove some lemmas that will be useful in studying Carter subgroups in finite groups, in particular, in almost simple groups.

Lemma 2.4.1. *Let K be a Carter and N be a normal subgroups of a finite group G. Assume that KN satisfies* **(C)** *(this condition holds if either G satisfies* **(C)** *or N is solvable) or $KN = G$.*
Then KN/N is a Carter subgroup of G/N.

Proof. If $KN = G$, then the statement is evident. Assume that $KN \neq G$, i. e., KN satisfies **(C)**. Consider $x \in G$ and assume that $xN \leq N_{G/N}(KN/N)$. Therefore $x \in N_G(KN)$. We have that K^x is a Carter subgroup of KN. Since KN satisfies **(C)**, we obtain that its Carter subgroups are conjugate. Thus there exists $y \in KN$ such that $K^y = K^x$. Since K is a Carter subgroup of G, it follows that $xy^{-1} \in N_G(K) = K$ and $x \in KN$. □

Lemma 2.4.2. *Let K be a Carter subgroup of a finite group G. Assume also that $e \neq z \in Z(K)$ and $C_G(z)$ satisfies* **(C)**.

(1) *Every subgroup Y which contains K and satisfies* **(C)***, is self-normalizing in G.*

(2) *No conjugate of z in G, except z, lies in $Z(G)$.*

(3) *If H is a Carter subgroup of G, non-conjugate to K, then z is not conjugate to any element in the center of H.*

In particular the centralizer $C_G(z)$ is self-normalizing in G, and z is not conjugate to any power $z^k \neq z$.

Proof. (1) Take $x \in N_G(Y)$. Then K^x is a Carter subgroup of Y. By Theorem 2.1.4 Carter subgroups of Y are conjugate. Therefore there exists $y \in Y$ with $K^x = K^y$. Hence

$$xy^{-1} \in N_G(K) = K \leq Y \text{ and } x \in Y.$$

(2) Assume $z^{x^{-1}} \in Z(K)$ for some $x \in G$. Then z belongs to the center of $\langle G, G^x\rangle \leq C_G(z)$. Since $C_G(z)$ satisfies **(C)**, there exists $y \in C_G(z)$ such that $K^x = K^y$. From $xy^{-1} \in C_G(z)$, we get $z^{xy^{-1}} = z$ hence $z^x = z^y = z$. We conclude $z^{x^{-1}} = z$.

(3) If our claim is false, substituting H with some conjugate H^x (if necessary), we may assume $z \in Z(K) \cap Z(H)$, i. e. $z \in Z(\langle K, H\rangle) \leq C_G(z)$. Again since $C_G(z)$ satisfies **(C)**, there exists $y \in C_G(z)$ such that $H = K^y$; a contradiction. □

Note that for every known finite simple group G (and hence almost simple, since the group of outer automorphisms is soluble) and for all elements $z \in G$ of prime order we see that composition factors of $C_G(z)$ are known simple groups. Indeed, for sporadic groups this statement can be checked by using [21]. Composition factors of $C_{A_n}(z)$ are alternating groups. If G is a finite simple group of Lie type over a field of characteristic p and $(|z|, p) = 1$, then z is semisimple and all composition factors of $C_G(z)$ are finite groups of Lie type. If $|z| = p$ and p is a good prime for G, then by Theorems 1.2 and 1.4 from [40], all composition factors of $C_G(z)$ are finite groups of Lie type. From papers of several authors it follows that in case when p is a bad prime for a finite adjoint group of Lie type G, all composition factors of the centralizer of an element of order p are known finite simple groups. Therefore, if we are classifying Carter subgroups of an almost simple group A, then by induction we may assume that $C_A(z)$ satisfies **(C)** for all elements $z \in A$ of prime order.

Lemma 2.4.3. *Let Q be a Sylow 2-subgroup of a finite group G.*
Then G contains a Carter subgroup K, satisfying $Q \leq K$, if and only if $N_G(Q) = QC_G(Q)$.

Proof. Assume that G contains a Carter subgroup K, satisfying $Q \leq K$. Since K is nilpotent, it follows that Q is normal in K and $K \leq QC_G(Q) \trianglelefteq N_G(Q)$. By Feit-Thompson theorem (see [29]), we obtain that $N_G(Q)$ is solvable. Thus by Lemma 2.4.2(1) we have that $QC_G(Q)$ is self-normalizing in $N_G(Q)$, so $N_G(Q) = QC_G(Q)$.

Assume now that $N_G(Q) = QC_G(Q)$, i. e., the equality $N_G(Q) = Q \times O(C_G(Q))$ holds. Since $O(C_G(S))$ is of odd order, it is solvable. Therefore it contains a Carter subgroup K_1. Consider a nilpotent subgroup $K = Q \times K_1$ of G. Assume that $x \in N_G(K)$, then $x \in N_G(Q)$. But K is a Carter subgroup of $N_G(Q)$, hence $x \in K$ and K is a Carter subgroup of G. □

Definition 2.4.4. A finite group G is said to satisfy **(ESyl2)**, if for its Sylow 2-subgroup Q the equality $N_G(Q) = QC_G(Q)$ holds. In other words, G satisfies **(ESyl2)**, if every element of odd order, normalizing a Sylow 2-subgroup Q of G, centralizes Q.

Lemma 2.4.5. *Let Q be a Sylow 2-subgroup of a finite group G and x be an element of odd order from $N_G(Q)$. Assume that there exist normal subgroups $G_1, \ldots, G_k$ of G such that $G_1 \cap \ldots \cap G_k \cap Q \leq Z(N_G(Q))$ and x centralizes Q modulo G_i for all i.*
Then x centralizes Q. In particular, if G/G_i satisfies **(ESyl2)** *for all i, then G satisfies* **(ESyl2)**.

Proof. Consider the normal series $Q \trianglerighteq Q_1 \trianglerighteq \ldots \trianglerighteq Q_k \trianglerighteq Q_{k+1} = \{e\}$, where $Q_i = Q \cap (G_1 \cap \ldots \cap G_i)$. The conditions of the lemma imply that x centralizes each factor Q_{i-1}/Q_i. Since x is an element of odd order, this implies that x centralizes Q. □

Lemma 2.4.6. *Let H be a subgroup of a finite group G such that $|G : H| = 2^t$, H satisfies* **(ESyl2)**, *and each element of odd order of G is in H (this condition is evidently equivalent to the subnormality of H).*
Then G satisfies **(ESyl2)**.

Proof. Let Q be a Sylow 2-subgroup of G such that $Q \cap H$ is a Sylow 2-subgroup of H. Consider an element $x \in N_G(Q)$ of odd order. Since $x \in H$, then $x \in N_H(Q) \leq N_H(Q \cap H) = (Q \cap H) \times O(N_H(Q \cap H))$, i. e., $x \in O(N_H(Q \cap H))$. Thus the set of elements of odd order in $N_G(Q)$ forms a subgroup $R = O(N_H(Q \cap H)) \cap N_G(Q)$ of $N_G(Q)$. Clearly R is normal in $N_G(Q)$, therefore $R = O(N_G(Q))$. On the other hand, Q is normal in $N_G(Q)$ by definition and $Q \cap R = \{e\}$, whence $N_G(Q) = Q \times O(N_G(Q))$. □

By using the result of this section, we shall improve the definition of minimal counter example.

Definition 2.4.7. A finite almost simple group A is called a *minimal counter example*, if it contains nonconjugate Carter subgroups, but Carter subgroups of every almost simple group, of order less than $|A|$ with simple socle being a *known* simple group, are conjugate.

3 Conjugacy in simple groups

3.1 Brief review of results of the section

Recall that in view of Lemma 2.4.2 none element from the center of a Carter subgroup can be conjugate to its nontrivial power (if the centralizer of the element satisfies **(C)**). Thus if we would be able to prove that each element of prime order r of G is conjugate to its nontrivial power and at the same time its centralizer satisfies **(C)**, then we may state that order of a Carter subgroup (if it exists) is not divisible by r.

In this section we obtain the information on the conjugacy of elements of prime order in finite simple groups and, by using this information, we obtain a description of Carter subgroups in a wide class of almost simple groups. Actually, in almost simple groups, distinct from $A_n^\varepsilon(q)$ ($\varepsilon = \pm$), Carter subgroups should be 2-groups, as made clear below. The results can be formulated as a list of almost simple groups A-s, that cannot be minimal counter example (see Theorem 3.3.5). This list is summarized in the Table 3, where Field(S) stands for the group generated by field and inner-diagonal automorphisms of a finite group of Lie type S.

Table 3: Finite almost simple groups, which are not minimal counter examples

Soc(A)$= G$	Conditions for A
alternating, sporadic; $A_1(p^t)$, $B_\ell(p^t)$, $C_\ell(p^t)$, t is even if $p = 3$; ${}^2B_2(2^{2n+1})$,$G_2(p^t)$, $F_4(p^t)$, ${}^2F_4(2^{2n+1})$; $E_7(p^t)$, $p \neq 3$; $E_8(p^t)$, $p \neq 3, 5$ ${}^3D_4(p^t)$, $D_{2\ell}(p^t)$, ${}^2D_{2\ell}(p^t)$, t is even if $p = 3$ in the last 2 cases and,	none

if $G = D_4(p^t)$, $\lvert(\mathrm{Field}(G) \cap A) : (\widehat{G} \cap A)\rvert_{2'} > 1$	
$B_\ell(3^t)$, $C_\ell(3^t)$, $D_{2\ell}(3^t)$, ${}^3D_4(3^t)$, ${}^2D_{2\ell}(3^t)$, $D_{2\ell+1}(r^t)$, ${}^2D_{2\ell+1}(r^t)$, ${}^2G_2(3^{2n+1})$, $E_6(p^t)$, ${}^2E_6(p^t)$, $E_7(3^t)$, $E_8(3^t)$, $E_8(5^t)$	$A = G$

In particular, A cannot be simple (case $A = A_\ell^\varepsilon(q)$ is excluded by Theorem 1.5.6).

3.2 Preliminary results

Lemma 3.2.1. *Let $\overline{G}$ be a simple connected algebraic group over a field of characteristic p, t be an element of order r of $\overline{G}$, not divisible by p.*
Then $C_{\overline{G}}(t)/C_{\overline{G}}(t)^0$ is a $\pi(r)$-group.

Proof. Since p does not divide r, then t is semisimple. By Lemma 1.5.1, $C_{\overline{G}}(t)^0$ is a connected reductive subgroup of maximal rank of $\overline{G}$ and every p-element of $C_{\overline{G}}(t)$ is contained in $C_{\overline{G}}(t)^0$. Assume that a prime $s \notin \pi(r)$ divides order $|C_{\overline{G}}(t)/(C_{\overline{G}}(t)^0)|$. Then $s \neq p$ and $C_{\overline{G}}(t)$ contains an element x of order s^k such that $x \notin C_{\overline{G}}(t)^0$. Since x, t commute, we have that $x \cdot t$ is a semisimple element of $\overline{G}$ (of order $r \cdots^k$). Therefore, there exists a maximal torus $\overline{T}$ of $\overline{G}$ containing $x \cdot t$. Then $(xt)^r = x^r \in \overline{T}$. Since $(s, r) = 1$, there exists m such that $rm \equiv 1 \pmod{s^k}$, thus $(x^r)^m = x \in \overline{T}$. Since $xt, x \in \overline{T}$, then $t \in \overline{T}$, so $\overline{T} \leq C_{\overline{G}}(t)^0$, hence $x \in C_{\overline{G}}(t)^0$; a contradiction. ☐

Lemma 3.2.2. *Let $s \in \overline{G}$ be a semisimple element of order r such that $(r, \Delta(\overline{G})) = 1$.*
Then $C_{\overline{G}}(s)$ is connected. In particular, it follows that for every Frobenius map σ of $\overline{G}$, two semisimple elements $s, s' \in \overline{G}_\sigma$ are conjugate in $\overline{G}_\sigma$ if and only if they are conjugate in $\overline{G}$.

Proof. Follows from Lemmas 1.5.2 and 3.2.1. ☐

The following lemma plays an important role, since it shows that a semisimple element of odd prime order is usually conjugate to its inverse.

Lemma 3.2.3. *Let $G = O^{r'}(\overline{G}_\sigma)$, $\overline{G}$ has an adjoint type and the root system of $\overline{G}$ has type distinct from A_ℓ $(\ell > 1), D_{2\ell+1}, E_6$.*
Then each semisimple element of odd order $s \in \widehat{G}$ is conjugate to its inverse by an element of G.

Proof. There exists some σ-stable maximal torus $\overline{T}$ of $\overline{G}$ with $s \in \overline{T}$. $\overline{T}$ is generated by the set $\{h_\alpha(\lambda) \mid \alpha \in \Phi, \lambda \in \overline{\mathbb{F}}_p^*\}$ and the factor group $N_{\overline{G}}(\overline{T})/\overline{T}$ is isomorphic to the Weyl group W of $\overline{G}$. If $w \in W$ and n_w is a preimage of w under the natural epimorphism $N_{\overline{G}}(\overline{T}) \to W$, then $h_\alpha(\lambda)^{n_w} = h_{\alpha^w}(\lambda)$. Now let w_0 be the unique involution of W such that $w_0(\Phi^+) = \Phi^-$ and let n_0 be a preimage of w_0. Since we are assuming $\Phi \neq A_\ell$ $(\ell > 1)$, $D_{2\ell+1}$, and E_6, we have $\alpha^{w_0} = -\alpha$ for all $\alpha \in \Phi$, hence $h_\alpha(\lambda)^{n_0} = h_{-\alpha}(\lambda) = h_\alpha(\lambda)^{-1}$. We conclude that $s^{n_0} = s^{-1}$, i.e. that s is conjugate to s^{-1} in $\overline{G}$. Thus, by the previous Lemma, s and s^{-1} are conjugate in $\overline{G}_\sigma$. Finally, from $\overline{G}_\sigma = \overline{T}_\sigma G$, we conclude that s and s^{-1} are conjugate in G. ☐

Lemma 3.2.4. *Let $\overline{C}$ be a connected reductive subgroup of maximal rank of $\overline{G}$. Denote by W and $W_{\overline{C}}$ the Weyl groups of $\overline{G}$ and $\overline{C}$ respectively, by $W_{\overline{C}}^\perp$ the subgroup of W, generated by reflections in roots orthogonal to all roots from $\Phi(\overline{C})$, and by $\Delta_{\overline{C}}$ the Dynkin diagram of $\overline{C}$. Then:*

(a) $N_W(W_{\overline{C}})/(W_{\overline{C}} \times W_{\overline{C}}^{\perp}) \simeq \mathrm{Aut}_W(\Delta_{\overline{C}})$;

(b) $N_{\overline{G}}(\overline{C})/\overline{C} \simeq N_W(W_{\overline{C}})/W_{\overline{C}}$.

Let $G = O^{p'}(\overline{G}_\sigma)$ *be split or one of the groups* ${}^2A_\ell(p^t)$, ${}^2D_{2\ell+1}(p^t)$, ${}^2E_6(p^t)$. *If* $s \in G$ *is a semisimple element such that* $C_{\overline{G}}(s)$ *is connected and* $N_G(C_{\overline{G}}(s)) > C_{\overline{G}}(s)$, *then* $N_G(C_G(s)) > C_G(s)$.

Proof. Point (a) may be found in [17, Proposition 4]. As to point (б), let $\overline{T}$ be a maximal torus of $\overline{G}$ contained in $\overline{C}$, so that we may assume $W = N_{\overline{G}}(\overline{T})/\overline{T}$ and $W_{\overline{C}} = N_{\overline{C}}(\overline{T})/\overline{T}$. All maximal tori of $\overline{C}$ are conjugate in $\overline{C}$, since $\overline{C}$ is connected. It follows easily that $N_{\overline{G}}(\overline{C}) = \overline{C} N_{N_{\overline{G}}(\overline{T})}(\overline{C})$. Moreover it is shown in [17, Proposition 5] that $N_{N_{\overline{G}}(\overline{T})}(\overline{C}) = N_{N_{\overline{G}}(\overline{T})}(N_{\overline{C}}(\overline{T}))$. Hence,

$$\frac{N_{\overline{G}}(\overline{C})}{\overline{C}} = \frac{\overline{C} N_{N_{\overline{G}}(\overline{T})}(N_{\overline{C}}(\overline{T}))}{\overline{C}} \simeq \frac{N_{N_{\overline{G}}(\overline{T})(N_{\overline{C}}(\overline{T}))}}{N_{\overline{C}}(\overline{T})} \simeq \frac{N_W(W_{\overline{C}})}{W_{\overline{C}}}. \tag{3}$$

Now, let $G = O^{p'}(\overline{G}_\sigma)$ be as in the statement, and set $\overline{C} = C_{\overline{G}}(s)$. Write $\sigma = \tau\varphi$, where τ is the graph automorphism of $\overline{G}$ induced by a symmetry ρ of the Dynkin diagram of $\Phi = \Phi(\overline{G})$ and φ is a field automorphism. Now let τ be the isometry which extends ρ on Euclidean space $\mathbb{R} \otimes_{\mathbb{Z}} \mathbb{Z}\Phi$. If $\overline{T}_1$ is a σ-stable maximal split torus of $\overline{G}$, then for each $x \in N_{\overline{G}}(\overline{T}_1)/\overline{T}_1$, we have $x^\sigma = {}^\tau x$ (considering $N_{\overline{G}}(\overline{T}_1)/\overline{T}_1 = W_1$ as a group of isometries of $\mathbb{R} \otimes_{\mathbb{Z}} \mathbb{Z}\Phi$). Thus if G is split, i. e., $\rho = \tau = e$, then σ acts trivially on W_1. If G is twisted, hence of type A_ℓ, $D_{2\ell+1}$, or E_6, it is possible to show directly, that $-\tau \in W_1$. Thus we may twist $\overline{T}_1$ by $-\tau$, obtaining the σ-stable torus $(\overline{T}_1)_{-\tau}$. By equation (2):

$$\frac{(N_{\overline{G}}((\overline{T}_1)_{-\tau})))_\sigma}{((\overline{T}_1)_{-\tau}))_\sigma} \simeq C_{W_1,\sigma}(-\tau) = \{x \in W_1 \mid {}^\tau x(-\tau)x^{-1} = -\tau\} = W_1.$$

Let $\{X_\alpha \mid \alpha \in \Phi\}$ be the set of $\overline{T}_1$-root subgroups and set $\overline{C}_1 = \langle \overline{T}_1, X_\alpha \mid \alpha \in \Phi(\overline{C}) \rangle$. Since $\Phi(\overline{C})$ is σ-invariant it follows that $\overline{C}_1$ is σ-stable. Moreover, since $\tau(\Phi(\overline{C})) = \Phi(\overline{C})$, we have that $-\tau \in N_{W_1}(W_{\overline{C}_1})$. By [17, Proposition 1 and 2], it follows that there exists $(\overline{C}_1)_{-\tau}$ obtained from $\overline{C}_1$ by twisting with $-\tau$. Up to conjugation in G we may assume that $(\overline{T}_1)_{-\tau} \leq (\overline{C}_1)_{-\tau}$. Define $\overline{T}_0 = \overline{T}_1$ and $\overline{C}_0 = \overline{C}_1$ if G is split, and $\overline{T}_0 = (\overline{T}_1)_{-\tau}$ and $\overline{C}_0 = (\overline{C}_1)_{-\tau}$ if G is twisted.

Since $\Phi(\overline{C}) = \Phi(\overline{C}_0)$, there exists $g \in \overline{G}$, such that ${}^g\overline{C}_0 = C$ and ${}^g\overline{T}_0 = \overline{T}$. It follows that $\dot{w} = g^{-1}\sigma(g) \in N_{\overline{G}}(\overline{C}_0) \cap N_{\overline{G}}(\overline{T}_0)$. So the image w of $\dot{w}$ in $W_0 = N_{\overline{G}}(\overline{T}_0)/\overline{T}_0$ belongs to $N_{W_0}(W_{C_0})$.

From $\overline{G}_\sigma = \overline{T}_\sigma G$ it follows $(N_{\overline{G}}(\overline{C}))_\sigma = N_{\overline{G}_\sigma}(\overline{C}) = \overline{T}_\sigma N_G(\overline{C})$. Hence we are done if we can show that it is nontrivial the group $\frac{(N_{\overline{G}}(\overline{C}))_\sigma}{\overline{C}_\sigma} = \frac{\overline{T}_\sigma N_G(\overline{C})}{\overline{T}_\sigma C_G(s)} \simeq \frac{N_G(\overline{C})}{C_G(s)}$, which is a subgroup of $\frac{N_G(C_G(s))}{C_G(s)}$.

Using equation (3) we get

$$\frac{(N_{\overline{G}}(\overline{C}))_\sigma}{\overline{C}_\sigma} \simeq \frac{(N_{N_{\overline{G}}(\overline{T})}(N_{\overline{C}}(\overline{T}))/\overline{T})_\sigma}{(N_{\overline{C}}(\overline{T})/\overline{T})_\sigma} \simeq \frac{N_{N_{\overline{G}}(\overline{T})}(N_{\overline{C}}(\overline{T}))/\overline{T} \cap (N_{\overline{G}}(\overline{T})/\overline{T})_\sigma}{N_{\overline{C}}(\overline{T})/\overline{T} \cap (N_{\overline{G}}(\overline{T})/\overline{T})_\sigma}.$$

By our choice of $\overline{T}_0$ we have $\frac{N_{\overline{G}}(\overline{T}_0)}{\overline{T}_0} = \left(\frac{N_{\overline{G}}(\overline{T}_0)}{\overline{T}_0}\right)_\sigma$, i. e. σ acts trivially on the finite group $\frac{N_{\overline{G}}(\overline{T}_0)}{\overline{T}_0}$. Now, if $w \in W_{\overline{C}_0}$, by [17, Proposition 1] we may assume $w = e$, $\overline{T} = \overline{T}_0$, $\overline{C} = \overline{C}_0$. It follows

$N_{\overline{G}}(\overline{T})/\overline{T} = (N_{\overline{G}}(\overline{T})/\overline{T})_\sigma$, hence

$$\frac{(N_{\overline{G}}(\overline{C}))_\sigma}{\overline{C}_\sigma} \simeq \frac{N_W(W_{\overline{C}})}{W_{\overline{C}}} \simeq \frac{N_{\overline{G}}(\overline{C})}{\overline{C}}$$

which is non-trivial by assumption. Finally assume that $w \notin W_{\overline{C}_0}$, i. e. $\dot{w} = g^{-1}\sigma(g) \notin \overline{C}_0$. It follows that ${}^g\dot{w} = \sigma(g)g^{-1} \notin \overline{C}$, i. e. ${}^g\dot{w}\overline{T} \notin N_{\overline{C}}(\overline{T})/\overline{T}$. On the other hand ${}^g\dot{w} \in N_{\overline{G}}(\overline{C}) \cap N_{\overline{G}}(\overline{T})$. Moreover, since σ acts trivially on $N_{\overline{G}}(\overline{T}_0)/\overline{T}_0$, we have that $\sigma(\dot{w}\overline{T}_0) = \dot{w}\overline{T}_0$, i. e. $\sigma(g)^{-1}g\sigma(g)^{-1}\sigma^2(g) = t_0 \in \overline{T}_0$. Hence ${}^gt_0 = t \in \overline{T}$ and ${}^{\sigma(g)g^{-1}}t = ({}^g\dot{w})^{-1}\sigma({}^g\dot{w}) \in \overline{T}$. It follows that $\sigma({}^g\dot{w}\overline{T}) = {}^g\dot{w}\overline{T}$. So, if $w \notin W_{\overline{C}_0}$, we conclude that ${}^g\dot{w}\overline{T}$ maps onto a non trivial element of the group $\frac{N_{N_{\overline{G}}(\overline{T})}(N_{\overline{C}}(\overline{T}))/\overline{T}\cap(N_{\overline{G}}(\overline{T})/\overline{T})_\sigma}{N_{\overline{C}}(\overline{T})/\overline{T}\cap(N_{\overline{G}}(\overline{T})/\overline{T})_\sigma}$. ☐

The rest of this subsection is devoted to unipotent elements in groups of Lie type.

Lemma 3.2.5. *Let $G = O^{p'}(\overline{G}_\sigma)$ be a finite group of Lie type with the base field $\mathbb{F}_{p^t}$, with p odd. If $p = 3$, suppose t even. Assume further that $\Phi(\overline{G}) \neq G_2, F_4, E_6, E_7, E_8$ if $p = 3$, and $\Phi(\overline{G}) \neq E_8$ if $p = 5$.*
Then every unipotent element u of order p is conjugate in G to some power $u^k \neq u$.

Proof. Under our assumptions p is a good prime. By point (i) of [40, Theorem 1.4], there exists a closed σ-stable subgroup $A_1(\overline{\mathbb{F}}_p)$ of $\overline{G}$ such that $u \in A_1(\overline{\mathbb{F}}_p)$. Clearly $O^{p'}((A_1(\overline{\mathbb{F}}_p))_\sigma)$ is isomorphic either to $\mathrm{SL}_2(p^{tm})$, or to $\mathbf{PSL}_2(p^{tm})$, for some positive integer $m > 0$. Up to conjugation inside $A_1(\overline{\mathbb{F}}_p)$, we may assume $u = \begin{pmatrix} 1 & \zeta \\ 0 & 1 \end{pmatrix}$ (or is equal to the projective image of this matrix) for some $\zeta \in \mathbb{F}_{p^{tm}}$. Under our assumptions, there exist $\eta \in \mathbb{F}_{p^t}$ such that $1 \neq \eta^2 = k \in \mathbb{F}_p$. Let x be the matrix $\begin{pmatrix} \eta^{-1} & 0 \\ 0 & \eta \end{pmatrix}$ or its projective image. Then $x \in G$, and u, $u^x = \begin{pmatrix} 1 & k\zeta \\ 0 & 1 \end{pmatrix} = u^k$ are conjugate in G. ☐

Lemma 3.2.6. *Let $u \in G = G_2(3^t)$ be an element of order 3.*
Then u is conjugate to u^{-1} in G.

Proof. By [28, Proposition 6.4] there exist 9 unipotent conjugacy classes in G. All of them may be found in Table 4, where α, β denote respectively a short and a long fundamental root of G_2, ζ is an element of $\mathbb{F}_{3^t}$ such that the polynomial $x^3 - x + \zeta$ is irreducible in $\mathbb{F}_{3^t}[x]$ and η is a non-square of F_{3^t}. Since $|x_1| = 9$ and x_2, x_3 are conjugate to x_1 in $G_2(\overline{\mathbb{F}}_3)$, we only need to verify that x_4, x_5, x_6, x_7, x_8 are conjugate to their inverses. Using the formulae $x_\beta(u)^{h_\alpha(t)} = x_\beta(t^{\frac{2(\alpha,\beta)}{(\alpha,\alpha)}}u)$ for each $\alpha, \beta \in \Phi$ (see [15, Proposition 6.4.1]), we get: $x_6^{h_\alpha(-1)} = x_6^{-1}$, $x_8^{h_\beta(-1)} = x_8^{-1}$, $x_4^{h_\beta(-1)} = x_4^{-1}$, and $x_5^{h_\beta(-1)} = x_5^{-1}$. Finally $|C_K(x_7)| \neq |C_K(x_i)|$ for all $i \neq 7$: thus also x_7 is conjugate to its inverse. ☐

Table 4: Unipotent classes in $G_2(q)$, $q = 3^t$.

representative x	$\|C_K(x)\|$

$x_0 = 1$	$q^6(q^2-1)(q^6-1)$
$x_1 = x_\alpha(1)x_\beta(1)$	$3q^2$
$x_2 = x_\alpha(1)x_\beta(1)x_{3\alpha+\beta}(\zeta)$	$3q^2$
$x_3 = x_\alpha(1)x_\beta(1)x_{3\alpha+\beta}(-\zeta)$	$3q^2$
$x_4 = x_{\alpha+\beta}(1)x_{3\alpha+\beta}(1)$	$2q^4$
$x_5 = x_{\alpha+\beta}(1)x_{3\alpha+\beta}(\eta)$	$2q^4$
$x_6 = x_{2\alpha+\beta}(1)$	$q^6(q^2-1)$
$x_7 = x_{2\alpha+\beta}(1)x_{3\alpha+2\beta}(1)$	q^6
$x_8 = x_{3\alpha+2\beta}(1)$	$q^6(q^2-1)$

Lemma 3.2.7. *Let $u \in G = F_4(3^t)$ be an element of order 3. Then u is conjugate to u^{-1} in G.*

Proof. By [39, Table 6] there exist 28 unipotent conjugacy classes of G. All of them may be found in Table 5. Recall that in an Euclidean 4-dimensional space with orthonormal base $\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4$ all roots of F_4 may be written as $\{\pm\varepsilon_i \pm \varepsilon_j, \pm\varepsilon_i, \frac{1}{2}(\pm\varepsilon_1 \pm \varepsilon_2 \pm \varepsilon_3 \pm \varepsilon_4)\}$. In Table 5 the symbols $\pm i \pm j$, $\pm i$, and $\pm 1 \pm 2 \pm 3 \pm 4$ denote the roots $\pm\varepsilon_i \pm \varepsilon_j$, $\pm\varepsilon_i$, and $\frac{1}{2}(\pm\varepsilon_1 \pm \varepsilon_2 \pm \varepsilon_3 \pm \varepsilon_4)$ respectively, η is a fixed non-square element of $\mathbb{F}_{3^t}$, ξ is a fixed element of $\mathbb{F}_{3^t}$ such that $x^2+\xi x+\eta$ is an irreducible polynomial in $\mathbb{F}_{3^t}[x]$, ζ is a fixed element of $\mathbb{F}_{3^t}$ such that $x^3 - x + \zeta$ is an irreducible polynomial in $\mathbb{F}_{3^t}[x]$. By using [39, Table 7] one may easily verify that $|x_9| = |x_{10}| > 3, |x_i| > 3$ for all $i \geq 12$. Indeed, by [39, Table 7] we have that elements x_9 and x_{10} are conjugate in $F_4(\overline{\mathbb{F}}_3)$. They also are conjugate to an element $c_7 = x_{r_1}(1)x_{r_2}(1)x_{r_3}(1)$, where the roots r_1, r_2 and r_3 я are fundamental roots in a root system of type A_3. But it is evident, that $|c_7| > 3$. In all cases when $|x_i| > 3$, we proceed in a similar way. In the remaining cases one can see that $|C_K(x_i)| \neq |C_K(x_j)|$ for all $i \neq j$. So if $|x_i| = 3$, then $i = 1, 2, 3, 4, 5, 6, 7, 8, 11$, and x_i is conjugate to its inverse under G. ☐

Table 5: Unipotent classes $F_4(q)$, $q = 3^t$

representative x	$\lvert C_K(x)\rvert$
$x_0 = 1$	$\lvert K\rvert$
$x_1 = x_{1+2}(1)$	$q^{24}(q^2-1)(q^4-1)(q^6-1)$
$x_2 = x_{1-2}(1)x_{1+2}(-1)$	$2q^{21}(q^2-1)(q^3-1)(q^4-1)$
$x_3 = x_{1-2}(1)x_{1+2}(-\eta)$	$2q^{21}(q^2-1)(q^3+1)(q^4-1)$
$x_4 = x_2(1)x_{3+4}(1)$	$q^{20}(q^2-1)^2$
$x_5 = x_{2-3}(1)x_4(1)x_{2+3}(1)$	$2q^{17}(q^2-1)(q^3-1)$
$x_6 = x_{2-3}(1)x_4(1)x_{2+3}(\eta)$	$2q^{17}(q^2-1)(q^3+1)$
$x_7 = x_2(1)x_{1-2+3+4}(1)$	$q^{14}(q^2-1)(q^6-1)$
$x_8 = x_{2-3}(1)x_4(1)x_{1-2}(1)$	$q^{16}(q^2-1)$
$x_9 = x_{2-3}(1)x_{3-4}(1)x_{3+4}(-1)$	$2q^{12}(q^2-1)^2$
$x_{10} = x_{2-3}(1)x_{3-4}(1)x_{3+4}(-\eta)$	$2q^{12}(q^4-1)$
$x_{11} = x_{2+3}(1)x_{1+2-3-4}(1)x_{1-2+3+4}(1)$	$q^{14}(q^2-1)$
$x_{12} = x_{2-3}(1)x_4(1)x_{1-4}(1)$	$2q^{12}(q^2-1)$
$x_{13} = x_{2-3}(1)x_4(1)x_{1-4}(\eta)$	$2q^{12}(q^2-1)$
$x_{14} = x_{2-4}(1)x_{3+4}(1)x_{1-2}(-1)x_{1-3}(-1)$	$24q^{12}$
$x_{15} = x_{2-4}(1)x_{3+4}(1)x_{1-2}(-\eta)x_{1-3}(-1)$	$8q^{12}$
$x_{16} = x_{2-4}(1)x_{2+4}(-\eta)x_{1-2+3+4}(1)x_{1-3}(-1)$	$4q^{12}$
$x_{17} = x_{2-4}(1)x_{3+4}(1)x_{1-2-3+4}(1)x_{1-2}(-\eta)x_{1-3}(\xi)$	$4q^{12}$
$x_{18} = x_2(1)x_{3+4}(1)x_{1-2+3-4}(1)x_{1-2}(-1)x_{1-3}(\zeta)$	$3q^{12}$
$x_{19}x_{2-3}(1)x_{3-4}(1)x_4(1)$	$q^8(q^2-1)$

$x_{20} = x_2(1)x_{3+4}(1)x_{1-2-3-4}(1)$	$q^8(q^2-1)$
$x_{21} = x_{2-4}(1)x_3(1)x_{2+4}(1)x_{1-2-3+4}(1)$	$2q^8$
$x_{22} = x_{2-4}(1)x_3(1)x_{2+4}(\eta)x_{1-2-3+4}(1)$	$2q^8$
$x_{23} = x_{2-3}(1)x_{3-4}(1)x_4(1)x_{1-2}(1)$	$2q^6$
$x_{24} = x_{2-3}(1)x_{3-4}(1)x_4(1)x_{1-2}(\eta)$	$2q^6$
$x_{25} = x_{2-3}(1)x_{3-4}(1)x_4(1)x_{1-2-3-4}(1)$	$3q^4$
$x_{26} = x_{2-3}(1)x_{3-4}(1)x_4(1)x_{1-2-3-4}(1)x_{1-2+3+4}(\zeta)$	$3q^4$
$x_{27} = x_{2-3}(1)x_{3-4}(1)x_4(1)x_{1-2-3-4}(1)x_{1-2+3+4}(-\zeta)$	$3q^4$

Lemma 3.2.8. *Let $u \in G$ be an element of order 3, where $G = E_6(3^t)$ or $G = {}^2E_6(3^t)$ is a canonical finite group of Lie type.*
Then u is conjugate to u^{-1} in G.

Proof. Let $\overline{G}$ and σ be such that $G = O^{p'}(\overline{G})$. Since the characteristic equals 3, we have that $Z(\overline{G}_{sc}) = 1$. So we may assume $\overline{G} = \overline{G}_{sc}$ to be universal. Thus $\overline{G}$ is simply connected and $G = \overline{G}_\sigma$. We assemble the information from [36, Lemmas 4.2, 4.3, 4.4, and Theorem 4.13] on conjugacy classes of unipotent elements of G in Table 6. In Table 6 we substitute the root $\alpha_1 r_1 + \alpha_2 r_2 + \alpha_3 r_3 + \alpha_4 r_4 + \alpha_5 r_5 + \alpha_6 r_6$, where $r_1, r_2, r_3, r_4, r_5, r_6$ form a fundamental system of E_6, by the 6-tuple $\alpha_1\alpha_2\alpha_3\alpha_4\alpha_5\alpha_6$ of its coefficients.

Note that if $n \geqslant 3$ and $r_1, r_2, \dots, r_n$ are fundamental roots of root system of type A_n, then $|x_{r_1}(1)x_{r_2}(1)\dots x_{r_n}(1)| > 3$. By using this fact we obtain, that $|x_4| > 3$, $|x_7| > 3$, $|x_8| > 3$, $|x_i| > 3$, where $i \geqslant 10$, $i \neq 12, 16$. Thus we have to consider remaining cases only. We have that $x_1^{h_{r_1}(\lambda)} = x_1^{-1}$, where λ is a square root of -1 in $\overline{\mathbb{F}}_3$. For each $x \in \overline{G}$ denote by $\mathrm{Ccl}(x)$ its conjugacy class in $\overline{G}$. Since $C_{\overline{G}}(x_1) = C_{\overline{G}}(x_1)^0$, from [34, Theorem 8.5] we have that for every Frobenius map σ and for every $x \in \mathrm{Ccl}(x_1) \cap \overline{G}_\sigma$, the elements x and x^{-1} are conjugate under $\overline{G}_\sigma$. So if $x \in \mathrm{Ccl}(x_1) \cap G$, then x is conjugate to its inverse.

Table 6: Unipotent classes in $E_6(\overline{\mathbb{F}}_3)$

representative x	$C = C_G(x)$ $\lvert C : C^0\rvert$
$x_1 = x_{10000}(1)$	1
$x_2 = x_{100000}(1)x_{001000}(1)$	2
$x_3 = x_{100000}(1)x_{000100}(1)$	1
$x_4 = x_{100000}(1)x_{001000}(1)x_{000100}(1)$	1
$x_5 = x_{100000}(1)x_{001000}(1)x_{000010}(1)$	1
$x_6 = x_{100000}(1)x_{000100}(1)x_{000001}(1)$	1
$x_7 = x_{100000}(1)x_{001000}(1)x_{000100}(1)x_{000010}(1)$	1
$x_8 = x_{100000}(1)x_{001000}(1)x_{000100}(1)x_{000001}(1)$	1
$x_9 = x_{100000}(1)x_{001000}(1)x_{000010}(1)x_{000001}(1)$	1
$x_{10} = x_{100000}(1)x_{001000}(1)x_{010000}(1)x_{000010}(1)$	1
$x_{11} = x_{100000}(1)x_{001000}(1)x_{000100}(1)x_{010000}(1)x_{000001}(1)$	1
$x_{12} = x_{100000}(1)x_{001000}(1)x_{000010}(1)x_{000001}(1)x_{010000}(1)$	1
$x_{13} = x_{100000}(1)x_{001000}(1)x_{000100}(1)x_{000010}(1)x_{000001}(1)$	1
$x_{14} = x_{010000}(1)x_{001000}(1)x_{000100}(1)x_{000010}(1)$	1
$x_{15} = x_{010000}(1)x_{001000}(1)x_{000100}(1)x_{010110}(1)$	6
$x_{16} = x_{000001}(1)x_{000010}(1)x_{001000}(1)x_{010000}(1)$	1
$x_{17} = x_{010000}(1)x_{001000}(1)x_{000010}(1)x_{101100}(1)$	1
$x_{18} = x_{000010}(1)x_{000100}(1)x_{001000}(1)x_{100000}(1)x_{000001}(1)x_{111111}(1)$	2
$x_{19} = x_{010000}(1)x_{000100}(1)x_{000010}(1)x_{000001}(1)x_{101000}(1)x_{001110}(1)$	1
$x_{20} = x_{100000}(1)x_{010000}(1)x_{001000}(1)x_{000100}(1)x_{000010}(1)x_{000001}(1)$	3

For the other x_i-s such that 3, with $i \neq 2$, we proceed in the same way. We are left with x_2.

By [34, Theorem 8.5] we have that, for every Frobenius map σ, $\mathrm{Ccl}(x_2)\cap\overline{G}_\sigma$ consists of two conjugacy classes of $G=\overline{G}_\sigma$. Assume first that $G=E_6(3^t)$. Then, by [36, Lemmas 4.2 and 4.4] we have, that if $x\in\mathrm{Ccl}(x_2)\cap G$, then x is conjugate under G either to $y_1=x_{100000}(1)x_{001000}(1)$, or to $y_2=x_{100000}(1)x_{001000}(1)x_{000001}(1)x_{122321}(\eta)$, where η is a nonsquare in $\mathbb{F}_{3^t}$. By [36, Lemma 4.2] $|C_G(y_1)|=2q^{26}(q^2-1)^2(q^3-1)^2$, by [36, Lemma 4.4] $|C_G(y_2)|=2y^{26}(q^4-1)(q^6-1)$. For $i=1,2$, let $\mathrm{Ccl}_G(y_i)$ be the conjugacy class of y_i in G. Since $|C_G(y_1)|\neq|C_G(y_2)|$ we have, that y_i is conjugate to its inverse under G for $i=1,2$. So if $x\in\mathrm{Ccl}_G(y_1)$, or $x\in\mathrm{Ccl}_G(y_2)$, then x is conjugate to its inverse under G. Now assume that $G={}^2E_6(3^t)$ and denote $E_6(3^{2t})$ by G_1. Then $G=(G_1)_\tau$ for some graph automorphism τ of G_1. There exists a Frobenius map σ such that $G_1=\overline{G}_\sigma$, $G=\overline{G}_{\sigma\tau}$ (see [32, (7-2)]). Let Ccl_1 and Ccl_2 be two conjugacy classes of G_1 contained in $\mathrm{Ccl}(x_2)\cap G_1$. We prove that every $x\in\mathrm{Ccl}_i$, $i=1,2$, is conjugate to x^{-1} в G_1. Since $\mathrm{Ccl}(x_2)\cap G$ consists of two conjugacy classes of G, we have that $\mathrm{Ccl}_1\cap G$ consists of one conjugacy class and $\mathrm{Ccl}_2\cap G$ consists of one conjugacy class. So, every $x\in\mathrm{Ccl}_i\cap G$, $i=1,2$ is conjugate to its inverse under G. ☐

Lemma 3.2.9. *Let $O^{p'}(\overline{G}_\sigma)\leq G\leq\overline{G}_\sigma$ be a finite adjoint group of Lie type over a field of odd characteristic p and the root system Φ of $\overline{G}$ is one of the following: A_n ($n\geqslant 2$), D_n ($n\geqslant 4$), B_n ($n\geqslant 3$), G_2, F_4, E_6, E_7 or E_8; and $G\not\simeq{}^2G_2(3^{2n+1})$. Let U be a maximal unipotent subgroup of G, H be a Cartan subgroup of G, normalizing U, and Q is a Sylow 2-subgroup of H.*
Then $C_U(Q)=\{e\}$.

Proof. Clearly we enough to prove the lemma for the case $G=O^{p'}(\overline{G}_\sigma)=O^{p'}(G)$, i. e., we may assume that G is a canonical adjoint group of Lie type.

First assume that G is split. Assume that $C_U(Q)\neq\{e\}$ and $u\in C_U(Q)\setminus\{e\}$. Consider decomposition (1) of $u=\prod_{r\in\Phi^+}x_r(t_r)$, where t_r are from the definition field $\mathbb{F}_q$ of G. In view of [15, Theorem 5.3.3(ii)] this decomposition is unique. Since for every $h(\chi)\in H$, $r\in\Phi$, $t\in\mathbb{F}_q$ the formulae $h(\chi)x_r(t)h(\chi)^{-1}=x_r(\chi(r)t)$ holds (see [15, p. 100]), then we obtain that each multiplier $x_r(t_r)$ in decomposition (1) of u is in $C_U(Q)$. So we may assume that $u=x_r(t)$ for some $r\in\Phi^+$ and $t\in\mathbb{F}_q^*$. Under our restriction on Φ, by Hartley-Shute lemma 1.5.5, there exists $h(\chi)\in H$ such that $\chi(r)=-1$. Since $h(\chi)^2=h(\chi^2)$ (see [15, p. 98]), then we have that $\chi^2(r)=1$, i. e., $|h(\chi)^2|<|h(\chi)|$. Hence, $|h(\chi)|$ is even and we may write $h(\chi)=h_2\cdot h_{2'}=h(\chi_1)\cdot h(\chi_2)$, a decomposition of $h(\chi)$ as a product of its 2- and $2'$- parts. Now $\chi(r)=\chi_1(r)\cdot\chi_2(r)$, therefore $\chi_1(r)=-1$ and $\chi_2(r)=1$. Thus $h(\chi_1)x_r(t)h(\chi_1)^{-1}=x_r(-t)\neq x_r(t)$. Since $h(\chi_1)\in Q$, the obtained equation contradicts to the choice of $x_r(t)\in C_U(Q)$.

Assume that $G\simeq{}^2A_n(q)$, $G\simeq{}^2D_n(q)$, or $G\simeq{}^2E_6(q)$, then $\Phi(\overline{G})$ equals A_n, D_n and E_6 respectively. Denote by $\bar{r}$ the image of r of Φ under the corresponding symmetry. In terms of [15], the root system $\Phi(\overline{G})$ is expressible as a union of equivalency classes Ψ_i-s, while each Ψ_i has type either A_1, or $A_1\times A_1$, or A_2. In view of [15, Proposition 13.6.1], the equality $U=\prod_i X_{\Psi_i}$ holds, where

$$X_{\Psi_i}=\{x_r(t)\mid t\in\mathbb{F}_q\},$$

if $\Psi_i=\{r\}$ has type A_1 (here $r=\bar{r}$);

$$X_{\Psi_i}=\{x_r(t)x_{\bar{r}}(t^q)\mid t\in\mathbb{F}_{q^2}\},$$

if $\Psi_i=\{r,\bar{r}\}$ has type $A_1\times A_1$ (here $r\neq\bar{r}$, and $r+\bar{r}\notin\Phi(\overline{G})$);

$$X_{\Psi_i}=\{x_r(t)x_{\bar{r}}(t^q)x_{r+\bar{r}}(u)\mid t\in\mathbb{F}_{q^2},u+u^q=-N_{r,\bar{r}}tt^q\},$$

if $\Psi_i = \{r, \bar{r}, r+\bar{r}\}$ has type A_2 (here $r \neq \bar{r}$ and $r+\bar{r} \in \Phi(\overline{G})$). Now if $h(\chi)$ is an element of H, then the following equalities hold (see [15, p. 263]):

$$h(\chi)x_r(t)h(\chi)^{-1} = x_r(\chi(r)t),$$

if $r = \bar{r}$ and $\Psi_i = \{r\}$ has type A_1;

$$h(\chi)x_r(t)x_{\bar{r}}(t^q)h(\chi)^{-1} = x_r(\chi(r)t)x_{\bar{r}}(\chi(\bar{r})t^q),$$

if $r \neq \bar{r}$, $r+\bar{r} \notin \Phi(\overline{G})$ and $\Psi_i = \{r, \bar{r}\}$ has type $A_1 \times A_1$;

$$h(\chi)x_r(t)x_{\bar{r}}(t^q)x_{r+\bar{r}}(u)h(\chi)^{-1} = x_r(\chi(r)t)x_{\bar{r}}(\chi(\bar{r})t^q)x_{r+\bar{r}}(\chi(r+\bar{r})u),$$

if $r \neq \bar{r}$, $r+\bar{r} \in \Phi(\overline{G})$ and $\Psi_i = \{r, \bar{r}, r+\bar{r}\}$ has type A_2.

Let u be a nontrivial element from $C_U(Q)$. Then u contains a nontrivial multiplier from X_{Ψ_i} for some i. In view of uniqueness of decomposition into the product $\prod_i X_{\Psi_i}$ (see [15, Proposition 13.6.1]) we may assume that $u \in X_\Psi$.

Assume that Ψ has type A_1, i. e., $u = x_r(t)$, $t \in \mathbb{F}_q$, $r = \bar{r}$. In view of Hartley-Shute lemma 1.5.5, for each $s \in \mathbb{F}_q$ there exists $h(\chi) \in H$ such that $\chi(r) = s$. Take $s = -1$. Then there exists $h(\chi) \in H$ such that $\chi(r) = -1$. Since $h(\chi)^2 = h(\chi^2)$ (see formulae on p. 98 from [15]), then we have that $\chi^2(r) = 1$, i. e. $|h(\chi)^2| < |h(\chi)|$. Hence order $|h(\chi)|$ is even and we may write $h(\chi) = h_2 \cdot h_{2'} = h(\chi_1) \cdot h(\chi_2)$, a decomposition of $h(\chi)$ into the product of its 2- and $2'$- parts. Now $\chi(r) = \chi_1(r) \cdot \chi_2(r)$, therefore $\chi_1(r) = -1$ and $\chi_2(r) = 1$. Thus $h(\chi_1)x_r(t)h(\chi_1)^{-1} = x_r(-t) \neq x_r(t)$. So the case $u = x_r(t)$ and $\Psi = \{r\}$ has type A_1 is impossible.

Assume that $\Psi = \{r, \bar{r}\}$ has type $A_1 \times A_1$. By Hartley-Shute lemma 1.5.5 for every $s \in \mathbb{F}_{q^2}$ there exists $h(\chi) \in H$ such that $\chi(r) = s^2$. Since there exists $s \in \mathbb{F}_{q^2}$ such that $s^2 = -1$, then there exists $h(\chi) \in H$ such that $\chi(r) = -1$. As above $h(\chi)$ can be written as $h(\chi_1) \cdot h(\chi_2)$, a product of its 2- and $2'$- parts. Then $\chi_1(r) = -1$, so

$$h(\chi_1)x_r(t)x_{\bar{r}}(t^q)h(\chi_1)^{-1} = x_r(-t)x_{\bar{r}}(-t^q) \neq x_r(t)x_{\bar{r}}(t^q).$$

Thus the case $u = x_r(t)x_{\bar{r}}(t^q)$ and $\Psi = \{r, \bar{r}\}$ has type $A_1 \times A_1$ is impossible.

Assume that $\Psi = \{r, \bar{r}, r+\bar{r}\}$ has type A_2. By Hurtley-Shute lemma 1.5.5, for each $s \in \mathbb{F}_{q^2}$ there exists $h(\chi) \in H$ such that $\chi(r) = s^3$. Choose $s = -1$, then there exists $h(\chi) \in H$ such that $\chi(r) = -1$. Again $h(\chi) = h(\chi_1) \cdot h(\chi_2)$ is expressible as the product of its 2- and $2'$- parts and $\chi_1(r) \neq 1$. Then

$$\begin{aligned} &h(\chi_1)x_r(t)x_{\bar{r}}(t^q)x_{r+\bar{r}}(u)h(\chi_1)^{-1} = \\ &\qquad x_r(-t)x_{\bar{r}}(\chi_1(-t^q)x_{r+\bar{r}}(\chi_1(r+\bar{r})u) \neq \\ &\qquad\qquad x_r(t)x_{\bar{r}}(t^q)x_{r+\bar{r}}(u) \end{aligned}$$

for $t \neq 0$. If $t = 0$, then choose s so that $s^2 = -1$. Then $\chi_1(r+\bar{r}) = -1$ and, as above, we obtain the inequality. Hence this case is impossible.

Assume at last that $G \simeq {}^3D_4(q)$. In terms from [15], a root system $\Phi(\overline{G})$ is expressible as a union of equivalency classes Ψ_i, when each Ψ_i has type either A_1, or $A_1 \times A_1 \times A_1$. In view of [15, Proposition 13.6.1], the equality $U = \prod_i X_{\Psi_i}$ holds, where

$$X_{\Psi_i} = \{x_r(t) \mid t \in \mathbb{F}_q\},$$

if $\Psi_i = \{r\}$ has type A_1 (here $r = \bar{r}$);

$$X_{\Psi_i} = \{x_r(t)x_{\bar{r}}(t^q)x_{\bar{\bar{r}}}(t^{q^2}) \mid t \in \mathbb{F}_{q^3}\},$$

if $\Psi_i = \{r, \bar{r}, \bar{\bar{r}}\}$ has type $A_1 \times A_1 \times A_1$ (here $r \neq \bar{r}$ and $r + \bar{r} \notin \Phi(\overline{G})$). In both cases, by Hartley-Shute lemma 1.5.5, there exists $h(\chi) \in H$ such that $\chi(r) = -1$. As above we may assume that $h(\chi)$ is a 2-element, i. e. $h(\chi) \in Q$ and $h(\chi)$ does not centralizes nonidentical elements from X_{Ψ_i}, and the statement of Lemma 3.2.9 follows in the last case. ☐

Lemma 3.2.10. *In the notations of Lemma* 3.2.9*, with p odd, let K be a Carter subgroup of G such that $|K| = 2^a p^b$.*
Then $a > 0$. More precisely, up to conjugation, $O_p(K) \leq C_U(Q)$. In particular, under the assumptions of Lemma 3.2.9*, K is a 2-group.*

Proof. The condition $a = 0$ would imply $K = U$. But U is normalized by H which is non-trivial as p is odd and G is simple. Thus $a > 0$. Now, assume $b > 0$. By Borel-Tits theorem (Lemma 1.5.4), K i contained in a proper parabolic subgroup P of G and $O_p(K) \leq O_p(P)$. Since $P = LO_p(P)$, where L is a Levi factor of P, from Lemma 2.4.1 it follows that $KO_p(P)/O_p(P) \cong O_2(K)$ is a Carter subgroup of $P/O_p(P) \cong L$. Thus $O_2(K)$ is a Sylow 2-subgroup of L. But L contains H, therefore we may assume that $Q \leq K$. It follows that $O_p(K) \leq C_U(Q)$. ☐

Lemma 3.2.11. *Let G be a non-Abelian simple group not of Lie type.*
Then every element z of odd order is conjugate to some $z^k \neq z$.

Proof. By the classification of finite simple groups, G is either alternating, or sporadic. Our claim can be checked directly in first case, and using the description of the conjugacy classes given in [21] in the second case. ☐

3.3 Almost simple groups which are not minimal counter examples

In this subsection A denotes a minimal almost simple group that is a minimal counter example (see definition 2.4.7). If G is a group of Lie type, we denote by $\mathrm{Field}(G)$ the subgroup of $\mathrm{Aut}(G)$ generated by inner, diagonal, and field automorphisms. If G is a simple group which is not of Lie type, we set $\widehat{G} = G$ to unify notations. More over for each $x \in G$ we assume that composition factors of the centralizer $C_G(s)$ are known simple groups, and so $C_G(s)$ satisfies **(C)**. As we noted in subsection 2.4, this assumption is always true. We say it here in order to emphasize that all results do not depend on the classification of finite simple groups.

Lemma 3.3.1. *Let A be a minimal counter example and $G = F^*(A)$. Assume that for every element $z \in \widehat{G}$ of odd prime order, z is conjugate to some $z^k \neq z$ in G.*
Then A is not a minimal counter example if one of the following holds:

(a) *$|A : \widehat{G} \cap A|$ is a 2-power;*

(b) *$|\widehat{G} : (\widehat{G} \cap A)|$ is a 2-power and , if $\Phi(\overline{G})$ has type D_4, then $|(\mathrm{Field}(G)\cap) : (\widehat{G} \cap A)|_{2'} > 1$;*

(c) *for every odd prime r and every Sylow r-subgroup R of A, either $R \cap G$ has no complement in R, or all such complements are conjugate in A.*

Proof. Let K, H be nonconjugate Carter subgroups of A. Note that by Lemma 2.4.2(b) it follows that $K \cap \widehat{G}$ and $H \cap \widehat{G}$ are 2-groups. We prove (c) first, that we show that (a) and (b) follows from (c).

(c) By Theorem 2.1.4 and Lemma 2.4.1, we obtain that $KG/G = HG/G = A/G$. In particular, if r is a prime divisor of $|A/G|$, then r divides both $|K|$ and $|H|$. By Lemma 2.4.2 and by conditions of this lemma, it follows that $K \cap \widehat{G}$ and $H \cap \widehat{G}$ do not contain elements of odd prime order, i. e., are 2-groups. If $R \cap G$ has no a complement in R we get a contradiction immediately, if all such complements are conjugate in A, we obtain a contradiction with Lemma 2.4.2(c). Thus we obtain that $|A/G|$ is a 2-power, hence K and H are 2-groups, that is impossible.

Now (a) evidently follows from (c). As to (b), then it also follows from (c), by using the conjugacy of complements, that follows by Lemma 4.2.6. □

Note that all non-Abelian composition factors of the centralizer of every element of the alternating group Alt_n are alternating groups of lower degree. So Lemmas 3.2.11 and 3.3.1 and induction by n imply immediately that Carter subgroups of $\mathrm{Aut}(\mathrm{Alt}_n)$ with $n \geqslant 5$ either are Sylow 2-subgroups or do not exist. The same statement holds for sporadic groups. Thus the following statement is true.

Lemma 3.3.2. *Let S be a finite non-Abelian simple group, that is either sporadic, or alternating.*
Then, for every subgroup A of $\mathrm{Aut}(S)$, a Carter subgroup either does not exist, or is a Sylow 2-subgroup.

Theorem 3.3.3. *Let G be a finite adjoint group of Lie type such that $G = \mathbf{P}\Omega^{\pm}_{2(2\ell+1)}(p^t)$, and assume that $\ell \geqslant 2$.*
Then G is not a minimal counter example.

Proof. Assume that our statement is false. Then G contains a Carter subgroup K, that is not a 2-group. Let $s \in Z(K)$ be an element of odd prime order r. Then we may assume that s is semisimple, except, probably, the case, when $p \neq 2$ and $|K| = 2^a p^b$. But this is impossible in view of Lemmas 3.2.9 and 3.2.10. Hence s is semisimple and from $K \leq C_G(s)$ it follows that $C_G(s)$ is self-normalizing in G (see Lemma 2.4.2(a)). Now let $\overline{G} = \Omega_{2(2\ell+1)}(\overline{\mathbb{F}}_p)$ and σ be such that $\overline{G}_\sigma = \Omega^{\pm}_{2(2\ell+1)}(p^t)$. More over set K_0 to be equal to the preimage of K in $\overline{G}_\sigma$. Clearly K_0 is a Carter subgroup of $\overline{G}_\sigma$ and we may identify s with its preimage in $\overline{G}_\sigma$, since the center of $\overline{G}_\sigma$ has order 2 or 4. Since $|s|$ is odd, Lemma 3.2.1 implies that $C = C_{\overline{G}}(s)$ is a connected reductive subgroup of maximal rank of $\overline{G}$ (see Lemmas 1.5.1 and 1.5.2). More over C is a proper subgroup of $\overline{G}$, since $s \notin Z(\overline{G})$. By Lemma 3.2.4 the group $N_W(W_C)/W_C$ is isomorphic to $N_{\overline{G}}(C)/C$. By using the description of $N_W(W_C)/W_C$, given in [18, Proposition 10] and Lemma 3.2.4, we conclude that $N_G(C_G(s))/C_G(s)$ is trivial only if $W_C^{\perp}$ and $\mathrm{Aut}_W(\Delta_C)$ are both trivial. From assumption $\ell \geq 2$ it follows that this occurs precisely when $m_1 = 0$ and $m_{2\ell+1} = 1$ (in the notations from [18]). In this case $C = A_{2\ell}(\overline{\mathbb{F}}_p) * S$, where S is a 1-dimensional torus. By using the fact that $\overline{G}$ contains exactly one class of connected reductive subgroups isomorphic to C, and assuming that $\overline{G}$ preserves the bilinear form induced by $J = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$, we may identify C with the image of $\mathrm{GL}_{2\ell+1}(\overline{\mathbb{F}}_p)$ under a monomorphism φ such that

$$A \mapsto \begin{pmatrix} A & 0 \\ 0 & (A^{-1})^t \end{pmatrix}.$$

By Lang-Steinberg theorem (Lemma 1.5.3), we may assume that either $C_\sigma = \varphi(\mathrm{GL}_{2\ell+1}(p^t))$, or $C_\sigma = \varphi(\mathrm{GU}_{2\ell+1}(p^{2t}))$. Since K_0 is a Carter subgroup of C_σ and $\ell \geq 2$, by [26], [27], and Theorem 1.5.6 it follows that K_0 is the normalizer of a Sylow 2-subgroup P of C_σ, and either $p^t = 2$ (and $C_\sigma = \varphi(\mathrm{GL}_{2\ell+1}(p^t))$), or p is odd. From $s \in Z(C_\sigma)$ it follows that $r = |s|$ divides $p^t - 1$ if $C_\sigma \simeq \mathrm{GL}_{2\ell+1}(p^t)$, and that r divides $p^t + 1$ if $C_\sigma \simeq \mathrm{GU}_{2\ell+1}(p^{2t})$. In particular p is odd. By using known structure of normalizers of Sylow 2-subgroups in classical groups (see [9] and [20]), we may assume that K_0 is a subgroup of the following group

$$L = \left\{ \begin{pmatrix} B & 0 \\ 0 & \beta \end{pmatrix}^{\varphi} \mid B \in \mathrm{GL}_{2\ell}(p^t)\ ,\ \beta \in \overline{\mathbb{F}}_q^{*} \right\} \text{ if } C_\sigma \simeq \mathrm{GL}_{2\ell+1}(p^t)$$

$$L = \left\{ \begin{pmatrix} B & 0 \\ 0 & \beta \end{pmatrix}^{\varphi} \mid B \in \mathrm{GU}_{2\ell}(p^{2t})\ ,\ \beta^{p^t+1} = 1 \right\} \text{ if } C_\sigma \simeq \mathrm{GU}_{2\ell+1}(p^{2t}).$$

As we noted above, there exists $y \in L$ such that $y = \begin{pmatrix} I_{2\ell} & 0 \\ 0 & \gamma \end{pmatrix}^{\varphi}$ where γ has order r. Since y is in the center of L, it is also in the center of K_0. Thus $K_0 \leq C_{C_{G_\sigma}(s)}(y) = \big(C_C(y)\big)_\sigma$. From isomorphism $C \simeq \mathrm{GL}_{2\ell+1}(\overline{\mathbb{F}}_p)$ it follows that $C_C(y)$ is a connected reductive σ-invariant subgroup of maximal rank of $\overline{G}$. Thus, in view of above mentioned result by Carter [18, Proposition 10], $\big(C_C(y)\big)_\sigma$ is self-normalizing in $\overline{G}_\sigma$ only if $C_C(y)$ is conjugate to C. But $\dim(C_C(y)) < \dim(C)$, since y is not in the center of C. Thus $(C_C(y))_\sigma$ is not self-normalizing in $\overline{G}_\sigma$. Since $Z(\overline{G}) \leq C_C(y)$, it follows that the factor group $(C_C(y))_\sigma/(Z(\overline{G}))_\sigma$ is not self-normalizing in $\overline{G}_\sigma/(Z(\overline{G}))_\sigma = G$. Thus we have obtained a contradiction with Lemma 2.4.2(a), since K is contained in $(C_C(y))_\sigma/(Z(G))_\sigma$, and $(C_C(y))_\sigma/(Z(G))_\sigma$ satisfies **(C)**. ☐

Theorem 3.3.4. *Let $E_6^\varepsilon(p^t) \leq G \leq \widehat{E_6^\varepsilon(p^t)}$. Then G is not a minimal counter example.*

Proof. Assume that our claim is false. Then, by Lemma 2.4.2(c), G admits a Carter subgroup K, which does not contain any Sylow 2-subgroup of G. In particular K is not a 2-group. Let $s \in Z(K)$ have odd prime order r. By Lemmas 3.2.5, 3.2.8, and 2.4.2, p does not divide $|K|$. Hence s is semisimple and K is contained in $C_G(s)$, which, in virtue of Lemma 2.4.2(a), is self-normalizing. If $|s| \neq 3$, then, by Lemma 3.2.2, it follows that $C_{\overline{G}}(s)$ is connected. If $|s| = 3$, then, by Lemma 1.5.2, it follows that $|C : C^0|$ divides $\Delta = 3$. Direct calculations by using [24] and [36] show that $C_G(s)$ is not self-normalizing, if $|s| = 3$. Therefore we may assume that $|s| \neq 3$ and $C_{\overline{G}}(s)$ is connected. Since $C_G(s)$ is self-normalizing, Lemma 3.2.4 shows that $C = C_{\overline{G}}(s)$ is self-normalizing as well. By [36], we obtain that C is self-normalizing if and only if $C = A_4(\overline{\mathbb{F}}_p) \circ A_1(\overline{\mathbb{F}}_p) \circ S$, or $C = D_5(\overline{\mathbb{F}}_p) \circ S$, where S is a 1-dimensional torus of $\overline{G}$.

If $C = A_4(\overline{\mathbb{F}}_p) \circ A_1(\overline{\mathbb{F}}_p) \circ S$, then like in proof of Theorem 3.3.3, we may find an element $y \in Z(K)$ such that $|y| = r$ and $C_G(\langle s \rangle \times \langle y \rangle)$ is not self-normalizing; a contradiction with Lemma 2.4.2.

So, assume that $C = D_5(\overline{\mathbb{F}}_p) \circ S$. Then $C_G(s) = C \cap G = HL$, where H is a Cartan subgroup of G and $L = O^{p'}(C_G(s))$ is either $D_5(p^t)$ or ${}^2D_5(p^t)$. Since $|\hat{L} : L|$ divides 4, then

$$O_{2'}(H) = (O_{2'}(H) \cap Z(C_G(s))) \times (O_{2'}(H) \cap L).$$

Denoting by Q a Sylow 2-subgroup of $C_G(s)$, we claim that $N_{C_G(s)}(Q) = QZ(C_G(s))$. Indeed, let x be an element of $N_{C_G(s)}(Q)$. From $H = O_2(H) \times O_{2'}(H)$ and $C_G(s) = HL$, we can write $x = h_1 z l$ with $h_1 \in O_2(H)$, $z \in O_{2'}(H) \cap Z(C_G(s))$, $l \in L$. We may clearly assume $O_2(H) \leq Q$: thus $l \in N_{C_G(s)}(Q)$. From L normal in $C_G(s)$, it follows $l \in N_L(Q \cap L)$. By [9], $N_L(Q \cap L) =$

$Q\cap L$, so, $l\in Q$. We conclude that $N_{C_G(s)}(Q)=QZ(C_G(s))$ is nilpotent, hence a Carter subgroup of $C_G(s)$. Since $C_G(s)<G$, all Carter subgroups in $C_G(s)$ are conjugate. Therefore, up to conjugation, $K=N_{C_G(s)}(Q)$. In virtue of the formula $|(C)_\sigma|=|M_\sigma|\cdot|(Z(C)^0)_\sigma|$, where $M_\sigma=L$ in our notation (see [18]), we have that $|G:C_G(s)|$ is odd, so Q is a Sylow 2-subgroup of G, a contradiction. ☐

Our results are summarized in the following theorem.

Theorem 3.3.5. *An almost simple group A, with socle с цоколем G is not a minimal counter example in the following cases:*

(a) *G is alternating, sporadic, or one of the following groups: $A_1(p^t)$, $B_\ell(p^t)$, $C_\ell(p^t)$, where t is even if $p=3$; ${}^2B_2(2^{2n+1})$, $G_2(p^t)$, $F_4(p^t)$, ${}^2F_4(2^{2n+1})$, ${}^3D_4(q)$; $E_7(p^t)$, where $p\neq 3$; $E_8(p^t)$, where $p\neq 3,5$, $D_{2\ell}(p^t)$, ${}^3D_4(p^t)$, ${}^2D_{2\ell}(p^t)$, where t is even if $p=3$ and, more over, if $G=D_4(p^t)$, then $|(\mathrm{Field}(G)\cap A):(\widehat{G}\cap A)|_{2'}>1$;*

(b) *A is one of the following groups: $B_\ell(3^t)$, $D_{2\ell}(3^t)$, ${}^2D_{2\ell}(3^t)$, $D_{2\ell+1}(p^t)$, ${}^2D_{2\ell+1}(r^t)$, ${}^3D_4(3^t)$, ${}^2G_2(3^{2n+1})$, $E_6^\varepsilon(r^t)$, $\widehat{E_6^\varepsilon(r^t)}$, $E_7(3^t)$, $E_8(3^t)$, $E_8(5^t)$, $C_\ell(3^t)$;*

In particular, no simple group, can be a minimal counter example. More over, if each almost simple group with known simple normal subgroup satisfies **(C)**, *then in all above mentioned groups a Carter subgroup (if exists) contains a Sylow 2-subgroup.*

Proof. (a) We claim that every element $z\in\widehat{G}$ of prime odd order is conjugate, under G, to some power $z^k\neq z$. When G is alternating or sporadic this is true by Lemma 3.2.11, and when G is of Lie type and z is semisimple, this is true by Lemma 3.2.3. On the other hand, when z is unipotent (hence p is odd), our claim follows from Lemmas 3.2.6, 3.2.7 if $G=G_2(3^t)$, $F_4(3^t)$ and from Lemma 3.2.5 in the remaining cases. Finally, if $G\simeq {}^3D_4(q)$, then by [44, Theorem 1.2(vi)] each element of G is conjugate to its inverse. Thus (a) follows from Lemma 3.3.1, since for all groups under consideration we have either that $|\widehat{G}:G|$ is a power of 2, and so by Lemma 4.2.6 all complements of odd order are conjugate, or that $|A:A\cap\widehat{G}|$ is a power of 2 (see [21], for example).

(b) Our statement follows from the results obtained in [27] and Theorem 1.5.6, when $G=B_2(3^t)\simeq C_2(3^t)$ or $G=C_\ell(3^t)$, and from Theorems 3.3.3 and 3.3.4, when G is one of the groups $D_{2\ell+1}^\varepsilon(p^t)$, $E_6^\varepsilon(p^t)$ or $\widehat{E_6^\varepsilon(p^t)}$. So assume that we are in the remaining cases. Every semisimple element $z\in\widehat{G}$ of prime odd order is conjugate to some z^{-1} by Lemma 3.2.3. Thus, in characteristic 2 a Carter subgroup K of G can only be a Sylow 2-subgroup and, in odd characteristic, K can only have order 2^ap^b. If $G\neq {}^2G_2(3^{2n+1})$, then the assumptions of Lemma 3.2.9 are satisfied and, by using Lemma 3.2.10, we conclude that K is again a 2-group.

Now assume $G={}^2G_2(3^{2n+1})$ (here $n\geqslant 1$). Then $|K|=2^a3^b$. Since the normalizer of a Sylow 2-subgroup of G contains an element of order 7 (see [11]), then we obtain that $b>0$. By Lemma 1.5.4, K is contained in a proper parabolic subgroup P of G. Since Lie rank of G equal 1, then P is a Borel subgroup, i. e. $P=U\rtimes H$, where H is a Cartan subgroup and U is a maximal unipotent subgroup of G. Since P is solvable, it satisfies **(C)** and by Lemma 2.4.1, KU/U is a Carter subgroup of $P/U\simeq H$. But for $n\geqslant 1$ the subgroup H contains an element of odd order, so K contains an element of prime odd order. A contradiction with $|K|=2^a3^b$. ☐

Note that after proving the statement that for every known finite simple group S and a nilpotent subgroup $N \leq \mathrm{Aut}(S)$, Carter subgroups of $\langle N, S \rangle$ are conjugate, Theorem 3.3.5 would imply that Carter subgroups in the groups mentioned in the theorem should contain a Sylow 2-subgroup. By Lemma 2.4.3 this is possible only if the normalizer of a Sylow 2-subgroup Q in A satisfies $N_A(Q) = QC_A(Q)$, i. e. if and only if A satisfies **(ESyl2)**. In [9] and subsequent results of the present paper, simple groups satisfying **(ESyl2)** are completely determined. More over Lemmas 4.3.1 and 4.3.3 allow to "lift" the property **(ESyl2)** from a simple group to an almost simple group. Thus a complete classification of Carter subgroups in groups mentioned in Theorem 3.3.5 is known.

4 Semilinear groups of Lie type

In this section we shall give a definition of semilinear groups of Lie type and generalize results about the structure of finite groups of Lie type for them. We need this theory to finding Carter subgroups in extensions of groups of Lie type by field, graph, or graph-field automorphisms in section 5. In the last subsection of this paragraph we shall consider the existence of Carter subgroup in semilinear groups, either containing a Sylow 2-subgroup, or contained in the normalizer of a Borel subgroup.

4.1 Basic definitions

Now we define some overgroups of finite groups of Lie type. We first give a more detailed description of a Frobenius map σ. Note that all maps in this section are automorphisms, if $\overline{G}$ is considered as an abstract group, and they are endomorphisms, if it is considered as an algebraic group. Since we use the maps to construct connected automorphisms of finite groups and groups over algebraically closed field, we find it appropriate to call all maps in this section by automorphisms. Let $\overline{G}$ be a connected simple linear algebraic group of adjoint type over the algebraic closure $\overline{\mathbb{F}}_p$ of a finite field of positive characteristic p. Below, if we do not say opposite, we shall consider groups of adjoint type. Choose a Borel subgroup $\overline{B}$ of $\overline{G}$, let $\overline{U} = R_u(\overline{B})$ be the unipotent radical of $\overline{B}$. There exists a Borel subgroup $\overline{B}^-$, satisfying $\overline{B} \cap \overline{B}^- = \overline{T}$, where $\overline{T}$ is a maximal torus of $\overline{B}$ (hence of $\overline{G}$). We partially duplicate the notations and the definitions of subsection 1.3 here. Let Φ be the root system of $\overline{G}$ and let $\{X_r \mid r \in \Phi^+\}$ be the set of $\overline{T}$-invariant 1-dimensional subgroups of $\overline{U}$. Each X_r is isomorphic to the additive group of $\overline{\mathbb{F}}_p$, so each element of X_r can be written as $x_r(t)$, where t is the image of $x_r(t)$ under this isomorphism. Denote by $\overline{U}^- = R_u(\overline{B}^-)$ the unipotent radical of $\overline{B}^-$. Define like above $\overline{T}$-invariant 1-dimensional subgroups $\{\overline{X}_r \mid r \in \Phi^-\}$ of $\overline{U}^-$. Then $\overline{G} = \langle \overline{U}, \overline{U}^- \rangle$. Let $\bar{\varphi}$ be a field automorphism of $\overline{G}$ (as an abstract group) and $\bar{\gamma}$ be a graph automorphism of $\overline{G}$. An automorphism $\bar{\varphi}$ is known to may be chosen so that it acts by $x_r(t)^{\bar{\varphi}} = x_r(t^p)$ (see [15, 12.2] and [19, 1.7], for example). In view of [15, Propositions 12.2.3 and 12.3.3], we may choose $\bar{\gamma}$ so that it acts by $x_r(t)^{\bar{\gamma}} = x_{\bar{r}}(t)$, if Φ has no roots of distinct length, and by $x_r(t)^{\bar{\gamma}} = x_{\bar{r}}(t^{\lambda_r})$ for suitable $\lambda_r \in \{1, 2, 3\}$, if Φ has roots of distinct length. Recall that $\bar{r}$ is the image of r under the symmetry ρ (corresponding to $\bar{\gamma}$) of a root system Φ. In both cases we can write $x_r(t)^{\bar{\gamma}} = x_{\bar{r}}(t^{\lambda_r})$, where $\lambda_r \in \{1, 2, 3\}$. From this formulas it is evident that $\bar{\varphi} \cdot \bar{\gamma} = \bar{\gamma} \cdot \bar{\varphi}$. Let $n_r(t) = x_r(t)x_{-r}(-t^{-1})x_r(t)$ and $\overline{N} = \langle n_r(t) \mid r \in \Phi, t \in \overline{\mathbb{F}}_p^* \rangle$. Let $h_r(t) = n_r(t)n_r(-1)$ and $\overline{H} = \langle h_r(t) \mid r \in \Phi, t \in \overline{\mathbb{F}}_p^* \rangle$. By [15, Chapters 6 and 7], $\overline{H}$ is a maximal torus of $\overline{G}$, $\overline{N} = N_{\overline{G}}(\overline{H})$, and subgroups $\overline{X}_r$ are root subgroups with respect to $\overline{H}$. So we can substitute

$\overline{T}$ by $\overline{H}$ and assume that $\overline{T}$ is $\bar{\varphi}$- and $\bar{\gamma}$- invariant under our choice. More over, $\bar{\varphi}$ induces a trivial automorphism of $\overline{N}/\overline{H}$. Note that $\overline{H} \leq \overline{B} \cap \overline{B}^-$, therefore $\overline{H} = \overline{T}$.

An automorphism $\bar{\varphi}^k$ with $k \in \mathbb{N}$ is called a *classical Frobenius automorphism.* We shall call an automorphism σ by a *Frobenius automorphism*, if σ is conjugate under $\overline{G}$ with $\bar{\gamma}^\epsilon\bar{\varphi}^k, \epsilon \in \{0,1\}, k \in \mathbb{N}$. By Lang-Steinberg theorem (Lemma 1.5.3) it follows that for every $\bar{g} \in \overline{G}$ elements σ and $\sigma\bar{g}$ are conjugate under $\overline{G}$. Thus by [42, 11.6], we have that a Frobenius map, defined in subsection 1.4, coinsides with a Frobenius automorphism, defined here.

Now fix $\overline{G}$, $\bar{\varphi}$, $\bar{\gamma}$, and $\sigma = \bar{\gamma}^\epsilon\bar{\varphi}^k$; and assume that $|\bar{\gamma}| \leqslant 2$, i. e., we do not consider a triality automorphism of a group $\overline{G}$ with root system $\Phi(\overline{G}) = D_4$. Set $B = \overline{B}_\sigma$, $H = \overline{H}_\sigma$, and $U = \overline{U}_\sigma$. Since $\overline{B}, \overline{H}$, and $\overline{U}$ are $\bar{\varphi}$- and $\bar{\gamma}$- invariant, they give us respectively a Borel subgroup, a Cartan subgroup, and a maximal unipotent subgroup (a Sylow p-subgroup) of $\overline{G}_\sigma$ (for more details see [19, 1.7–1.9] or [31, Chapter 2]).

Assume that $\epsilon = 0$, i. e., $O^{p'}(\overline{G}_\sigma)$ is not twisted (is split). Then $U = \langle X_r \mid r \in \Phi^+\rangle$, where X_r is isomorphic to the additive group of $\mathbb{F}_{p^k} = \mathbb{F}_q$, and each element of X_r can be written as $x_r(t), t \in \mathbb{F}_q$. Set also $U^- = \overline{U}^-_\sigma$. As for U, we may write $U^- = \langle X_r \mid r \in \Phi^-\rangle$ and each element of X_r can be written as $x_r(t), t \in \mathbb{F}_q$. Now we can define an automorphism φ as a restriction of $\bar{\varphi}$ on $\overline{G}_\sigma$ and an automorphism γ as a restriction of $\bar{\gamma}$ on $\overline{G}_\sigma$. By definition the equalities $x_r(t)^\varphi = x_r(t^p)$ and $x_r(t)^\gamma = x_{\bar{r}}(t^{\lambda_r})$ hold for all $r \in \Phi$ (see the definition of $\bar{\gamma}$ above). Define an automorphism ζ of $\overline{G}_\sigma$ by $\zeta = \gamma^\varepsilon\varphi^\ell$, $\varphi^\ell \neq e$, $\varepsilon \in \{0,1\}$, and an automorphism $\bar{\zeta}$ of $\overline{G}$ by $\bar{\zeta} = \bar{\gamma}^\varepsilon \cdot \bar{\varphi}^\ell$. Choose a ζ-invariant subgroup G with $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$. Note that if the root system Φ of $\overline{G}$ is not equal to D_{2n}, then $\overline{G}_\sigma/(O^{p'}(\overline{G}_\sigma))$ is cyclic. Thus ro most groups and automorphisms, except groups of type D_{2n} over a field of odd characteristic, every subgroup G of $\overline{G}_\sigma$, with $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$, is γ- and φ- invariant. Define ΓG as a set of subgroups of type $\langle G, \zeta g\rangle \leq \overline{G}_\sigma \leftthreetimes \langle\zeta\rangle$, where $g \in \overline{G}_\sigma$, $\langle\zeta g\rangle \cap \overline{G}_\sigma \leq G$; and $\Gamma\overline{G}$ as a set of subgroups of type $\overline{G} \leftthreetimes \langle\bar{\zeta}\rangle$. Following [31, Definition 2.5.13], an automorphism ζ is called a *field* automorphism, if $\varepsilon = 0$, i. e., $\zeta = \varphi^\ell$ and is called a *graph-field* automorphism in the remaining cases (recall that we are assuming $\varphi^\ell \neq e$).

Now assume that $\epsilon = 1$, i. e., $O^{p'}(\overline{G}_\sigma)$ is twisted. Then $U = \overline{U}_\sigma$ и $U^- = \overline{U}^-_\sigma$. Define φ on $U^\pm$ as a restriction of $\bar{\varphi}$ on $U^\pm$. Since $O^{p'}(\overline{G}_\sigma) = \langle U^+, U^-\rangle$, we obtain an automorphism φ of $O^{p'}(\overline{G}_\sigma)$. Consider $\zeta = \varphi^\ell \neq e$, and let G be a ζ-invariant group with $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$. Then $\bar{\zeta} = \bar{\varphi}^\ell$ is an automorphism of $\overline{G}$. Define ΓG as a set of subgroups of type $\langle G, \zeta g\rangle \leq \overline{G}_\sigma \leftthreetimes \langle\zeta\rangle$, where $g \in \overline{G}_\sigma$, $\langle\zeta g\rangle \cap \overline{G}_\sigma \leq G$; and $\Gamma\overline{G}$ as a set of subgroups of type $\overline{G} \leftthreetimes \langle\bar{\zeta}\rangle$. Following [31, Definition 2.5.13], we say that ζ is a *field* automorphism, if $|\zeta|$ is not divisible by $|\gamma|$ (this definition is used also in the case, when $|\gamma| = 3$ and $\overline{G}_\sigma \simeq {}^3D_4(q)$), and that ζ is a *graph* automorphism in the remaining cases.

Groups from above defined set ΓG are called *semilinear finite groups of Lie type* (they are also called *semilinear canonical finite groups of Lie type*, if $G = O^{p'}(\overline{G}_\sigma)$), while groups from the set $\Gamma\overline{G}$ are called *semilinear algebraic groups.* Note that $\Gamma\overline{G}$ cannot be defined without ΓG, since we need to know that $\varphi^\ell \neq e$. If G is written in the notations from [15], i. e., $O^{p'}(G) = G = A_n(q)$ or $O^{p'}(G) = G = {}^2A_n(q)$ etc., then we shall write ΓG as $\Gamma A_n(q)$, $\Gamma^2 A_n(q)$ etc.

Consider $A \in \Gamma G$ and $x \in A \setminus G$. Then $x = \zeta^k y$ for some $k \in \mathbb{N}$ and $y \in \overline{G}_\sigma$. Define $\bar{x}$ to be equal to $\bar{\zeta}^k y$. Conversely, if $\bar{x} = \bar{\zeta}^k y$ for some $y \in \overline{G}_\sigma$, $\zeta^k \neq e$, and $\langle\zeta^k y\rangle \cap \overline{G}_\sigma \leq G$, define x to be equal to $\zeta^k y$. Note that we need not to assume that $\bar{x} \notin \overline{G}$, since $|\bar{\zeta}| = \infty$. If $x \in G$, set $\bar{x} = x$.

Lemma 4.1.1. *In above notations consider a subgroup X of G. An element x normalizes X*

if and only if $\bar{x}$ normalizes X as a subgroup of $\overline{G}$.

Proof. Since ζ is a restriction of $\bar{\zeta}$ on G, our statement is evident. $\square$

Let X_1 be a subgroup of $A \in \Gamma G$. Then X_1 is generated by a normal subgroup $X = X_1 \cap G$ and an element $x = \zeta^k y$. By Lemma 4.1.1, we may consider the subgroup $\overline{X}_1 = \langle \bar{x}, X\rangle$ of $\overline{G} \leftthreetimes \langle\bar{\zeta}\rangle$. Now we find in reasonable to explain, why we use so complicated notations and definitions. We have that order of ζ is always finite, but order of $\bar{\zeta}$ is always infinite. Thus, even if $Z(\overline{G})$ is trivial, we cannot consider $G \leftthreetimes \langle\bar{\zeta}\rangle$ as a subgroup of $\mathrm{Aut}(\overline{G})$. Therefore we need to define in a some way (one possible way is just given) the connection between elements from $\mathrm{Aut}(G)$ and elements from $\mathrm{Aut}(\overline{G})$, in order to use the machinery of linear algebraic groups.

Let $\overline{R}$ be a σ-stable maximal torus (respectively a reductive subgroup of maximal rank, a parabolic subgroup) of $\overline{G}$, and an element $y \in N_{\overline{G}\leftthreetimes\langle\bar{\zeta}\rangle}(\overline{R})$, is chosen so that there exists $x \in \langle G, \zeta g\rangle$ with $y = \bar{x}$. Then $R_1 = \langle x, \overline{R} \cap G\rangle$ is called a *maximal torus* (respectively a *reductive subgroup of maximal rank*, a *parabolic subgroup*) of $\langle G, \zeta g\rangle$.

4.2 Translation of basic results

Lemma 4.2.1. *Let $M = \langle x, X\rangle$, where $X = M \cap G \trianglelefteq M$ is a subgroup of $\langle G, \zeta g\rangle$ such that $O_p(X)$ is nontrivial. Then there exists a proper σ- and $\bar{x}$- invariant parabolic subgroup $\overline{P}$ of $\overline{G}$ such that $X \leq \overline{P}$ and $O_p(X) \leq R_u(\overline{P})$.*

Proof. Define $U_0 = O_p(X)$, $N_0 = N_{\overline{G}}(U_0)$ and by induction $U_i = U_0 R_u(N_{i-1})$ and $N_i = N_{\overline{G}}(U_i)$. Clearly U_i, N_i are $\bar{x}$- and σ- invariant for all i. By [12, Proposition 30.3], the chain of subgroups $N_0 \leq N_1 \leq \ldots \leq N_k \leq \ldots$ is finite and $\overline{P} = \cup_i N_i$ is a proper parabolic subgroup of $\overline{G}$. Clearly $\overline{P}$ is σ- and $\bar{x}$- invariant, $X \leq \overline{P}$ and $O_p(X) \leq R_u(\overline{P})$. $\square$

Lemma 4.2.2. *Let $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$ be a finite adjoint group of Lie type with a base field of characteristic p and order q. Assume also that $O^{p'}(G)$ is not isomorphic to ${}^2D_{2n}(q)$, ${}^3D_4(q)$, ${}^2B_2(2^{2n+1})$, ${}^2G_2(3^{2n+1})$, ${}^2F_4(2^{2n+1})$. Then there exists a maximal σ-stable torus $\overline{T}$ of $\overline{G}$ such that*

- (a) *$(N_{\overline{G}}(\overline{T})/\overline{T})_\sigma \simeq (N_{\overline{G}}(\overline{T}))_\sigma/(\overline{T}_\sigma) = N(\overline{G}_\sigma, \overline{T}_\sigma)/\overline{T}_\sigma \simeq W$, where W is the Weyl group of $\overline{G}$;*
- (b) *if r is an odd prime divisor of $q - (\varepsilon 1)$, where $\varepsilon = +$, if G is split and $\varepsilon = -$, if G is twisted, then $N(\overline{G}_\sigma, \overline{T}_\sigma)$ contains a Sylow r-subgroup $\overline{G}_\sigma$;*
- (c) *if r is a prime divisor of $q - (\varepsilon 1)$, and s is an element of order r of G such that $C_{\overline{G}}(s)$ is connected, then, up to conjugation by an element of G, an element s is contained in $T = \overline{T}_\sigma \cap G$;*

A torus $\overline{T}$ is unique, up to conjugation, in $O^{p'}(\overline{G}_\sigma)$ and $|\overline{T}_\sigma| = (q - \varepsilon 1)^n$, where n is a rank of $\overline{G}$.

Proof. Since for every maximal torus T of $\overline{G}_\sigma$ the equality $\overline{G}_\sigma = TO^{p'}(\overline{G}_\sigma)$ holds, without lost of generality we may assume that $G = \overline{G}_\sigma$. If G is split, then the lemma can be easily proven. In this case $\overline{T}$ is a maximal torus such that $\overline{T}_\sigma$ is a Cartan subgroup of $\overline{G}_\sigma$ (i. e. $\overline{T}$ is a maximal split torus) and (a) is evident. Point (b) follows from [32, (10.1)]. More over, from [32, (10.2)] it follows that order of $\overline{T}_\sigma$ is uniquely defined and is equal to $(q-1)^n$, where n is a rank of $\overline{G}$. By [1, F, §6] we have that each element of order r of $\overline{T}$ is contained in $\overline{G}_\sigma$. Now there exists $g \in \overline{G}$ such that $s^g \in \overline{T}$, hence $s^g \in G$. In view of connectedness of the centralizer of s, elements s and s^g are conjugate in $\overline{G}$ if and only if they are conjugate in G, so s and s^g are conjugate in G, whence (c). The information about classes of maximal tori, given in [1, G] and [16], implies that, up to conjugation by an element from G, there exists a unique torus $\overline{T}$ such that $|\overline{T}_\sigma| = (q-1)^n$.

Assume that $O^{p'}(G) \simeq {}^2A_n(q)$. Then $\overline{T}$ is a maximal torus such that $|\overline{T}_\sigma| = (q+1)^n$. Note that $\overline{T}_\sigma$ can be obtained from a maximal split torus by twisting by the element $w_0\sigma$. Direct calculations by using [19, Proposition 3.3.6] show that $N(\overline{G}_\sigma, \overline{T}_\sigma)/\overline{T}_\sigma$ is isomorphic to $W(\overline{G})$, which in turn is isomorphic to Sym_{n+1}. The uniqueness follows from [18, Proposition 8]. Point (b) follows from [32, (10.1)]. As to point (c) we show first that each element of order r from $\overline{T}$ is in G. Assume that t is an element of order r in $\overline{T}$ (recall that in this case r divides $q+1$). Let $\overline{H}$ be a σ-stable maximal split torus of $\overline{G}$. The torus $\overline{T}_\sigma$ is obtained from $\overline{H}$ by twisting by $w_0\sigma$, where $w_0 \in W(\overline{G})$ is a unique element mapping all positive roots into negatives and $\overline{T}_\sigma \simeq \overline{H}_{\sigma w_0}$. Let $r_1, \ldots, r_n$ be a set of fundamental roots of A_n. Then t as an element of $\overline{H}$, can be written as $h_{r_1}(\lambda_1) \cdot \ldots \cdot h_{r_n}(\lambda_n)$. Now for each i we have $\sigma w_0 : h_{r_i}(\lambda) \mapsto h_{-r_i}(\lambda^q) = h_{r_i}(\lambda^{-q})$, i. e., $t^{\sigma w_0} = t^{-q}$. Since r divides $q+1$, we obtain that $t^{q+1} = e$, i. e., $t = t^{-q}$. Hence $t^{\sigma w_0} = t$ and $t \in \overline{T}_\sigma$. Now as in nontwisted case, there exists $g \in \overline{G}$ such that $s^g \in \overline{T}$, therefore $s^g \in \overline{T}_\sigma$. In view of connectedness of $C_{\overline{G}}(s)$, elements s and s^g are conjugate in G.

For $O^{p'}(G) = {}^2D_{2n+1}(q)$ we take $\overline{T}$ to be equal to a unique (up to conjugation in G) maximal torus, which has order $|\overline{T}_\sigma|$ equals $(q+1)^{2n+1}$ (uniqueness follows from [18, Proposition 10]), and for $O^{p'}(G) = {}^2E_6(q)$ we take $\overline{T}$ to be equal to a unique (again up to conjugation in G) maximal torus, which has order $|\overline{T}_\sigma|$ equals $(q+1)^6$ (uniqueness follows from [24, Table 1, p. 128]). As in the case of $G = {}^2A_n(q)$ it is easy to show that $\overline{T}$ satisfies (a), (b), and (c) of the lemma. ☐

Lemma 4.2.3. *Let G be a finite group of Lie type and $\overline{G}$, σ are chosen so that $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$. Let s be a regular semisimple element of odd prime order r of G.*
Then $N_G(C_{\overline{G}}(s)) \neq C_G(s)$.

Proof. In view of [34, Proposition 2.10] we have that $C_{\overline{G}}(s)/C_{\overline{G}}(s)^0$ is isomorphic to a subgroup of $\Delta(\overline{G})$. Now, if the root system Φ of $\overline{G}$ is not equal to either A_n, or E_6, then $|\Delta(\Phi)|$ is a power of 2. Since $\Delta(\overline{G})$ is a quotient of $\Delta(\Phi(\overline{G}))$, then Lemma 3.2.1 implies that $C_{\overline{G}}(s) = C_{\overline{G}}(s)^0 = \overline{T}$ is a maximal torus and $C_G(s) = C_{\overline{G}}(s) \cap G = T$. Since $N_G(T) \geq N(G,T) \neq T$ we obtain the statement of the lemma in this case. Thus we may assume that either $\Phi = A_n$, or $\Phi = E_6$.

Assume first that $\Phi = A_n$, i. e., $O^{p'}(G) = A_n^\varepsilon(q)$, where $\varepsilon \in \{+,-\}$. Clearly $T = C_{\overline{G}}(s)^0 \cap G$ is a normal subgroup of $C_G(s)$, hence $C_G(s) \leq N(G,T)$. Assume that $N_G(C_{\overline{G}}(s)) = C_G(s)$. Then $C_G(s) = N_{N(G,T)}(C_G(s))$ and $C_G(s)/T$ is a self-normalizing subgroup of $N(G,T)/T$. As we noted above $C_G(s)/T$ is isomorphic to a subgroup of $\Delta(A_n)$, i. e., it is cyclic. By Lemma 3.2.1, we also have that $C_G(s)/T$ is an r-group, thus $C_G(s)/T = \langle x \rangle$ for some r-element $x \in N(G,T)/T$. Thus $\langle x \rangle$ is a Carter subgroup of $N(G,T)/T$. Now, in view of [19, Proposition 3.3.6], we have that $N(G,T)/T \simeq C_{W(\overline{G})}(y)$ for some $y \in W(\overline{G}) \simeq \mathrm{Sym}_{n+1}$. Clearly $C_{C_{W(\overline{G})}(y)}(x)$ contains y, thus y must be an r-element, otherwise $N_{C_{W(\overline{G})}(y)}(\langle x \rangle)$ would contain an element of order coprime to r, i. e., $N_{C_{W(\overline{G})}(y)}(\langle x \rangle) \neq \langle x \rangle$. A contradiction with the fact that $\langle x \rangle$ is a Carter subgroup of $C_{W(\overline{G})}(y)$.

Now let $y = \tau_1 \cdot \ldots$ be the decomposition of y into the product of independent cycles and $l_1, \ldots$ be the lengths of $\tau_1, \ldots$ respectively. Assume first that m_1 cycles has the same length l_1, m_2 cycles has the length l_2 etc. Let $m_0 = n+1-(l_1m_1 + \ldots + l_km_k)$. Then

$$C_{\mathrm{Sym}_{n+1}}(y) \simeq \big(Z_{l_1} \wr \mathrm{Sym}_{m_1}\big) \times \ldots \times \big(Z_{l_k} \wr \mathrm{Sym}_{m_k}\big) \times \mathrm{Sym}_{m_0},$$

where Z_{l_i} is a cyclic group of order l_i. If $m_j > 1$ for some $j \geqslant 0$, then there exists a normal subgroup N of $C_{\mathrm{Sym}_{n+1}}(y)$ such that $C_{\mathrm{Sym}_{n+1}}(y)/N \simeq \mathrm{Sym}_{m_j} \neq \{e\}$. By Lemma 3.3.2, Carter subgroup in a group S satisfying $\mathrm{Alt}_\ell \leq S \leq \mathrm{Sym}_\ell$ are conjugate for all $\ell \geqslant 1$. Thus $C_{W(\overline{G})}(y)$ and N satisfy **(C)** and $\langle x\rangle$ is the unique, up to conjugation, Carter subgroup of $C_{W(\overline{G})}(y)$. By Lemma 2.4.1 we obtain that $\langle x\rangle$ maps onto a Carter subgroup of $C_{W(\overline{G})}(y)/N \simeq \mathrm{Sym}_{m_j}$. By Lemma 3.3.2 only a Sylow 2-subgroup of Sym_{m_j} can be a Carter subgroup of Sym_{m_j}. A contradiction with the fact that x is an r-element and r is odd.

Thus we may assume that $C_{W(\overline{G})}(y) \simeq (Z_{l_1} \times \ldots \times Z_{l_k})$ and $l_i \neq l_j$ if $i \neq j$. From the known structure of maximal tori and their normalizers in $A_n^\varepsilon(q)$ (see [18, Propositions 7,8], for example) we obtain the structure of T and $N(G,T)$, which we explain by using matrices. Below a group $\mathrm{GL}_n^\varepsilon(q)$ is isomorphic to $\mathrm{GL}_n(q)$ if $\varepsilon = +$ and is isomorphic to $\mathrm{GU}_n(q)$ if $\varepsilon = -$. For the decomposition $l_1 + l_2 + \ldots + l_k = n+1$ in $\mathrm{GL}_{n+1}^\varepsilon(q)$ consider a subgroup L, consisting of block-diagonal matrices of view

$$\begin{pmatrix} A_1 & 0 & \ldots & 0 \\ 0 & A_1 & \ldots & 0 \\ \ldots & \ldots & \ldots & \ldots \\ 0 & 0 & \ldots & A_k \end{pmatrix},$$

where $A_i \in \mathrm{GL}_{l_i}^\varepsilon(q)$. Then $L \simeq \mathrm{GL}_{l_1}^\varepsilon(q) \times \ldots \times \mathrm{GL}_{l_k}^\varepsilon(q)$. Denote, for brevity, $\mathrm{GL}_{l_i}^\varepsilon(q)$ by G_i. In every group G_i consider a Singer cycle T_i. $N_{G_i}(T_i)/T_i$ is known to be a cyclic group of order l_i and $N(G_i,T_i) = N_{G_i}(T_i)$. There exists a subgroup Z of $Z(\mathrm{SL}_{n+1}^\varepsilon(q))$ such that $O^{p'}(G) \simeq \mathrm{SL}_{n+1}^\varepsilon(q)/Z$. Then $T \simeq \big((T_1 \times \ldots \times T_k) \cap \mathrm{SL}_{n+1}^\varepsilon(q)\big)/Z$ and $N(G,T) \simeq \big((N(G_1,T_1) \times \ldots \times N(G_k,T_k)) \cap \mathrm{SL}_{n+1}^\varepsilon(q)\big)/Z$. Since for every Singer cycle T_i the group $N(G_i,T_i)/T_i$ is cyclic, we may assume that $N(G,T) = C_G(s)$ and T is a Singer cycle, i. e., is a cyclic group of order $\frac{q^{n+1}-(\varepsilon 1)^{n+1}}{q-(\varepsilon 1)}$ and $n+1 = r^k$ for some $k \geqslant 1$ (the last equality holds, since $N(G,T)/T$ is an r-group). But $q^{r^k} \equiv q \pmod r$, hence, r divides $q - (\varepsilon 1)$. By Lemma 4.2.2 we obtain that s is in $N(G,H)$, where H is a maximal torus such that the factor group $N(G,H)/H$ is isomorphic to Sym_{n+1} and $|H| = (q-\varepsilon 1)^n$. In particular, H is not a Singer cycle. If $s \in H$, this immediately implies a contradiction with the choice of s. If $s \notin H$, then, since the order of s is prime, the intersection $\langle s\rangle \cap H$ is trivial. Hence, under the natural homomorphism $N(G,H) \to N(G,H)/H \simeq \mathrm{Sym}_{n+1}$ the element s maps on an element of order r. But in Sym_{n+1} every element of odd order is conjugate to its inverse. Thus, there exists a 2-element z of G, which normalizes, but not centralizes $\langle s\rangle$. Therefore, $z \leq N_{\overline{G}}(C_{\overline{G}}(s)) \leq N_{\overline{G}}(C_{\overline{G}}(s)^0)$ and $|N(G,T)/T|$ is divisible by 2, that contradicts the above proven statement that $N(G,T)/T$ is an r-group. This final contradiction finish the case $\Phi(\overline{G}) = A_n$.

In the remaining case $\Phi = E_6$ it is easy to see, that for every $y \in W(E_6)$, the group $C_{W(E_6)}(y)$ does not contain Carter subgroup of order 3. Indeed, if $C_{W(E_6)}(y)$ has a Carter subgroup of order 3, then it is generated by y. But it is known (and can be easily checked by using [16, Table 9]), that in $W(E_6)$ there is no elements of order 3, which centralizer has order 3. Since $|C_G(s)/T|$ divides 3 and the group $C_G(s)/T$ is a Carter subgroup of $C_{W(E_6)}(y)$ for some y, we get a contradiction. □

By using Lemma 4.2.3 we can obtain a similar result for semilinear groups.

LEMMA 4.2.4. *Let $\langle G, \zeta g\rangle$ be a finite semilinear group of Lie type and $\overline{G}$, σ are chosen so that $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$. Let s be a regular semisimple element of odd order of G. Then $N_{\langle G,\zeta g\rangle}(C_{\langle G,\zeta g\rangle}(s)) \neq C_{\langle G,\zeta g\rangle}(s)$.*

Proof. Since s is semisimple, there exists σ-stable maximal torus $\overline{S}$ of $\overline{G}$ containing s. Since $\overline{G}_\sigma = O^{p'}(\overline{G}_\sigma)\overline{S}_\sigma$ we may assume that $g \in \overline{S}_\sigma$, i. e. elements g and s commutes. If $C_{\langle G,\zeta g\rangle}(s)G \neq \langle G,\zeta g\rangle$, then we can substitute $\langle G,\zeta g\rangle$ by $C_{\langle G,\zeta g\rangle}(s)G$ and prove the lemma for this group. Moreover, if $C_{\langle G,\zeta g\rangle}(s) = C_G(s)$, then the lemma follows from Lemma 4.2.3, so we may assume that ζ centralizes s. If either G is not twisted, or $|\zeta|$ is odd, then by [31, Proposition 2.5.17] it follows that we may assume $\sigma = \bar{\zeta}^k$ for some $k > 0$. By Lemma 4.2.3 there exists an element of $N_{G_{\zeta g}}(C_{\overline{G}}(s))$, not contained in $C_{G_{\zeta g}}(s)$, and the lemma follows.

Assume that G is twisted and $|\zeta|$ is even. Then $\sigma = \bar{\gamma}\bar{\varphi}^k$, $\bar{\zeta} = \bar{\varphi}^\ell$, where k divides ℓ. Therefore s is in $\overline{G}_{\bar{\gamma}}$. Depending on the root system $\Phi(\overline{G})$, we obtain that $\overline{G}_{\bar{\gamma}}$ is isomorphic to a simple algebraic group with root system equal to B_m (for some $m > 1$), C_m (for some $m > 2$), or F_4. By Lemma 3.2.3 the element s is conjugate with its inverse under $O^{p'}((\overline{G}_{\bar{\gamma}})_{\sigma\bar{\zeta}g}) \leq G_{\zeta g}$, so $N_{\langle G,\zeta g\rangle}(C_{\langle G,\zeta g\rangle}(s)) \neq C_{\langle G,\zeta g\rangle}(s)$. □

Lemma 4.2.5. *Let G be a finite group of Lie type over a field of odd characteristic p. Assume that $\overline{G}$ and σ are chosen so that $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$. Let ψ be a field automorphism of odd order of $O^{p'}(\overline{G}_\sigma)$.*

Then ψ centralizes a Sylow 2-subgroup of G, and there exists a ψ-stable Cartan subgroup H such that ψ centralizes a Sylow 2-subgroup of H. Moreover, if $G \not\simeq {}^2G_2(3^{2n+1})$, ${}^3D_4(q^3)$, ${}^2D_{2n}(q^2)$, then there exists a ψ-stable torus T of G such that ψ centralizes a Sylow 2-subgroup of T and the factor group $N(G,T)/T$ is isomorphic to $N_{\overline{G}}(\overline{T})/\overline{T}$.

Proof. Clearly we need to prove the lemma only for the case $G = \overline{G}_\sigma$. Assume that $|\psi| = k$. Let $\mathbb{F}_q$ be the base field of G. Then $q = p^\alpha$ and $\alpha = k \cdot m$. Now $|G|$ can be written as $|G| = q^N(q^{m_1} + \varepsilon_1 1) \cdot \ldots \cdot (q^{m_n} + \varepsilon_n 1)$ for some N, where n is the rank of G, $\varepsilon_i = \pm$ (see [15, Theorems 9.4.10 and 14.3.1]). Similarly we have that $|G_\psi| = (p^m)^N((p^m)^{m_1} + \varepsilon_1 1) \cdot \ldots \cdot ((p^m)^{m_n} + \varepsilon_n 1)$, i. e., $|G|_2 = |G_\psi|_2$ and a Sylow 2-subgroup of G_ψ is a Sylow 2-subgroup of G. By [31, Proposition 2.5.17] there exists an automorphism ψ_1 of $\overline{G}$ such that $\sigma = \psi_1^k$ and ψ coincides with the restriction of ψ_1 on $\overline{G}_\sigma$. Note that ψ_1, in general, is not equal to $\bar{\psi}$ defined above. Consider a maximal split torus $\overline{H}_{\psi_1}$ of $\overline{G}_{\psi_1}$. Then $H = \overline{H}_\sigma$ is a ψ-stable Cartan subgroup of G. Since $|H| = (q^{k_1} + \varepsilon 1) \cdot \ldots \cdot (q^{k_l} + \varepsilon_l 1)$, where $\varepsilon_i = \pm$, the equality $|H|_2 = |H_\psi|_2$ can be proven in the same way.

Now assume that $G \not\simeq {}^2G_2(3^{2n+1})$, ${}^3D_4(q^3)$, ${}^2D_{2n}(q^2)$. By Lemma 4.2.2, there exists a maximal torus T of G_ψ such that $N(G_\psi,T)/T \simeq N_{\overline{G}}(\overline{T})/\overline{T}$ and $|T_\psi| = (p^m - \varepsilon 1)^n$. Since $|\psi|$ is odd and $\overline{T}_{\psi_1}$ is obtained from a maximal splittorus $\overline{H}$ by twisting by an element w_0, then $\overline{T}_\sigma$ is also obtained from a maximal split torus $\overline{H}$ by twisting by element w_0 (see proof of Lemma 4.3.1). Therefore $|\overline{T}_\sigma| = (q - \varepsilon 1)^n$, $|\overline{T}_{\psi_1}| = (p^m - \varepsilon 1)^n$, hence $|\overline{T}_\sigma|_2 = |T|_2 = |T_\psi|_2$. □

Lemma 4.2.6. [32, (7-2)] *Let $\overline{G}$ be a connected simple linear algebraic group over a field of characteristic p, σ be a Frobenius map of $\overline{G}$ and $G = \overline{G}_\sigma$ be a finite group of Lie type. Let φ be a field or a graph-field automorphism of G (if φ is graph-field, then corresponding graph automorphism has order 2) and let φ' be an element of $(G \leftthreetimes \langle\varphi\rangle) \setminus G$ such that $|\varphi'| = |\varphi|$ and $G \leftthreetimes \langle\varphi\rangle = G \leftthreetimes \langle\varphi'\rangle$.*
Then there exists $g \in G$ such that $\langle\varphi\rangle^g = \langle\varphi'\rangle$. In particular, if $G/O^{p'}(G)$ is a 2-group and φ is of odd order, then g can be chosen in $O^{p'}(G)$.

A particular case of the following lemma is proven in [30, Theorem A]).

Lemma 4.2.7. *Let G be a finite adjoint split group of Lie type, $\overline{G}$, σ are chosen so that $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$. Assume that τ is a graph automorphism of order 2 of $O^{p'}(G)$.*

Then every semisimple element $s \in G$ *is conjugate to its inverse under* $\langle O^{p'}(\overline{G}_\sigma), \tau a\rangle$, *where* a *is an element of* $\overline{G}_\sigma$.

Proof. If $\Phi(\overline{G})$ is not of type A_n, D_{2n+1}, E_6, then the lemma follows from Lemma 3.2.3, thus we need to consider groups of type A_n, D_{2n+1}, E_6. Denote by $\bar{\tau}$ the graph automorphism of $\overline{G}$ such that $\bar{\tau}|_G = \tau$. Let $\overline{T}$ be a maximal σ-stable torus of $\overline{G}$ such that $\overline{T}_\sigma \cap G$ is a Cartan subgroup of G. Let $r_1, \ldots, r_n$ be fundamental roots of $\Phi(\overline{G})$ and ρ be the symmetry corresponding to $\bar{\tau}$. Denote r_i^ρ by $\bar{r}_i$. Then $\overline{T} = \langle h_{r_i}(t_i) \mid 1 \leqslant i \leqslant n,\ t_i \neq 0\rangle$ and $h_{r_i}(t_i)^{\bar{\tau}} = h_{\bar{r}_i}(t_i)$. Denote by W the Weyl group of $\overline{G}$. Let w_0 be the unique element of W mapping all positive roots onto negative roots and let n_0 be its preimage in $N_{\overline{G}}(\overline{T})$ under the natural homomorphism $N_{\overline{G}}(\overline{T}) \to N_{\overline{G}}(\overline{T})/\overline{T} \simeq W$. Since σ acts trivially on $W = N(G,T)/T$ (see Lemma 4.2.2), we can take $n_0 \in G$, i. e., $n_0^\sigma = n_0$. Then for all r_i and t we have that

$$h_{r_i}(t)^{n_0\bar{\tau}} = h_{r_i^{w_0\rho}}(t) = h_{-r_i}(t) = h_{r_i}(t^{-1}).$$

Thus $x^{n_0\bar{\tau}} = x^{-1}$ for all $x \in \overline{T}$.

Now let s be a semisimple element of G. Then there exists a maximal σ-stable torus $\overline{S}$ of $\overline{G}$ containing s. Since all maximal tori of $\overline{G}$ are conjugate, we have that there exists $g \in \overline{G}$ such that $\overline{S}^g = \overline{T}$. Since $\overline{G}_\sigma = O^{p'}(\overline{G}_\sigma)\overline{T}_\sigma$, then we may assume that $a \in \overline{T}_\sigma$. Therefore $s^{gn_0\bar{\tau}ag^{-1}} = s^{-1}$. Since $n_0^\sigma = n_0$ and $\bar{\tau}^\sigma = \bar{\tau}$ we have that $(gn_0\bar{\tau}ag^{-1})^\sigma = g^\sigma n_0\bar{\tau}a(g^{-1})^\sigma$. Moreover, since $\overline{S}$ is σ-stable, then for every $x \in \overline{S}$ we have that $x^{gn_0\bar{\tau}ag^{-1}} = x^{g^\sigma n_0\bar{\tau}a(g^{-1})^\sigma} = x^{-1}$, i. e., $gn_0\bar{\tau}ag^{-1}\overline{S} = g^\sigma n_0\bar{\tau}a(g^{-1})^\sigma\overline{S}$. In particular, there exists $t \in \overline{S}$ such that $gn_0\bar{\tau}ag^{-1}t = g^\sigma n_0\bar{\tau}a(g^{-1})^\sigma$. In view of Lang-Steinberg Theorem (Lemma 1.5.3) there exists $y \in \overline{S}$ such that $t = y \cdot (y^{-1})^\sigma$. Therefore, $gn_0\bar{\tau}ag^{-1}y = (gn_0\bar{\tau}ag^{-1}y)^\sigma$, i. e., $gn_0\tau ag^{-1}y \in \overline{G}_\sigma \leftthreetimes \langle\tau\rangle$, and $s^{gn_0\tau ag^{-1}y} = s^{-1}$. Since $O^{p'}(\overline{G}_\sigma)\overline{S}_\sigma = \overline{G}_\sigma$, and $\overline{S}_\sigma$ is Abelian, we may find $z \in \overline{S}_\sigma$ such that $gn_0\tau ag^{-1}yz \in \langle O^{p'}(\overline{G}_\sigma), \tau a\rangle$. □

4.3 Carter subgroups of special type

In this subsection we consider problems of structure and existence of Carter subgroups in semilinear groups, containing a Sylow 2-subgroup or is contained in the normalizer of a Borel subgroup.

Lemma 4.3.1. *Let* G *be a finite group of Lie type over a field of odd characteristic and* $\overline{G}$, σ *are chosen so that* $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$. *If* G *satisfies* **(ESyl2)**, *then every subgroup* L *with* $G \leq L \leq O^{p'}(\overline{G}_\sigma)$ *satisfies* **(ESyl2)**.

Proof. Let Q be a Sylow 2-subgroup of $\overline{G}_\sigma$ and $Q^0 = O^{p'}(\overline{G}_\sigma) \cap Q$ be a Sylow 2-subgroup of $O^{p'}(\overline{G}_\sigma)$. If $N_{\overline{G}_\sigma}(Q^0) = QC_{\overline{G}_\sigma}(Q)$, then the statement of the lemma is clearly true. In view of [9, Theorem 1], for a classical group $\overline{G}_\sigma$ the equality $N_{\overline{G}_\sigma}(Q^0) = QC_{\overline{G}_\sigma}(Q)$ can fail to be true only if the root system of $\overline{G}$ has type A_1 or C_n. If the root system of $\overline{G}$ has type A_1 or C_n, then $|\overline{G}_\sigma : O^{p'}(\overline{G}_\sigma)| = 2$ and the statement of the lemma follows from Lemma 2.4.6.

Assume now that G is a group of exceptional type. If $\overline{G}_\sigma = O^{p'}(\overline{G}_\sigma)$, then the statement of the lemma is clearly true. The equality $N_{\overline{G}_\sigma}(Q^0) = QC_{\overline{G}_\sigma}(Q)$ might fail to be true only if the root system of $\overline{G}$ has type E_6 or E_7. If the root system of $\overline{G}$ has type E_7, then $|\overline{G}_\sigma : O^{p'}(\overline{G}_\sigma)| = 2$ and the statement of the lemma follows from Lemma 2.4.6.

Assume that the root system of $\overline{G}$ has type E_6. Then either $\overline{G}_\sigma = O^{p'}(\overline{G}_\sigma)$ or $|\overline{G}_\sigma : O^{p'}(\overline{G}_\sigma)| = 3$. In the first case we have nothing to prove, so assume that $|\overline{G}_\sigma : O^{p'}(\overline{G}_\sigma)| = 3$.

Since the group G coincides either with $\overline{G}_\sigma$, or with $O^{p'}(\overline{G}_\sigma)$, and since in case $G=\overline{G}_\sigma$ there is nothing to prove, we may assume that $G=O^{p'}(\overline{G}_\sigma)$. By [31, Theorem 4.10.2] there exists a maximal torus T of $\overline{G}_\sigma$ such that Q is contained in $N(\overline{G}_\sigma,T)$. Since $|\overline{G}_\sigma:G|=3$, then $Q=Q^0\leq N(G,T\cap G)$. By [10, Theorem 6] the equality $N_G(Q)=Q\times R^0$ holds, where $R^0\leq T$ is a cyclic group of odd order. Now since $\overline{G}_\sigma=TG$, then $N_{\overline{G}_\sigma}(Q)=\langle N_T(Q),N_G(Q)\rangle$. Indeed, $N(G,T\cap G)/(T\cap G)\simeq N(G,T)/T$. Hence, a Sylow 2-subgroup QT/T of $N(G,T)/T$ coincides with its normalizer. Since the factor group $\overline{G}_\sigma/G$ is cyclic of order 3, then $N_{\overline{G}_\sigma}(Q)=\langle tg,N_G(Q)\rangle$, where $t\in T$ and $g\in G$. Moreover, since $|\overline{G}_\sigma:G|=3$, we may assume that tg is an element of order 3^k for some $k>0$. Since $t\in T\leq N(\overline{G}_\sigma,T)$, then $Q^t\leq N(G,T\cap G)$. So there exists an element $g_1\in N(G,T\cap G)$ such that $Q^t=Q^{g_1^{-1}}$. Therefore we may assume that $tg=tg_1\in N(\overline{G}_\sigma,T)$. Under the natural epimorphism $\pi:N(\overline{G}_\sigma,T)\rightarrow N(\overline{G}_\sigma,T)/T$ the image of $N_{N(\overline{G}_\sigma,T)}(Q)$ coincides with Q. Hence, $(tg)^\pi=e$, so $tg\in T$. Thus each element of odd order of $\overline{G}_\sigma$ normalizing Q lies in T. Since T is a torus, then T is Abelian, hence the set of elements of odd order of $N_{\overline{G}_\sigma}(Q)$ forms a normal subgroup $R\leq T$. Therefore $N_{\overline{G}_\sigma}(Q)=Q\times R$, i. e., $\overline{G}_\sigma$ satisfies **(ESyl2)**. □

The following lemma is immediate from [9, теорема 1].

Lemma 4.3.2. *Let $O^{p'}(\overline{G}_\sigma)=G$ be a canonical finite group of Lie type and $\overline{G}$ is either of type A_1 or of type C_n, p is odd, $q=p^\alpha$ is the order of the base field of G. Then G satisfies* **(ESyl2)** *if and only if $q\equiv\pm1\pmod 8$.*

Note that Lemma 4.3.1 together with [9, Theorem 1] and [10, Theorem 6] implies that every group of Lie type over a field of odd characteristic, distinct from a Ree group and groups from Lemma 4.3.2, satisfies **(ESyl2)**.

Lemma 4.3.3. *Let G be a finite adjoint group of Lie type over a field of odd characteristic, $G\not\simeq {}^3D_4(q^3)$, and $\overline{G}$, σ are chosen so that $O^{p'}(\overline{G}_\sigma)\leq G\leq\overline{G}_\sigma$. Let A be a subgroup of $\mathrm{Aut}(O^{p'}(\overline{G}_\sigma))$ such that $A\cap\overline{G}_\sigma=G$. If $O^{p'}(G)\simeq D_4(q)$, assume also that A is contained in the group generated by inner-diagonal, field automorphisms and a graph automorphism of order 2. Then A satisfies* **(ESyl2)** *if and only if G satisfies* **(ESyl2)**.

Proof. Assume that G satisfies **(ESyl2)**. In the conditions of the lemma we have that the factor group A/G is Abelian, so $A/G=\overline{A}_1\times\overline{A}_2$, where $\overline{A}_1$ is a Hall $2'$-subgroup of A/G and $\overline{A}_2$ is a Sylow 2-subgroup of A/G. Denote by A_1 the complete preimage of $\overline{A}_1$ in A. If A_1 satisfies **(ESyl2)**, then by Lemma 2.4.6 A satisfies **(ESyl2)** as well. Thus we may assume that the order $|A/G|$ is odd. Since we are assuming that a graph automorphism of order 3 is not contained in A, then A/G is cyclic, hence $A=\langle G,\psi g\rangle$, where ψ is a field automorphism of odd order and $g\in\overline{G}_\sigma$. Since $|A:G|=|\psi|$ is odd, we may assume that $|\psi g|$ is also odd. By Lemma 4.2.5, ψ centralizes a Sylow 2-subgroup of $\overline{G}_\sigma$, therefore g is of odd order. Now the quotient $\overline{G}_\sigma/G$ is Abelian and can be written as $\overline{L}\times\overline{Q}$, where $\overline{L}$ is a Hall $2'$-subgroup of $\overline{G}_\sigma/G$ and $\overline{Q}$ is a Sylow 2-subgroup of $\overline{G}_\sigma/G$. Let L be the complete preimage of $\overline{L}$ in $\overline{G}_\sigma$ under the natural homomorphism. Then $g\in L$. Consider $L\leftthreetimes\langle\psi\rangle\geq A$. By construction, $|L\leftthreetimes\langle\psi\rangle:A|=|L:G|$ is odd. By Lemma 4.3.1 the group L satisfies **(ESyl2)**. By Lemma 4.2.5 the field automorphism ψ centralizes a Sylow 2-subgroup Q of L, Thus

$$N_{L\leftthreetimes\langle\psi\rangle}(Q)=N_L(Q)\times\langle\psi\rangle=QC_L(Q)\times\langle\psi\rangle=QC_{L\leftthreetimes\langle\psi\rangle}(Q),$$

i. e., the group $L\leftthreetimes\langle\psi\rangle$ satisfies **(ESyl2)**. Since $|L\leftthreetimes\langle\psi\rangle:A|$ is odd, then A satisfies **(ESyl2)**.

Now assume that A satisfies **(ESyl2)**. If G does not satisfies **(ESyl2)**, then [9, Theorem 1] and [10, Theorem 6] imply that the root system of $\overline{G}$ either has type A_1, or has type C_n. In both cases the factor group $\mathrm{Aut}(O^{p'}(\overline{G}_\sigma)/\overline{G}_\sigma)$ is cyclic and is generated by a field automorphism φ. Further, from [9, Theorem 1] it follows that the order of the base field (that is equal to the field of definition in this case, since G is not twisted) is equal to $q = p^t$ and $q \equiv \pm 3 \pmod 8$. Therefore t is odd and, so $|\mathrm{Aut}(O^{p'}(\overline{G}_\sigma))/\overline{G}_\sigma|$ is odd. Thus $|A : G|$ is odd, hence G satisfies **(ESyl2)**. $\square$

Lemma 4.3.4. *Let $\langle G, \zeta g\rangle$ be a finite semilinear group of Lie type over a field of characteristic p (we do not exclude the case $\langle G, \zeta g\rangle = G$) and G is of adjoint type (recall that $g \in \overline{G}_\sigma$, but not necessary $g \in G$). Assume that $B = U \leftthreetimes H$, where H is a Cartan subgroup of G, is a ζg-invariant Borel subgroup of G and $\langle B, \zeta g\rangle$ contains a Carter subgroup K of $\langle G, \zeta g\rangle$. Assume that $K \cap U \neq \{e\}$. Then one of the following statements holds:*

(a) *either $\langle G, \zeta g\rangle = \langle {}^2A_2(2^{2t}), \zeta g\rangle$, or $\langle G, \zeta g\rangle = {}^2\widehat{A_2(2^{2t})} \leftthreetimes \langle \zeta\rangle$; the order $|\zeta| = t$ is odd and is not divisible by 3, $C_G(\zeta) \simeq {}^2\widehat{A_2(2^2)}$, $K \cap G$ is Abelian and has order $2 \cdot 3$;*

(b) *$\langle G, \zeta g\rangle = \langle {}^2A_2(2^{2t})\zeta g\rangle$, the order $|\zeta| = t$ is odd, $C_G(\zeta) \simeq {}^2A_2(2^2)$, the subgroup $K \cap G$ is a Sylow 2-subgroup of G_ζ;*

(c) *either $\langle G, \zeta g\rangle = \langle A_2(2^{2t}), \zeta g\rangle$, or $\langle G, \zeta g\rangle = \widehat{A_2(2^{2t})} \leftthreetimes \langle \zeta\rangle$, ζ is a graph-field automorphism of order $2t$, t is not divisible by 3, and $C_G(\zeta) \simeq {}^2\widehat{A_2(2^2)}$, the subgroup $K \cap G$ is Abelian and has order $2^{|\zeta_{2'}|} \cdot 3$;*

(d) *$\langle G, \zeta g\rangle = \langle A_2(2^{2t}), \zeta g\rangle$, ζ is a graph-field automorphism and $C_G(\zeta) \simeq {}^2A_2(2^2)$, the subgroup $K \cap G$ is a Sylow 2-subgroup of $G_{\zeta_{2'}}$;*

(e) *G is defined over $\mathbb{F}_{2^t}$, $\langle G, \zeta g\rangle = G \leftthreetimes \langle \zeta g\rangle$, ζ is either a field automorphism of order t of $O^{2'}(G)$, if $O^{2'}(G)$ is split, or a graph automorphism of order t, if $O^{2'}(G)$ is twisted, and, up to conjugation in G, $K = Q \leftthreetimes \langle \zeta g\rangle$, where Q is a Sylow 2-subgroup of $G_{(\zeta g)_{2'}}$;*

(f) *G is split and defined over $\mathbb{F}_{2^t}$, $\langle G, \zeta g\rangle = G \leftthreetimes \langle \zeta g\rangle$, ζ is a product of a field automorphism of odd order t of $O^{2'}(G)$ and a graph automorphism of order 2, ζ and ζg are conjugate under $\overline{G}_\sigma$, and, up to a conjugation in G, $K = Q \leftthreetimes \langle \zeta g\rangle$, where Q is a Sylow 2-subgroup of $G_{(\zeta g)_{2'}}$;*

(g) *$G/Z(G) \simeq \mathbf{P}SL_2(3^t)$, the order $|\zeta| = t$ is odd (hence $\zeta \in \langle G, \zeta g\rangle$), and K contains a Sylow 3-subgroup of $G_{\zeta_{3'}}$;*

(h) *$\langle G, \zeta g\rangle = {}^2G_2(3^{2n+1}) \leftthreetimes \langle \zeta\rangle$, $|\zeta| = 2n+1$, $K \cap {}^2G_2(3^{2n+1}) = Q \times P$, where Q is of order 2 and $|P| = 3^{|\zeta|_3}$.*

Note that in all points (a)–(h) of the lemma Carter subgroups, having given structure, do exist. The existence of Carter subgroups in points (a) and (c) follows from the existence of a Carter subgroup of order 6 in $\mathbf{P}\mathrm{GU}_3(2)$ (see [27]). The existence of Carter subgroups in points (b), (d)-(f) follows from the fact that a Sylow 2-subgroup in a group of Lie type defined over a field of order 2, coincides with its normalizer. The existence of Carter subgroups in point (g) follows from the fact that a Sylow 3-subgroup of $\mathbf{P}\mathrm{SL}_2(3)$ coincides with its normalizer. The existence of a Carter subgroup, satisfying point (h) of the lemma, follows from the existence of a Carter subgroup K of order 6 in a (non simple) group ${}^2G_2(3)$. The existence of a Carter subgroup K of order 6 in ${}^2G_2(3)$ follows from the results given in [11] and [45].

Proof. If G is one of the groups $A_1(q)$, $G_2(q)$, $F_4(q)$, ${}^2B_2(2^{2n+1})$, or ${}^2F_4(2^{2n+1})$, then the lemma follows from Table 3. If $\langle G, \zeta g\rangle = G$, then the lemma follows from the results of section 3 and Theorem 1.5.6. So we may assume that $\langle G, \zeta g\rangle \neq G$, i. e., that ζ is a nontrivial field, graph-field, or graph automorphism. If $\Phi(\overline{G}) = C_n$, the lemma follows from Theorem 5.2.3 below, that does not use Lemma 4.3.4, so we assume that $\Phi(\overline{G}) \neq C_n$. If $\Phi(\overline{G}) = D_4$ and either a graph-field automorphism ζ is a product of a field automorphism and a graph automorphism of order 3, or $G \simeq {}^3D_4(q^3)$, then the lemma follows from Theorem 5.3.1 below, that does not use Lemma 4.3.4, so we assume that $\langle G, \zeta g\rangle$ is contained in the group A_1 defined in Theorem 5.3.1, and $G \not\simeq {}^3D_4(q^3)$. Since we shall use Lemma 4.3.4 in the proof of Theorem 5.4.1, after Theorems 5.2.3 and 5.3.1, it is possible to make such additional assumptions.

Assume that q is odd and $\Phi(\overline{G})$ is one of the following types: A_n $(n \geqslant 2)$, D_n $(n \geqslant 4)$, B_n $(n \geqslant 3)$, E_6, E_7 or E_8. By Lemma 2.4.1 we have that KU/U is a Carter subgroup of $\langle B, \zeta g\rangle/U \simeq \langle H, \zeta g\rangle$. Since $\overline{G}_\sigma = G\overline{H}_\sigma$, where $\overline{H}$ is a maximal split torus of $\overline{G}$ and $\overline{H}_\sigma \cap G = H$, then we may assume that $g \in \overline{H}_\sigma$, in particular g centralizes H. So $H_\zeta \leq Z(\langle H, \zeta g\rangle)$, and we obtain, up to conjugation in B, that $H_\zeta \leq K$. By Lemma 4.2.5, the automorphism $\zeta_{2'}$ centralizes a Sylow 2-subgroup Q of H. Thus, each element of odd order of $\langle H, \zeta g\rangle$ centralizes Q and Lemma 2.4.3 implies, that, up to conjugation in B, the inclusion $Q \leq K$ holds. By Lemma 3.2.9 it follows that $C_U(Q) = \{e\}$, a contradiction with the fact that $K \cap U$ is nontrivial.

Assume that $G \simeq {}^2G_2(3^{2n+1})$ and $\langle G, \zeta g\rangle = G \rtimes \langle\zeta\rangle$ (in this case $O^{p'}(\overline{G}_\sigma) = \overline{G}_\sigma$). Again by Lemma 2.4.1 we have that KU/U is a Carter subgroup of $(B \rtimes \langle\zeta\rangle)/U \simeq H \rtimes \langle\zeta\rangle$. By Lemma 3.2.3 every semisimple element of G is conjugate to its inverse. Since non-Abelian composition factors of every semisimple element of G can be isomorphic only to groups $A_1(q)$, by Table 3 it follows that the centralizer of every semisimple element of G satisfies condition **(C)**. So Lemma 2.4.2 implies that $KU/U \cap B/U$ is a 2-group. On the other hand, $|H|_2 = 2$ and $KU/U \geq Z(B/U) \geq H_\zeta$, hence $|H_\zeta| = 2$ and $|\zeta| = 2n+1$. Thus $K \cap G = (K \cap U) \times \langle t\rangle$, where t is an involution. It follows that $K \cap U = C_G(t) \cap G_{\zeta_{3'}}$. Now the structure results from [11, Theorem 1] and [45] imply point (h) of the lemma.

Assume now that $q = 2^t$. Assume first that $\Phi(\overline{G})$ has one of the types A_n $(n \geqslant 2)$, D_n $(n \geqslant 4)$, B_n $(n \geqslant 3)$, E_6, E_7 or E_8, G is split, and ζ is a field automorphism. Like above we obtain that $H_\zeta \leq K$, and $O^{2'}(G_\zeta)$ is a split group of Lie type with definition field of order $q = 2^{t/|\zeta|}$. By Hartley-Shute Lemma 1.5.5, for every $r \in \Phi(\overline{G})$ and for every $s \in GF(2^{t/|\zeta|})^*$ there exists $h(\chi) \in H_\zeta \cap O^{2'}(G_\zeta)$ such that $\chi(r) = s$. The same arguments as in Lemma 3.2.9 imply that if $\frac{t}{|\zeta|} \neq 1$, inequality $K \cap U \leq C_U(H_\zeta) = \{e\}$ holds, a contradiction. So $|\zeta| = t$ and $H_\zeta = \{e\}$. Since g can be chosen in $\overline{H}_\sigma$ and $\langle \zeta g\rangle \cap \overline{G}_\sigma \leq \langle\zeta g\rangle \cap \overline{H}_\sigma \leq H_\zeta = \{e\}$, then $\langle\zeta g\rangle \cap \overline{G}_\sigma = \{e\}$. By Lemma 4.2.6 elements ζg and ζ are conjugate under $\overline{G}_\sigma$, and point (e) of the lemma follows.

Now assume that $\Phi(\overline{G})$ is of type A_n $(n \geqslant 3)$, D_n $(n \geqslant 4)$, or E_6; and either ζ is a graph-field automorphism and G is split, or G is twisted. Let ρ be the symmetry of the Dynkin diagram of $\Phi(\overline{G})$ corresponding to γ (recall that $\zeta = \gamma^\varepsilon\varphi^\ell$ by definition), and $\bar{r}$ denotes r^ρ for $r \in \Phi(\overline{G})$. Like above it is possible to prove that, up to conjugation, $H_\zeta \leq K$. If $|\zeta| = 2t$, then $H_\zeta \neq \{e\}$, then by Hartley-Shute Lemma 1.5.5 we obtain that $C_U(H_\zeta) = \{e\}$ that contradicts the condition $K \cap U \neq \{e\}$. If $H_\zeta = \{e\}$, then either G is twisted and $|\zeta| = t$, that implies statement (e) of the lemma; or G is twisted, $|\zeta| = 2t$, in particular, t is odd, that implies point (f) of the lemma.

Assume that $O^{2'}(G) \simeq A_2(2^t)$, ζ is a graph-field automorphism and t is odd. If $|\zeta| \neq 2t$, then arguments, using Hartley-Shute Lemma 1.5.5, similar to the proof of Lemma 2.4.2 show that $C_U(H_\zeta) = \{e\}$, that contradicts to the condition $K \cap U \neq \{e\}$. If $|\zeta| = 2t$, then we obtain point (f) of the lemma.

Assume now that $O^{2'}(G)\simeq A_2(2^{2t})$ and ζ is a graph-field automorphism. Again for $|\zeta|\neq 2t$ from Hartley-Shute Lemma 1.5.5 it follows that $C_U(H_\zeta)=\{e\}$, that contradicts to the condition $K\cap U\neq\{e\}$. If $|\zeta|=2t$, then either $G_\zeta\simeq {}^2A_2(2^2)$, or $G_\zeta\simeq \widehat{{}^2A_2(2^2)}$. If $G_\zeta\simeq {}^2A_2(2^2)$, then $H_\zeta=\{e\}$ and we obtain the statement (d) of the lemma. If $G_\zeta\simeq \widehat{{}^2A_2(2^2)}$, then $|H_\zeta|=3$, and so $KU/U\cap HU/U$ is a cyclic group $\langle y\rangle$ of order $(2^{t_3}+1)_3=3^k$, where $3^{k-1}=t_3$. If $k>1$, then Hartley-Shute Lemma 1.5.5 implies that $C_U(y)=\{e\}$, that is impossible. Thus t is nor divisible by 3 and $K\cap U$ is contained in the centralizer of an element x, generating H_ζ. Consider the homomorphism $GL_3(2^{2t})\rightarrow \mathbf{P}GL_3(2^{2t})$. Then some preimage of x is similar to the matrix

$$\begin{pmatrix} \lambda & 0 & 0\\ 0 & \lambda^2 & 0\\ 0 & 0 & \lambda \end{pmatrix},$$

where λ is the generating element of the multiplicative group of $GF(2^2)$. The preimage of U is similar with the set of upper triangular matrices with the same elements on the diagonal. Direct calculations show that $C_U(x)$ is isomorphic to the additive group of $GF(2^{2t})$. The nilpotency of K implies that $K\cap U=(C_U(x))_{\zeta_{2'}}$, and point (c) of the lemma follows.

Assume now that $O^{2'}(G)\simeq {}^2A_2(2^{2t})$. By Lemma 2.4.1 KU/U is a Carter subgroup of $\langle B,\zeta g\rangle/U\simeq\langle H,\zeta g\rangle$ and, as above, we may assume that $H_\zeta\leq K$. If $|\zeta|=2t$, then $G_\zeta\simeq SL_2(2)$ and $H_\zeta=\{e\}$, and point (e) of the lemma follows. Assume that t is even and $|\zeta|\leqslant t$. Then either $O^{2'}(G_\zeta)\simeq SL_2(2^{2t/|\zeta|})$ (if the order $|\zeta|$ is even), or $O^{2'}(G_\zeta)\simeq {}^2A_2(2^{2t/|\zeta|})$ (if the order $|\zeta|$ is odd, hence $|\zeta|<t$). Clearly H_ζ contains an element x such that $K\cap U\leq C_U(H_\zeta)=\{e\}$, and this gives a contradiction with the condition $K\cap U\neq\{e\}$. If t is odd and $t\neq|\zeta|$, then $O^{2'}(G_\zeta)\simeq {}^2A_2(2^{2t/|\zeta|})$, and it follows that H_ζ contains an element x such that $C_U(x)=\{e\}$. If $|\zeta|=t$ and t is odd, then the order $|KU/U\cap B/U|$ can be divisible only by 3 (otherwise by Hartley-Shute Lemma 1.5.5 it again follows that $C_U(H_\zeta)=\{e\}$). If $G_\zeta\simeq {}^2A_2(2^{2t/|\zeta|})$, then $H_\zeta=\{e\}$ and we obtain point (b) of the lemma. If $G_\zeta\simeq \widehat{{}^2A_2(2^{2t/|\zeta|})}$, then $KU/U\cap HU/U$ is a cyclic group $\langle y\rangle$ of order $(2^{t_3}+1)_3=3^k$, where $3^{k-1}=t_3$. If $k>1$, then Hartley-Shute Lemma 1.5.5 implies, that $C_U(y)=\{e\}$, that is impossible. Thus t is not divisible by 3 and $K\cap U$ is contained in the centralizer of an element x, generating H_ζ. As in the non-twisted case above, we obtain that $C_U(x)$ is isomorphic to the additive group of $GF(2^t)$. The nilpotency of K implies that $K\cap U=(C_U(x))_{\zeta_{2'}}$, and point (a) of the lemma follows. □

5 Carter subgroups of semilinear groups

5.1 Brief review of results of this section

In this section, by using notations and results obtained in section 4, we classify Carter subgroups in groups of automorphisms of finite groups of Lie type. First we give such a classification in the case, when a group of Lie type has type C_n or when a group of its automorphisms contains a triality automorphism, since the arguments in these two cases differ from the remainings. The we formulate the final theorem and we prove this theorem in two subsections. In the last subsection we prove that in every finite group with known composition factors Carter subgroups are conjugate.

5.2 Carter subgroups of symplectic groups

Consider a set $\mathcal{A}$ of almost simple groups A such that a unique non-Abelian composition factor $S = F^*(A)$ is a canonical simple group of Lie type and A contains nonconjugate Carter subgroups. If the set $\mathcal{A}$ is not empty, denote by **Cmin** the minimal possible order of $F^*(A)$, with $A \in \mathcal{A}$. If the set $\mathcal{A}$ is empty, then let $\mathbf{Cmin} = \infty$. We shall prove that $\mathbf{Cmin} = \infty$, i. e. that $\mathcal{A} = \varnothing$. Note that if $A \in \mathcal{A}$ and $G = F^*(A)$, then there exists a subgroup A_1 of A such that $A_1 \in \mathcal{A}$ and $A_1 = KG$ for a Carter subgroup K of A. Indeed, if for every nilpotent subgroup N of A Carter subgroups of NG are conjugate, then A satisfies **(C)**, hence Carter subgroups of A are conjugate, that contradicts to the choice of A. So there exists a nilpotent subgroup N of A such that Carter subgroups of NG are not conjugate. Let K be a Carter subgroup of NG. Then clearly KG/G is a Carter subgroup of NG/G, i. e., coincides with NG/G. Therefore Carter subgroups of KG are not conjugate and $KG = A_1 \in \mathcal{A}$. So the condition $A = KG$ in Theorems 5.2.3, 5.3.1, and 5.4.1 is not a restriction and is used only to simplify arguments.

In this section we consider Carter subgroups in an almost simple group A with simple socle $G = F^*(A) \simeq \mathbf{P}\mathrm{Sp}_{2n}(q)$. We consider such groups in the separate section, since for groups of type $\mathbf{P}\mathrm{Sp}_{2n}(q)$ Lemma 3.2.9 is not true and we use arguments slightly different from those that we use in the proof of Theorem 5.4.1.

We shall prove first two technical lemmas.

Lemma 5.2.1. *Let $O^{p'}(\overline{G}_\sigma) = G$ be a canonical adjoint finite group of Lie type over a field of odd characteristic p and -1 is not a square in the base field of G. Assume that the root system Φ of $\overline{G}$ equals C_n. Let U be a maximal unipotent subgroup of G, H be a Cartan subgroup of G, normalizing U, and Q is a Sylow 2-subgroup of H.*
Then $C_U(Q) = \langle X_r \mid r \text{ is a long root}\rangle$.

Proof. If r is a short root, then there exists a root s with $< s, r >= 1$. Thus

$$x_r(t)^{h_s(-1)} = x_r((-1)^{<s,r>}t) = x_r(-t)$$

(see [15, Proposition 6.4.1]). Therefore, if $x \in C_U(Q)$ and $x_r(t)$ is a nontrivial multiplier in decomposition (1) of x, then r is a long root. Now if r is a long root, then for every root s either $| < s, r > | = 2$, or $< s, r >= 0$, i. e., $x_r(t)^{h_s(-1)} = x_r(t)$. Under the condition that -1 is not a square in the base field of G (i. e., in the field $\mathbb{F}_q$) we obtain that $q \equiv -1 \pmod 4$, so $\langle h_s(-1) \mid s \in \Phi\rangle = Q$, and the lemma follows. □

Lemma 5.2.2. *Let $G = \mathbf{P}\mathrm{Sp}_{2n}(q)$ be a simple canonical group of Lie type, J be a subset of the set of fundamental roots, containing the long fundamental root r_n, P_J be a parabolic subgroup, generated by a Borel subgroup B and by groups X_r with $-r \in J$, L be a Levi factor of P_J. Denote by S a quasisimple normal subgroup of L, isomorphic to $\mathrm{Sp}_{2k}(q)$ (it always exists, since $r_n \in J$).*
Then $Aut_L(S/Z(S)) = S/Z(S)$.

Proof. This statement is known, it is proven in an unpublished paper by N.A.Vavilov. We give a proof here for the completeness. As we noted above, L is a reductive subgroup of maximal rank of G, ans so the following inclusions hold $S/Z(S) \leq \mathrm{Aut}_L(S/Z(S)) \leq \widehat{S/Z(S)}$. Since $|\widehat{C_n(q)} : C_n(q)| = (2, q-1)$, then for q even the statement is evident. If q is odd, then for $\mathrm{Aut}_L(S/Z(S))$ there can be only two possibilities: either $\mathrm{Aut}_L(S/Z(S)) = S/Z(S)$, or $\mathrm{Aut}_L(S/Z(S)) = \widehat{S/Z(S)}$. We shall show that the second equality is impossible.

In our notations fundamental roots of the root system of S are $r_{n-k+1},\ldots,r_n$. If the equality $\mathrm{Aut}_L(S/Z(S))=\widehat{S/Z(S)}$ holds, then there exist elements $s_1,\ldots,s_k$ of $\mathbb{Z}\Phi=\mathbb{Z}C_n$ such that

$$< s_i, r_{n-k+j} >= \frac{(s_i, r_{n-k+j})}{(s_i,s_i)} = \begin{cases} 1 & \text{if } i=j,\\ 0 & \text{if } i\neq j.\end{cases}$$

(They generate the lattice of fundamental weights, thus allow to obtain all diagonal automorphisms of S). But for each root s of C_n we have that either $< s, r_n >= 0$, or $< s, r_n >= \pm 2$, i. e., for each element $s\in\mathbb{Z}\Phi$ the number $< s, r_n >$ is even, in particular is distinct from 1. Therefore such a set of elements $s_1,\ldots,s_k$ does not exists. $\square$

Theorem 5.2.3. *Let G be a finite adjoint group of Lie type (not necessary simple) over a field of characteristic p, and $\overline{G}$, σ are chosen so that $\mathbf{P}\mathrm{Sp}_{2n}(p^t)\simeq O^{p'}(\overline{G}_\sigma)\leq G\leq \overline{G}_\sigma$. Choose a subgroup A of $A\cap\overline{G}_\sigma=G$. Let K be a Carter subgroup of A. Assume also that $|\mathbf{P}\mathrm{Sp}_{2n}(p^t)|\leqslant$ **Cmin** and $A=KG$.*
Then exactly one of the following statements holds:

1. *G is defined over $GF(2^t)$, a field automorphism ζ is in A, $|\zeta|=t$, and, up to conjugation in G, the equality $K=Q\leftthreetimes\langle\zeta\rangle$ holds, where Q is a Sylow 2-subgroup of $G_{\zeta_{2'}}$.*
2. *$G\simeq\mathbf{P}\mathrm{SL}_2(3^t)\simeq\mathbf{P}\mathrm{Sp}_2(3^t)$, a field automorphism ζ is in A, $|\zeta|=t$ is odd, and, up to conjugation in G, the equality $K=Q\leftthreetimes\langle\zeta\rangle$ holds, where Q is a Sylow 3-subgroup of $G_{\zeta_{3'}}$.*
3. *p does not divide $|K\cap G|$ and K is contained in the normalizer of a Sylow 2-subgroup of A.*

*In particular, Carter subgroups of A are conjugate, i. e., if $A_1\in\mathcal{A}$ and $F^*(A_1)=$ **Cmin**, then $F^*(A_1)\not\simeq\mathbf{P}\mathrm{Sp}_{2n}(p^t)$.*

Proof. Assume that the theorem is not true and A is a counter example such that $|F^*(A)|$ is minimal. Note that no more than one statement of the theorem can be fulfill, since if statement (2) holds, then, by Lemmas 4.3.2 and 4.3.3, for a Sylow 2-subgroup Q of A the condition $N_G(S)=SC_G(S)$ is not true, i. e., statement (3) of the theorem does not hold. Thus if A_1 is an almost simple group with $F^*(A_1)$ being a simple group of Lie type of order less, than $|F^*(A)|$, then Carter subgroups of A_1 are conjugate. In view of Theorem 1.5.6 we may assume that $A\neq G$. Moreover, by Theorem 3.3.5, we may assume that q is odd, i. e., that $\mathrm{Aut}(\mathbf{P}\mathrm{Sp}_{2n}(q))$ does not contain a graph automorphism. Thus we may assume that $A=\langle G,\zeta g\rangle$.

Assume that K is a Carter subgroup of $\langle G,\zeta g\rangle$ and K does not satisfy to the statement of the theorem. Write $K=\langle x,K\cap G\rangle$. If either $p\neq 3$ or t is even, then the theorem follows from Theorem 3.3.5. Thus we may assume that $q=3^t$ and t is odd. Since $|\overline{G}_\sigma:O^{p'}(\overline{G_\sigma})|=2$ and the order $|\zeta|$ is odd, we may assume that the order $|\zeta g|$ is also odd and so $\zeta\in\langle G,\zeta g\rangle$, i. e., $A=G\leftthreetimes\langle\zeta\rangle$. By Lemma 3.2.3 every semisimple element of odd order is conjugate to its inverse in G. Now, for every semisimple element $t\in G$, each non-Abelian composition factor of $C_G(t)$ is a simple group of Lie type (see [17]) of order less, than **Cmin**. Therefore, for every non-Abelian composition factor S of $C_A(t)$ and every nilpotent subgroup $N\leq C_A(t)$, Carter subgroups of $\langle\mathrm{Aut}_N(S),S\rangle$ are conjugate. It follows that $C_A(t)$ satisfies **(C)**. Hence, by Lemma 2.4.2, $|K\cap G|=2^\alpha\cdot 3^\beta$ for some $\alpha,\beta\geqslant 0$.

If $G=\widehat{\mathbf{P}\mathrm{Sp}_{2n}(q)}$ then by [46, Theorem 2] every unipotent element is conjugate to its inverse. Since 3 is a good prime for G, then [40, Theorems 1.2 and 1.4] imply that, for any element

$u\in G$ of order 3, all composition factors of $C_G(u)$ are simple groups of Lie type of order less, than **Cmin**. Thus $C_A(u)$ satisfies **(C)**, hence, by Lemma 2.4.2, we obtain that $K\cap G$ is a 2-group. By Lemmas 4.2.5 and 4.2.6 every element $x\in A\setminus G$ of odd order with $\langle x\rangle\cap G=\{e\}$ centralizes some Sylow 2-subgroup of G. Hence K contains a Sylow 2-subgroup of G, and hence of A, i. e., K satisfies statement (3) of the theorem.

Thus we may assume that $G=\mathbf{P}Sp_{2n}(q)$ and $\beta\geqslant 1$, i. e., a Sylow 3-subgroup $O_3(K\cap G)$ of $K\cap G$ is nontrivial. By Lemma 4.2.1 we obtain that $K\cap G$ is contained in some K-invariant parabolic subgroup P of G with a Levi factor L and, up to conjugation in P, a Sylow 2-subgroup $O_2(K\cap G)$ of $K\cap G$ is contained in L. Note that all non-Abelian composition factors of P are simple groups of Lie type of order less, than **Cmin**, so P and each its homomorphic image satisfy **(C)**. The group $\widetilde{K}=KO_3(P)/O_3(P)$ is isomorphic to $K/O_3(K\cap G)$ and, by Lemma 2.4.1, $\widetilde{K}$ is a Carter subgroup of $\langle\widetilde{K},P/O_3(P)\rangle$. Now $\widetilde{K}\cap P/O_3(P)\simeq O_2(K\cap G)$ is a 2-group and every element $x\in\langle\widetilde{K},P/O_3(P)\rangle\setminus P/O_3(P)$ of odd order with $\langle x\rangle\cap P/O_3(P)=\{e\}$ centralizes a Sylow 2-subgroup of $P/O_3(P)\simeq L$ (see Lemmas 4.2.5 and 4.2.6). Therefore $O_2(K\cap G)$ contains a Sylow 2-subgroup of L, in particular, contains a Sylow 2-subgroup H_2 of H. Since K is nilpotent, Lemma 5.2.1 implies that $O_3(K\cap G)\leq C_U(H_2)=\langle X_r\mid r\text{ is a long root of }\Phi(G)^+\rangle$. Since for every two long positive roots r,s in $\Phi(G)^+$ we have that $r+s\notin\Phi(G)$, Chevalley commutator formula [15, Theorem 5.2.2] (Lemma 1.3.1) implies that $\langle X_r\mid r\text{ is a long root of }\Phi(G)^+\rangle$ is Abelian.

Since ζ is a field automorphism, it normalizes each parabolic subgroup of G containing a ζ-stable Borel subgroup. Thus for every subset J of the set of fundamental roots $\Pi=\{r_1,\ldots,r_n\}$ of $\Phi=\Phi(G)$ the parabolic subgroup P_J is ζ-stable. Therefore we may suppose that $P=P_J$, where J is a proper subset of the set of fundamental roots Π of Φ. Choose the numbering of fundamental roots so that r_n is a long fundamental root, while the remaining fundamental roots r_i are short roots. If $r_n\in J$, then one of the components of the Levi factor, G_1 for example, is isomorphic to $Sp_{2k}(q)$ for some $k<n$ (note that since $A\neq G$ then $q\neq 3$). By Lemma 5.2.2 we obtain that $L/C_L(G_1)=\mathrm{Aut}_L(G_1/Z(G_1))=G_1/Z(G_1)$. By Lemma 2.4.1 $K_1=KC_L(G_1)O_3(P)/C_L(G_1)O_3(P)$ is a Carter subgroup of $(P\leftthreetimes\langle\zeta\rangle)/C_L(G_1)O_3(P)$. Since $|K_1\cap P/C_L(G_1)O_3(P)|$ is not divisible by 3, and ζ centralizes a Sylow 2-subgroup of $G_1/Z(G_1)$ (see Lemma 4.2.5), then K_1 contains a Sylow 2-subgroup of $P/C_L(G_1)O_3(P)\simeq G_1/Z(G_1)\simeq\mathbf{P}Sp_{2k}(q)$. Moreover by Lemma 4.2.5 a Sylow 2-subgroup of $(P/C_L(G_1)O_3(P))_\zeta$ is a Sylow 2-subgroup of $P/C_L(G_1)O_3(P)$. Thus $K_1\cap P/C_L(G_1)O_3(P)$ is a Sylow 2-subgroup of $(P/C_L(G_1)O_3(P))_\zeta\simeq\mathbf{P}Sp_{2k}(3)$. By Lemma 4.3.2 there exists an element x of odd order of $\mathbf{P}Sp_{2k}(3)$ that normalizes but not centralizes a Sylow 2-subgroup; a contradiction with the fact that K_1 is a Carter subgroup of $(P\leftthreetimes\langle\zeta\rangle)/C_L(G_1)O_3(P)$. Thus we may assume that $r_n\notin J$.

Consider the set $J_n=\Pi\setminus\{r_n\}$ and the parabolic subgroup P_{J_n}. From the above arguments it follows that $K\leq P_J\leftthreetimes\langle\zeta\rangle\leq P_{J_n}\leftthreetimes\langle\zeta\rangle$. Now the subgroup $\langle X_r\mid r\text{ is a long root of }\Phi(G)^+\rangle$ is contained in $O_3(P_{J_n})$ and $O_3(K\cap G)$ is contained in $\langle X_r\mid r\text{ is a long root of }\Phi(G)^+\rangle$, so $N_G(O_3(K\cap G))\leq O_3(P_{J_n})$ and we may assume that $P=P_{J_n}$. By Lemma 2.4.1, $\widetilde{K}=KO_3(P)/O_3(P)$ is a Carter subgroup of $(P\leftthreetimes\langle\zeta\rangle)/O_3(P)$. Note that a unique non-Abelian composition factor of $P\leftthreetimes\langle\zeta\rangle$ is isomorphic to $A_{n-1}(q)\simeq\mathbf{P}SL_n(q)$. By [9, Theorem 1] and [10, Theorem 4] we obtain that $\widetilde{K}=R\times\langle\zeta\rangle$, where R is a Sylow 2-subgroup of P centralized by ζ. Thus $O_3(K\cap G)\leq C_P(R)$. Consider $Q=O_3(K\cap G)\cap P_\zeta$. Since $O_3(K\cap G)$ is nontrivial and K is nilpotent, then $Q=O_3(K\cap G)\cap P_\zeta=Z(K)\cap O_3(K\cap G)$ is nontrivial. Therefore $N_G(Q)$ is a proper subgroup of G and by Lemma 4.2.1 $N_G(Q)$ is contained in a proper parabolic subgroup of G. On the other hand, $K\leq N_G(Q)$ and $P=P_{J_n}$ is a maximal proper parabolic subgroup of

G. If $N_G(Q)$ is not contained in P, then $N_G(Q)$ and K are contained in a parabolic subgroup P_J with $r_n \in J$. We have proved above that $r_n \notin J$, so $N_G(Q)$ is contained in P.

We shall show that $R \times Q$ is a Carter subgroup of G_ζ. Indeed, assume that an element $x \in G_\zeta$ normalizes $R \times Q$. Then x normalizes Q, so x is in P and normalizes $O_3(P)$. On the other hand x normalizes R, therefore normalizes $C_P(R)$, so x normalizes $C_{O_3(P)}(R)$. Moreover it is evident that x and ζ commute. Thus x normalizes $(R \times C_{O_3(P)}(P)) \leftthreetimes \langle\zeta\rangle$. As we noted above, $K \leq (R \times C_{O_3(P)}(P)) \leftthreetimes \langle\zeta\rangle$ and $(R \times C_{O_3(P)}(P)) \leftthreetimes \langle\zeta\rangle$ is solvable. Lemma 2.4.2(a) implies that $(R \times C_{O_3(P)}(P)) \leftthreetimes \langle\zeta\rangle$ coincides with its normalizer in $G \leftthreetimes \langle\zeta\rangle$, so $x \in R \times C_{O_3(P)}(R)$. The group $C_{O_3(P)}(R) \leq \langle X_r \mid r \text{ is a long root of } \Phi(G)^+\rangle$ is Abelian, so every element of $R \times C_{O_3(P)}(R)$ centralizes $C_{O_3(P)}(R) \geq O_3(K \cap G)$. Therefore x normalizes $(R \times C_{O_3(P)}(P)) \leftthreetimes \langle\zeta\rangle = K$, i. e., $x \in K$. By construction $R \times Q = K \cap G_\zeta$, so $x \in R \times Q$ and $R \times Q$ is a Carter subgroup of G_ζ. On the other hand $O^{3'}(G_\zeta) \simeq \mathbf{P}Sp_{2n}(3^{t/|\zeta|})$ and by induction groups $\mathbf{P}Sp_{2n}(3^{t/|\zeta|})$ and $\mathbf{P}\widehat{Sp_{2n}}(3^{t/|\zeta|})$ does not contain Carter subgroups of order divisible by 3. This final contradiction completes the proof of the theorem. ☐

5.3 Группы с автоморфизмом тройственности

Theorem 5.3.1. *Let G be a finite adjoint group of Lie type over a field of characteristic p, $\overline{G}$, σ are chosen so that $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$, and $O^{p'}(\overline{G}_\sigma)$ is isomorphic to either $D_4(q)$, or ${}^3D_4(q^3)$. Assume that τ is a graph automorphism of order 3 of $O^{p'}(G)$ (recall that for $G \simeq {}^3D_4(q^3)$ τ is an automorphism such that the set of its stable points is isomorphic to $G_2(q)$). Denote by A_1 the subgroup of $\mathrm{Aut}(D_4(q))$ generated by inner-diagonal and field automorphisms, and also by a graph automorphism of order 2. Let $A \leq \mathrm{Aut}(G)$ be such that $A \not\leq A_1$ (if $O^{p'}(G) \simeq D_4(q)$), and K be a Carter subgroup of A. Assume also that $|O^{p'}(G)| \leqslant$ **Cmin**, $G = A \cap \overline{G}_\sigma$ and $A = KG$. Then one of the following statements holds:*

(a) *$G \simeq {}^3D_4(q^3)$, $(|A : G|, 3) = 1$, q is odd and K contains a Sylow 2-subgroup of A;*

(b) *$(|A : G|, 3) = 3$, q is odd, $\tau \in A$ and, up to conjugation by an element of G, the subgroup K contains a Sylow 2-subgroup of $C_A(\tau) \in \Gamma G_2(q)$, and $\tau \in K$;*

(c) *$(|A : G|, 3) = 3$, $q = 2^t$, $|A : G| = 3t$, $A = G \leftthreetimes \langle\tau, \varphi\rangle$, where φ is a field automorphism of order t commuting with τ and, up to conjugation by an element of G, the subgroup K contains a Sylow 2-subgroup of $C_G(\langle\tau, \varphi\rangle_{2'}) \simeq G_2(2^{t_{2'}})$ and $\tau \in K$;*

(d) *$O^{p'}(G) \simeq D_4(p^{3t})$, p is odd, the factor group A/G is cyclic, $\tau \notin A$, $A = G \leftthreetimes \langle\zeta\rangle$, where for some natural m, $\zeta = \tau\varphi^m$ is a graph-field automorphism, and , up to conjugation by an element of G, $K = Q \leftthreetimes \langle\zeta\rangle$, where Q is a Sylow 2-subgroup of $C_G(\zeta_{2'}) \simeq {}^3D_4(p^{3t/|\zeta_{2'}|})$.*

*In particular, Carter subgroups of A are conjugate, i. e., if $A_2 \in \mathcal{A}$ and $|F^*(A_2)| =$ **Cmin**, then A_2 does not satisfy to the conditions of the theorem, so $F^*(A_2) \not\simeq {}^3D_4(q^3)$.*

Proof. Assume that the theorem is not true and A is a counter example such that $|O^{p'}(G)|$ is minimal. In view of [44, Theorem 1.2(vi)] we have that every element of G is conjugate to its inverse. By [17] and [40, Theorems 1.2 and 1.4] we obtain that for every element $t \in G$ of odd prime order, all non-Abelian composition factors of $C_G(t)$ are simple groups of Lie type of order less, than **Cmin**. Thus, $C_A(t)$ satisfies **(C)** and Lemma 2.4.2 implies that $K_G = K \cap G$ is a 2-group. Now Lemma 4.2.6 implies that all cyclic groups, generated by field automorphisms of the same odd order of G, are conjugate under G. Since the centralizer of every field automorphism in G is a group of Lie type of order less, than **Cmin**, we again use Lemma 2.4.2 and obtain the statement of the theorem by induction. Lemma 4.2.6 implies also that if $O^{p'}(G) \simeq D_4(q)$, then all cyclic groups generated by graph-field automorphisms are conjugate. Since the centralizes of each graph-field automorphism in G is a group of Lie type of order less than **Cmin**, we again use Lemma 2.4.2 and obtain statement (d) of the theorem by induction. Thus we may assume that A does not contain a field automorphism or a graph-field automorphism of odd order. Therefore either $G \simeq {}^3D_4(q^3)$ and A/G is a 2-group, or K contains an element s of order 3 such that $\langle s\rangle \cap A_1 = \{e\}$ (for groups ${}^3D_4(q^3)$ the equality $\langle s\rangle \cap G = \{e\}$ holds), $G \leftthreetimes \langle s\rangle = G \leftthreetimes \langle \tau\rangle$ and $K \cap G$ is a 2-group.

In the first case we obtain the statement (a) of the theorem with condition $(|A : G|, 3) = 1$. In the second case there exists two non-conjugate cyclic subgroups $\langle\tau\rangle$ and $\langle x\rangle$ of order 3 of A such that $\langle\tau\rangle \cap A_1 = \langle x\rangle \cap A_1 = \{e\}$ and $G \leftthreetimes \langle x\rangle = G \leftthreetimes \langle\tau\rangle$ (see [32, (9-1)]). Hence, either $s = \tau \in K$, or $s = x \in K$. Assume that $q \neq 3^t$. In the first case from the known structure of Carter subgroups in a group from the set $\Gamma G_2(q)$, obtained in Theorem 3.3.5, the statement (b) or (c) of the theorem follows, in the second case we have that $K \leq C_A(x)$. By [32, (9-1)] $C_G(x) \simeq \mathbf{P}GL_3^{\varepsilon}(q)$, where $q \equiv \varepsilon 1 \pmod 3$, $\varepsilon = \pm$ and $\mathbf{P}GL_3^{+}(q) = \mathbf{P}GL_3(q)$, $\mathbf{P}GL_3^{-}(q) = \mathbf{P}GU_3(q)$. Then $K = (K \cap G) \times \langle y, \varphi\rangle$, where φ is a field automorphism of $O^{p'}(G)$ of order equal to a power of 2 and y is a graph automorphism such that its order is a power of 3 and $x \in \langle y\rangle$. By nilpotency of K we obtain that $y\varphi = \varphi y$, it follows that $C_{C_G(\varphi)}(x) = C_{C_G(x)}(\varphi)$. Now we have that

$$C_G(\varphi) = \begin{cases} D_4(q^{1/|\varphi|}), & \text{if } O^{p'}(G) \simeq D_4(q), \\ {}^3D_4(q^{3/|\varphi|}), & \text{if } G \simeq {}^3D_4(q^3). \end{cases}$$

Hence $C_{C_G(x)}(\varphi) = C_{C_G(\varphi)}(x) \simeq \mathbf{P}GL_3^{\mu}(q^{1/|\varphi|})$, with $q^{1/|\varphi|} \equiv \mu 1 \pmod 3$, where $\mu = \pm$ (note that ε and μ can be different). As we noted above, $K \cap G$ is a 2-group. On the other hand, by [10, Theorem 4] there exists an element z of order 3 centralizing a Sylow 2-subgroup of $C_G(x) = \mathbf{P}GL_3^{\varepsilon}(q)$ and belonging to $C_{C_G(x)}(\varphi) \simeq \mathbf{P}GL_3^{\mu}(q^{1/|\varphi|})$. Thus z centralizes K, hence is in K. But $K \cap G$ does not contain elements of odd order, therefore this second case is impossible.

Assume now that $q = 3^t$. Then $C_G(\tau) \simeq G_2(q)$ and we obtain the theorem. In the second case $C_G(x) \simeq SL_2(q) \rightthreetimes U$, where U is a 3-group and $Z(C_G(x)) \cap U \neq \{e\}$, a contradiction with Lemma 2.4.2. □

5.4 Classification theorem

Theorem 5.4.1. *Let G be a finite adjoint group of Lie type (G is not necessary simple) over a field of characteristic p and $\overline{G}$, σ are chosen so that $O^{p'}(\overline{G}_\sigma) \leq G \leq \overline{G}_\sigma$. Assume also that $G \not\simeq {}^3D_4(q^3)$. Choose a subgroup A of $\mathrm{Aut}(O^{p'}(\overline{G}_\sigma))$ with $A \cap \overline{G}_\sigma = G$ and, if $O^{p'}(G) = D_4(q)$, assume that A is contained in the subgroup A_1 defined in Theorem* 5.3.1. *Let K be a Carter subgroup of A and assume that $A = KG$.*
Then exactly one of the following statements holds:

(a) *G is defined over a field of characteristic* 2, *$A = \langle\langle G, \zeta g, t\rangle$, where t is a 2-element, K is contained in the normalizer of a t-stable Borel subgroup of G $K \cap \langle G, \zeta g\rangle$ satisfies to one of the statements* (a)–(f) *of Lemma* 4.3.4;

(b) *$G \simeq \mathbf{P}SL_2(3^t)$, a field automorphism ζ is in A, $|\zeta| = t$ is odd, and, up to conjugation in G, the equality $K = Q \leftthreetimes \langle\zeta\rangle$ holds, where Q is a Sylow* 3*-subgroup of $G_{\zeta_{3'}}$;*

(c) $A = {}^2G_2(3^{2n+1}) \leftthreetimes \langle\zeta\rangle$, $|\zeta| = 2n+1$, *and, up to conjugation in* G *the equality* $K = (K\cap G)\leftthreetimes\langle\zeta\rangle$ *holds, and* $K\cap {}^2G_2(3^{2n+1}) = Q\times P$, *where* Q *is of order* 2 *and* $|P| = 3^{|\zeta|_3}$.

(d) p *does not divide* $|K\cap G|$ *and* K *contains a Sylow* 2*-subgroup of* A, *in view of Lemma 4.3.3* A *satisfies* **(ESyl2)** *if and only if* G *satisfies* **(ESyl2)**.

In particular, Carter subgroups of A *are conjugate.*

Remark. There exists a dichotomy for Carter subgroups in groups of automorphisms of finite groups of Lie type, not containing a graph, or a graph-field automorphism of order 3. They either are contained in the normalizer of a Borel subgroup, or the characteristic is odd and a Carter subgroup contains a Sylow 2-subgroup of the hole group.

Assume that the theorem is not true and A is a counter example to the theorem with $|F^*(A)|$ minimal. Among counterexamples with $|F^*(A)|$ minimal take those, for which $|A|$ is minimal. In this case for every almost simple group A_1 such that $|F^*(A_1)| < |F^*(A)|$, $F^*(A_1)$ is a finite simple group of Lie type and A_1 satisfies the conditions of Theorem 5.4.1, Carter subgroups are conjugate. Indeed, note that no more, than one statement of the theorem can be fulfill, since if either statement (b), or statement (c) of the theorem holds, then the condition $N_A(Q) = QC_A(Q)$ for a Sylow 2-subgroup Q of A is not true, i. e., the statement (d) of the theorem does not hold (the fact that other statements can not hold simultaneously is evident). Thus, Carter subgroups of A_1 are conjugate. Note also that from this fact we immediately obtain the inequality $|F^*(A)| \leqslant \mathbf{Cmin}$. Indeed, if $A_2 \in \mathcal{A}$ and $F^*(A_2) = \mathbf{Cmin}$, then either A_2 satisfies to the condition of Theorem 5.3.1, or A_2 satisfies conditions of Theorem 5.4.1. As we noted in Theorem 5.3.1, the first case is impossible. The second case, as we just noted, is possible only if $|F^*(A)| \leqslant |F^*(A_2)| = \mathbf{Cmin}$ (since A is a counterexample to the statement of the theorem with $|F^*(A)|$ is minimal).

We shall prove the theorem in the following way. If $F^*(A) \simeq \mathbf{P}Sp_{2n}(q)$, then the theorem follows from Theorem 5.2.3. If $A = G$, then the theorem follows from [26], [27] and results from sections 3 and 4 of the present paper. Thus we may assume, that $A/(A\cap G)$ is nontrivial. Let K be a Carter subgroup of A. We shall prove first that if p divides $|K\cap G|$, then one of the statements (a)–(c) of the theorem holds. Then we shall prove that if p does not divide $|K\cap G|$, then K contains a Sylow 2-subgroup of A. Since both of these steps are quite complicated, we divide them onto two sections. Note that, in view of [17], for every semisimple element $t\in G$, all non-Abelian composition factors of $C_G(t)$, so of $C_A(t)$, are simple groups of Lie type of order less, than $|F^*(A)|$, and hence of order less, than **Cmin**. Therefore $C_A(t)$ satisfies **(C)**. In order to apply Lemmas 2.4.1 and 2.4.2 we shall use this fact without future references.

5.5 Carter subgroups of order divisible by characteristic

Denote $K\cap G$ by K_G. For every group A, satisfying conditions of Theorem 5.4.1, the factor group A/G is Abelian and, for some natural t is isomorphic to a subgroup of $\mathbb{Z}_2\times\mathbb{Z}_t$, where $\mathbb{Z}_t$ denotes a cyclic group of order t. If the factor group A/G is not cyclic, then the group $O^{p'}(G)$ is split and A contains an element τa, where τ is a graph automorphism of $O^{p'}(G)$ and $a\in\overline{G}_\sigma$. Then every semisimple element of odd order is conjugate to its inverse in A (cf. Lemma 4.2.7). By Lemma 2.4.2 we obtain that $|K_G|$ is divisible only by 2 and p. If $p = 2$, then we obtain that K_G is a 2-group, it is contained in a proper K-invariant parabolic subgroup P of G and by Lemma 2.4.1 $KO_2(P)/O_2(P)$ is a Carter subgroup of $KP/O_2(P)$. Since $K_G\leq O_2(P)$, then $(KO_2(P)/O_2(P))\bigcap(P/O_2(P)) = \{e\}$. Hence P is a Borel subgroup of G, otherwise we would have $C_{P/O_2(P)}(KO_2(P)/O_2(P)) \neq \{e\}$, a contradiction with the fact that $KO_2(P)/O_2(P)$ is a Carter subgroup of $KP/O_2(P)$. Thus P is a Borel subgroup and the theorem follows from Lemma 4.3.4. Now if $p \neq 2$, then again K_G is contained in a proper parabolic subgroup P of G such that $O_p(K_G) \leq O_p(P)$ and $O_2(K_G) \leq L$. Then Lemmas 4.2.5 and 4.2.6 implies that $H_2 \leq O_2(K \cap G) \leq K$. Now Lemma 3.2.9 implies that $O_p(K_G) \leq C_U(H_2) = \{e\}$. Therefore $K \cap G$ is a 2-group. By Lemmas 4.2.5 and 4.2.6 every element $x \in A \setminus G$ of odd order such that $\langle x \rangle \cap G = \{e\}$ centralizes some Sylow 2-subgroup of G. Hence K contains a Sylow 2-subgroup of A, i. e., K satisfies statement (d) of the theorem. Therefore A/G is cyclic and we may assume that $A = \langle G, \zeta g \rangle \in \Gamma G$.

Recall that we are in the conditions of Theorem 5.4.1, $A = \langle G, \zeta g \rangle$ is supposed to be a counter example to the theorem with $|O^{p'}(G)|$ and $|A|$ minimal, and K is a Carter subgroup of $\langle G, \zeta g \rangle$ such that p divides $|K_G|$. We have that $K = \langle \zeta^k g, K_G \rangle$. Since $|O^{p'}(G)| \leqslant \mathbf{Cmin}$, Lemma 2.4.1 implies that KG/G is a Carter subgroup of $\langle G, \zeta g \rangle /G$. Therefore $|\zeta^k| = |\zeta|$, and we may assume that $k = 1$ and $K = \langle K_G, \zeta g \rangle$.

In view of Lemma 4.2.1 there exists a proper σ- and $\bar{\zeta} g$- invariant parabolic subgroup $\overline{P}$ of $\overline{G}$ such that $O_p(K_G) \leq R_u(\overline{P})$ and $K_G \leq \overline{P}$. In particular, $\overline{P}$ and $\overline{P}^{\bar{\zeta}}$ are conjugate in $\overline{G}$. Let Φ be the root system of $\overline{G}$ and Π be a set of fundamental roots of Φ. In view of [15, Proposition 8.3.1], $\overline{P}$ is conjugate to some $\overline{P}_J = \overline{B} \cdot \overline{N}_J \cdot \overline{B}$, where J is a subset of Π and $\overline{N}_J$ is a complete preimage of W_J in $\overline{N}$ under the natural homomorphism $\overline{N}/\overline{T} \to W$. Now $\overline{P}_J$ is $\bar{\varphi}$-invariant, hence $\overline{P}_J^{\bar{\zeta}} = \overline{P}_J^{\bar{\gamma}^\varepsilon}$ (recall that $\bar{\zeta} = \bar{\gamma}^\varepsilon \varphi^k$ by definition). Consider the symmetry ρ of the Dynkin diagram of Φ corresponding to $\bar{\gamma}$. Let $\overline{J}$ be the image of J under ρ. Clearly $\overline{P}_J^{\bar{\gamma}} = \overline{P}_{\overline{J}}$. Since $\overline{P}$ and $\overline{P}^{\bar{\zeta}}$ are conjugate in $\overline{G}$ we obtain that $\overline{P}_J$ and $\overline{P}_J^{\bar{\zeta}}$ are conjugate in $\overline{G}$. By [15, Theorem 8.3.3] it follows that either $\varepsilon = 0$, or $J = \overline{J}$; i. e., $\overline{P}_J$ is $\bar{\zeta}$-invariant.

Now we have that $\overline{P}^{\bar{y}} = \overline{P}_J$ for some $\bar{y} \in \overline{G}$. So $\langle \bar{\zeta} g, \overline{P} \rangle^{\bar{y}} = \langle (\bar{\zeta} g)^{\bar{y}}, \overline{P}_J \rangle$ and $\overline{P}_J^{(\bar{\zeta} g)^{\bar{y}}} = \overline{P}_J$. It follows

$$(\bar{\zeta} g)^{\bar{y}} = \bar{y}^{-1} \bar{\zeta} g \bar{y} = \bar{\zeta} \left(\bar{\zeta}^{-1} \bar{y}^{-1} \bar{\zeta} g \bar{y} \right) = \bar{\zeta} \cdot h,$$

where $h = \left(\bar{\zeta}^{-1} \bar{y}^{-1} \bar{\zeta} g \bar{y} \right) \in \overline{G}$. Since $\overline{P}_J^{\bar{\zeta}} = \overline{P}_J = \overline{P}_J^{h^{-1}}$ we obtain that $h \in N_{\overline{G}}(\overline{P}_J)$. By [15, Theorem 8.3.3], $N_{\overline{G}}(\overline{P}_J) = \overline{P}_J$, thus $\langle \bar{\zeta} g, \overline{P} \rangle^{y} = \langle \bar{\zeta}, \overline{P}_J \rangle$. Now both $\overline{P}$ and $\overline{P}_J$ are σ-invariant. Hence $\bar{y}\sigma(\bar{y}^{-1}) \in N_{\overline{G}}(\overline{P}) = \overline{P}$. Therefore, by Lang-Steinberg Theorem (Lemma 1.5.3), we may assume that $\bar{y} = \sigma(\bar{y})$, i. e., $\bar{y} \in \overline{G}_\sigma$. Since $\overline{G}_\sigma = \overline{T}_\sigma \cdot O^{p'}(\overline{G}_\sigma)$ and $\overline{T} \leq \overline{P}_J$, then we may assume that $\bar{y} \in O^{p'}(\overline{G}_\sigma)$. Thus, up to conjugation in G, we may assume that $\overline{K} \leq \langle \bar{\zeta}, \overline{P}_J \rangle = \overline{P}_J \leftthreetimes \langle \bar{\zeta} \rangle$ and

$$K \leq \langle (\overline{P}_J \cap G), \zeta g \rangle = \langle P_J, \zeta g \rangle,$$

in particular, $g \in (\overline{P}_J)_\sigma$. Further if $\overline{L}_J = \langle \overline{T}, \overline{X}_r | r \in J \cup -J \rangle$, then $\overline{L}_J$ is a σ- and $\bar{\zeta}$- invariant Levi factor of $\overline{P}_J$ and $L_J = \overline{L}_J \cap G$ is a ζ-invariant Levi factor of P_J. Then L_J^g is a ζg-stable factor Levi of P_J. Since all Levi factors are conjugate under $O_p(P_J)$, we may assume that L_J is a ζg-stable Levi factor. Lemma 2.4.1 implies that

$$KO_p(P_J)/O_p(P_J) = X$$

is a Carter subgroup of $\langle P_J, \zeta g \rangle / O_p(P_J)$ and

$$KZ(L_J)O_p(P_J)/Z(L_J)O_p(P_J) = \widetilde{X}$$

is a Carter subgroup of $\langle P_J, \zeta g \rangle / Z(L_J)O_p(P_J)$. Recall that $K = \langle \zeta g, K_G \rangle$, hence, if v and $\tilde{v}$ are the images of g under the natural homomorphisms

$$\omega : \langle P_J, \zeta g \rangle \to \langle L_J, \zeta g \rangle \simeq \langle P_J, \zeta g \rangle / O_p(P_J),$$

$$\tilde{\omega} : \langle P_J, \zeta g\rangle \to \langle P_J, \zeta g\rangle / Z(L_J)O_p(P_J) \simeq \langle L_J, \zeta g\rangle / Z(L_J),$$

then $X = \langle \zeta v, K_G^{\omega}\rangle$ and $\widetilde{X} = \langle \zeta\tilde{v}, K_G^{\tilde{\omega}}\rangle$. Note that $O_p(P)$ and $Z(L_J)$ are characteristic subgroups of P and L_J respectively, hence we may consider ζ as an automorphism of $L_J \simeq P/O_p(P)$ and $\widetilde{L} = L_J/Z(L_J)$. Note also that all non-Abelian composition factors of P are simple groups of Lie type of order less than **Cmin**, hence $\langle P, \zeta g\rangle$ satisfies **(C)**. Thus we may apply Lemma 2.4.1 to $\langle \widetilde{L}, \zeta g\rangle$, $\langle L, \zeta g\rangle$, and $\langle P, \zeta g\rangle$.

If P_J is a Borel subgroup of G, then the statement of the theorem follows from Lemma 4.3.4. So we may assume that $L_J \neq Z(L_J)$, i. e., that P_J is not a Borel subgroup of G. Then $L_J = H(G_1 * \ldots * G_k)$, where G_i are subsystem subgroups of G, $k \geqslant 1$, and H is a Cartan subgroup of G. Let $\zeta g = (\zeta_2 g_2)\cdot(\zeta_{2'} g_{2'})$ be the product of 2- and 2′- parts of ζg (with $g_2, g_{2'} \in (\overline{P}_J)_\zeta$). Now $\zeta_{2'} = \varphi^k$, for some k, is a field automorphism (recall that we do not consider the triality automorphism) and it normalizes each G_i, since φ normalizes each G_i. Moreover, in view of Lemma 4.2.5, we have that $\zeta_{2'}$ centralizes a Sylow 2-subgroup of H. In particular, it centralizes a Sylow 2-subgroup of $Z(L_J) \leq H$. Therefore, every element of odd order of $\langle L_J, \zeta_{2'} v_{2'}\rangle$ centralizes a Sylow 2-subgroup of $Z(L_J)$ (here $v_{2'}$ is the image of $g_{2'}$ under ω).

Now $\widetilde{L} = (\mathbf{P}G_1 \times \ldots \times \mathbf{P}G_k)\widetilde{H}$, where $\widetilde{H} = H^{\omega}$ and $\mathbf{P}G_1, \ldots, \mathbf{P}G_k$ are canonical finite groups of Lie type with trivial center. Set $M_i = C_{\widetilde{L}}(\mathbf{P}G_i)$, clearly $M_i = (\mathbf{P}G_1 \times \ldots \times \mathbf{P}G_{i-1} \times \mathbf{P}G_{i+1} \times \ldots \times \mathbf{P}G_k)C_{\widetilde{H}}(\mathbf{P}G_i)$; denote by L_i the factor group $\widetilde{L}/M_i$ and by π_i corresponding natural homomorphism. Then L_i is a finite group of Lie type and $\mathbf{P}G_i \leq L_i \leq \widehat{\mathbf{P}G_i}$.

Set $M_{i,j} = C_{\widetilde{L}}(\mathbf{P}G_i \times \mathbf{P}G_j)$, then

$$M_{i,j} = (\mathbf{P}G_1 \times \ldots \times \mathbf{P}G_{i-1} \times \mathbf{P}G_{i+1} \times \ldots \times \mathbf{P}G_{j-1} \times \mathbf{P}G_{j+1} \times \ldots \times \mathbf{P}G_k)C_{\widetilde{H}}(\mathbf{P}G_i \times \mathbf{P}G_j);$$

denote by $\pi_{i,j}$ corresponding natural homomorphism $\widetilde{L} \to \widetilde{L}/M_{i,j}$. If M_i (respectively $M_{i,j}$) is ζ-invariant, then M_i (resp. $M_{i,j}$) is normal in $\langle \widetilde{L}, \zeta\tilde{v}\rangle$ and we denote by π_i (resp. $\pi_{i,j}$) the natural homomorphism $\pi_i : \langle \widetilde{L}, \zeta\tilde{v}\rangle \to \langle \widetilde{L}, \zeta\tilde{v}\rangle/M_i$ ($\pi_{i,j} : \langle \widetilde{L}, \zeta\tilde{v}\rangle \to \langle \widetilde{L}, \zeta\tilde{v}\rangle/M_{i,j}$).

Now consider ζ. Since ζ^2 is a field automorphism, there can be two cases: either ζ normalizes $\mathbf{P}G_i$, or ζ^2 normalizes $\mathbf{P}G_i$ and $\mathbf{P}G_i^{\zeta} = \mathbf{P}G_j$ for some $j \neq i$. Consider these two cases separately.

Let ζ normalizes $\mathbf{P}G_i$. Then ζ normalizes M_i, and Lemma 2.4.1 implies that $\widetilde{X}^{\pi_i} = K_i$ is a Carter subgroup of $\langle L_i, (\zeta\tilde{v})^{\pi_i}\rangle$. Since $\langle L_i, (\zeta\tilde{v})^{\pi_i}\rangle$ is a semilinear group of Lie type satisfying the conditions of Theorem 5.4.1 (by definition, ζ^2 is a field automorphism, so we are not in the conditions of Theorem 5.3.1), $|L_i| < |G|$, and p does not divide $|K_i|$, we have that K_i contains a Sylow 2-subgroup Q_i of $\langle L_i, (\zeta\tilde{v})^{\pi_i}\rangle$ (in particular, $p \neq 2$) and, by Lemma 2.4.3, the group $\langle L_i, (\zeta\tilde{v})^{\pi_i}\rangle$ satisfies **(ESyl2)**.

Let ζ^2 normalizes $\mathbf{P}G_i$ and $\mathbf{P}G_i^{\zeta} = \mathbf{P}G_j$. Then $M_{i,j}$ is normal in $\langle \widetilde{L}, \zeta\tilde{v}\rangle$. We want to show that $\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}$ satisfies **(ESyl2)**. Since $M_{i,j}$ is a normal subgroup of $\langle \widetilde{L}, \zeta\tilde{v}\rangle$, then, by Lemma 2.4.1, $(\widetilde{X})^{\pi_{i,j}}$ is a Carter subgroup of $\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}$. Consider the subgroup

$$\langle (\mathbf{P}G_i)^{\pi_{i,j}} \times (\mathbf{P}G_j)^{\pi_{i,j}}, \widetilde{X}^{\pi_{i,j}}\rangle$$

of $\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}$ (note that $(\mathbf{P}G_i)^{\pi_{i,j}} \simeq \mathbf{P}G_i$ and $(\mathbf{P}G_j)^{\pi_{i,j}} \simeq \mathbf{P}G_i$, and till the end of this paragraph for brevity we shall identify these groups). Now we are in the conditions of Lemma 2.2.3, namely we have a finite group $\widetilde{G} = (\widetilde{X})^{\pi_{i,j}}(\mathbf{P}G_i \times \mathbf{P}G_j)$, where $\mathbf{P}G_i \simeq \mathbf{P}G_j$ has trivial center. Then $\mathrm{Aut}_{(\widetilde{X})^{\pi_{i,j}}}(\mathbf{P}G_i) \simeq \mathrm{Aut}_{\widetilde{X}}(\mathbf{P}G_i)$ is a Carter subgroup of $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_i)$. Now $\mathbf{P}G_i$ is a canonical finite group of Lie type and

$$\mathbf{P}G_i \leq \mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_i) \leq \mathrm{Aut}(\mathbf{P}G_i),$$

i. e., $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_i)$ satisfies to the conditions of Theorem 5.4.1 (by construction ζ^2 is a field automorphism and so we are not in the conditions of Theorem 5.3.1) and $(\widetilde{X})^{\pi_{i,j}} \cap (\mathbf{P}G_i \times \mathbf{P}G_j)$ is not divisible by the characteristic. By induction, $\mathrm{Aut}_{(\widetilde{X})^{\pi_{i,j}}}(\mathbf{P}G_i)$ contains a Sylow 2-subgroup of $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_i)$ (in particular, $p \neq 2$). The same arguments show that $\mathrm{Aut}_{\widetilde{X}}(\mathbf{P}G_j)$ contains a Sylow 2-subgroup of $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_j)$. Therefore, $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_i)$ and $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_j)$ satisfy **(ESyl2)**. Since $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_i) \leq \mathrm{Aut}_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_i)$ and $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_j) \leq \mathrm{Aut}_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_j)$, Lemmas 4.3.1 and 4.3.3 imply that groups of induced automorphisms $\mathrm{Aut}_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_i)$ and $\mathrm{Aut}_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_j)$ satisfy **(ESyl2)**. Consider $N_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_i)$ and $N_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_j)$. Since

$$|\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}} : N_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_i)| = |\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}} : N_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_j)| = 2,$$

it is easy to see that for every element h of $\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}$ the equality of cosets $hN_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_i) = hN_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_j)$ holds, it follows that $N_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_i) = N_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_j)$. By construction $C_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_i) \cap C_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_j) = \{e\}$, so Lemma 2.4.5 (with $C_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_i)$ and $C_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_j)$ as normal subgroups) implies that the normalizer $N_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_i)$ satisfies **(ESyl2)**. Now $|\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}} : N_{\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}}(\mathbf{P}G_1)| = 2$, thus Lemma 2.4.6 implies that $\langle \widetilde{L}, \zeta\tilde{v}\rangle^{\pi_{i,j}}$ satisfies **(ESyl2)**.

Now we shall show that $\langle L_J, \zeta v\rangle$ satisfies **(ESyl2)**. Since $\widetilde{L} \neq \{e\}$, then, as we noted above, $p \neq 2$. Let Q be a Sylow 2-subgroup of $\langle L_J, \zeta v\rangle$. Consider an element $x \in N_{\langle L_J, \zeta v\rangle}(Q)$ of odd order. We need to prove that x centralizes Q. As we noted above, every element of odd order of $\langle L_J, \zeta v\rangle$ centralizes $Q \cap Z(L_J)$, hence, if $\tilde{x} = x^{\tilde{\omega}}$ centralizes $\widetilde{Q} = Q^{\tilde{\omega}} \simeq Q/(Q \cap Z(L_J))$, then x centralizes Q. Now either M_i is normal in $\langle \widetilde{L}, \zeta\tilde{v}\rangle$, or $M_{i,j}$ is normal in $\langle \widetilde{L}, \zeta\tilde{v}\rangle$ and $(\cap_i M_i) \bigcap (\cap_{i,j} M_{i,j}) = \{e\}$. Moreover, as we proved above, x^{π_i} centralizes $\widetilde{Q}M_i/M_i$, and $x^{\pi_{i,j}}$ centralizes $\widetilde{Q}M_{i,j}/M_{i,j}$. By Lemma 2.4.5 (with normal subgroup M_i and $M_{i,j}$) we obtain that $\tilde{x}$ centralizes $\widetilde{Q}$.

Thus $\langle L_J, \zeta v\rangle$ satisfies **(ESyl2)** and by Lemma 2.4.3 there exists a Carter subgroup F of $\langle L_J, \zeta v\rangle$ containing Q. Since $\langle L_J, \zeta v\rangle$ satisfies **(C)**, Theorem 2.1.4 implies that $X = K^{\omega}$ and F are conjugate, i. e., X contains a Sylow 2-subgroup of $\langle L_J, \zeta v\rangle$ and, up to conjugation in $\langle P_J, \zeta v\rangle$, K contains a Sylow 2-subgroup of $\langle P_J, \zeta v\rangle$. In particular, a Sylow 2-subgroup Q_1 of a Cartan subgroup H is in K and Q_1 centralizes $K \cap O_p(P_J) \neq \{e\}$; a contradiction with Lemma 3.2.9.

5.6 Carter subgroups of order not divisible by characteristic

Again we are in the conditions of Theorem 5.4.1. As we noted in the previous section, for every group A satisfying conditions of Theorem 5.4.1, the factor group A/G is Abelian and, for some natural t is isomorphic to a subgroup of $\mathbb{Z}_2 \times \mathbb{Z}_t$. If the factor group A/G is not cyclic, then $O^{p'}(G)$ is split and A contains an element τa, where τ is a graph automorphism of $O^{p'}(G)$ and $a \in \overline{G}_\sigma$. Thus, if A/G is not cyclic, or $\Phi(\overline{G}) \neq A_n, D_{2n+1}, E_6$, then by Lemmas 3.2.3 and 4.2.7 every semisimple element of G is conjugate to its inverse. By Lemma 2.4.2 we obtain that $K_G = K \cap G$ is a 2-group. In the conditions of Theorem 5.4.1 the group A/G is Abelian and, if $\overline{A}_1$ is a Hall $2'$-subgroup of A/G, then $\overline{A}_1$ is cyclic. Let x be the preimage of the generating element of $\overline{A}_1$ taken in K. Then $\langle x\rangle \cap G \leq \langle x\rangle \cap \overline{G}_\sigma \leq K \cap \overline{G}_\sigma = K \cap (A \cap \overline{G}_\sigma) = K \cap G$. As we noted above, $K \cap G$ is a 2-group, hence $\langle x\rangle \cap \overline{G}_\sigma = \{e\}$. By Lemma 4.2.6, the element x under $\overline{G}_\sigma$ is conjugate to a field automorphism of odd order and by Lemma 4.2.5, the element x centralizes a Sylow 2-subgroup of G (in particular, $p \neq 2$) and, since A/G is Abelian, Lemma

2.4.5 implies that K contains a Sylow 2-subgroup of A. Thus Theorem 5.4.1 is true in this case. So we may assume that $A = \langle G, \zeta g\rangle$ is a semilinear group of Lie type, $K = \langle \zeta^k g, K_G\rangle$ is a Carter subgroup of A, and $\Phi(\overline{G}) \in \{A_n, D_{2n+1}, E_6\}$. Like in the previous section we may assume that $k = 1$. Since G_ζ is nontrivial, then the centralizer $C_G(\zeta g)$ is also nontrivial, we have that K_G is also nontrivial. Since G_ζ is nontrivial, then the centralizer $C_G(\zeta g)$ is nontrivial, so K_G is also nontrivial. Therefore $Z(K) \cap K_G$ is nontrivial. Consider an element $x \in Z(K) \cap K_G$ of prime order. Then $K \in C_A(x) = \langle \zeta g, C_G(x)\rangle$. Now $C_{\overline{G}}(x)^0 = \overline{C}$ is a connected σ-stable reductive subgroup of maximal rank of $\overline{G}$. Moreover $\overline{C}$ is a characteristic subgroup of $C_{\overline{G}}(x)$ and $C_{\overline{G}}(x)/\overline{C}$ is isomorphic to a subgroup of Δ (see [34, Proposition 2.10]). Thus K is contained in $\langle K, C\rangle$, where $C = \overline{C} \cap G$. Moreover, by Lemma 4.1.1, the subgroup $C = \overline{C} \cap G = T(G_1 * \ldots * G_m)$ is normal in $C_A(x)$ and $K_G C/C$ is isomorphic to a subgroup of Δ. Assume that $|K_G|$ is not divisible by 2.

If $m = 0$, then $C = T = Z(C)$ is a maximal torus. Then $\overline{T}$ is $\bar{\zeta}g$-stable. In view of Lemma 4.2.4 we obtain that $N_A(C_A(x)) \neq C_A(x)$. Since $C_A(x)$ is solvable in this case this gives a contradiction with Lemma 2.4.2.

If $m \geqslant 1$, then $Z(C)$ and $G_1 * \ldots * G_m$ are normal subgroups of $\langle K, C\rangle$. Hence we may consider $\widetilde{G} = \langle K, G_1 * \ldots * G_m * Z(C)\rangle/Z(C) \leq \langle K, C\rangle/Z(C)$. Then $\widetilde{G} = \widetilde{K}(\mathbf{P}G_1 \times \ldots \times \mathbf{P}G_m)$, where $\widetilde{K} = KZ(C)/Z(C)$ is a Carter subgroup of $\widetilde{G}$ (see Lemma 2.4.1) and $Z(\mathbf{P}G_i)$ is trivial. Now $\widetilde{K}$ acts by conjugation on $\{\mathbf{P}G_1, \ldots, \mathbf{P}G_m\}$ and without lost of generality we may assume that $\{\mathbf{P}G_1, \ldots, \mathbf{P}G_m\}$ is a $\widetilde{K}$-orbit. Thus we are in the condition of Lemma 2.2.3 and $\mathrm{Aut}_{\widetilde{K}}(\mathbf{P}G_1)$ is a Carter subgroup of $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_1)$. Moreover $|\widetilde{K} \cap \mathbf{P}G_1 \times \ldots \times \mathbf{P}G_m|$ is not divisible by the characteristic. By induction we have that either $\mathrm{Aut}_{\widetilde{K}}(\mathbf{P}G_1)$ contains a Sylow 2-subgroup of $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_1)$, or $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_1)$ satisfies to the conditions of Theorem 5.3.1 and $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_1) \cap \mathbf{P}G_1$ is a nontrivial 2-group, in particular p is odd. In any case $|K \cap G|$ is divisible by 2 that contradicts our assumption. Therefore the order $|K_G|$ is even and we may assume that $x \in Z(K) \cap K_G$ is an involution.

Write $\zeta g = \zeta_2 g_1 \cdot \zeta_{2'} g_2$, where $\zeta_2 g_1$ is the 2-part and $\zeta_{2'} g_2$ is the $2'$-part of ζg. By Lemma 4.2.5 the element $\zeta_{2'}$ centralizes a Sylow 2-subgroup Q_G of G, so we may assume that the order of g_2 is odd. Up to conjugation in G we may assume that $\zeta_{2'}$ centralizes a Sylow 2-subgroup of K_G. In particular, $\zeta_{2'}$ centralizes x. Let Q be a Sylow 2-subgroup of $C_G(x)$. Then there exists $y \in G$ such that $Q^y \leq Q_G$. Substituting the subgroup K by its conjugate K^y, we may assume that $\zeta_{2'}$ centralizes a Sylow 2-subgroup of $C_G(x)$. Since $\zeta_{2'} g_2$ centralizes x, we obtain that $g_2 \in C_{\overline{G}_\sigma}(x)$. Moreover, by Lemma 3.2.1 it follows that $g_2 \in C_{\overline{G}}(x)^0$. In particular, g_2 normalizes each G_i and centralizes $Z(C)$ and $Z(C_G(x))$.

Note that $\zeta_{2'}$ normalizes each G_i and centralizes a Sylow 2-subgroup of $Z(C_G(x))$ (recall that $\zeta_{2'}$ centralizes a Sylow 2-subgroup of $C_G(x)$). Indeed, $\zeta_{2'}$ normalizes C, hence normalizes characteristic subgroups $O^{p'}(C) = G_1 * \ldots * G_m$ and $Z(C)$ of C. So we may consider the induced automorphism $\zeta_{2'}$ of

$$O^{p'}(C)/(Z(C) \cap O^{p'}(C) = \mathbf{P}G_1 \times \ldots \times \mathbf{P}G_m.$$

Since each $\mathbf{P}G_i$ has trivial center and can not be written as a direct product of proper subgroups, corollary from Krull-Remak-Schmidt Theorem [38, 3.3.10] implies that $\zeta_{2'}$ permutes distinct $\mathbf{P}G_i$. Since $\zeta_{2'}$ centralizes a Sylow 2-subgroup of $C_G(x)$ and $C \trianglelefteq C_G(x)$, then $\zeta_{2'}$ centralizes a Sylow 2-subgroup of C, hence centralizes a Sylow 2-subgroup $Q_1 \times \ldots \times Q_m$ of $\mathbf{P}G_1 \times \ldots \times \mathbf{P}G_m$, where Q_i is a Sylow 2-subgroup of $\mathbf{P}G_i$. If $\zeta_{2'}$ would induce a nontrivial permutation on the set $\{\mathbf{P}G_1, \ldots, \mathbf{P}G_m\}$, then in would induce a nontrivial permutation on $\{Q_1, \ldots, Q_m\}$. Since each Q_i is nontrivial, this is impossible. Thus every element of odd order of $\langle K, C\rangle$ centralizes a Sylow 2-subgroup of $Z(C)$ and normalizes each G_i.

If $\Phi(\overline{G})=E_6$, then by Lemma 3.2.1 the centralizer of every involution of G in $\overline{G}$ is connected. By Lemma 4.2.2 every involution of G is contained in a maximal torus T such that $N(G,T)/T\simeq W$, where W is a Weyl group of $\overline{G}$. $\overline{C}$ is welknown to be generated by the torus $\overline{T}$ and $\overline{T}$-root subgroups. Write $\overline{C}=\overline{T}(\overline{G}_1*\ldots*\overline{G}_k)$. Since $\overline{T}_\sigma$ either is obtained from a maximal split torus $\overline{H}$ by twisting with an element w_0 of order 2, or is equal to $\overline{H}$, and each field automorphism acts trivially on the factor group $N_{\overline{G}}(\overline{H})/\overline{H}$, then $\bar{\zeta}_{2'}$ normalizes every subgroup $\overline{G}_i$. So, if $\Phi(\overline{G}_i)=D_4$, then $\bar{\zeta}_{2'}$ induces a field (but not a graph or a graph-field) automorphism of $\overline{G}_i$. Moreover, since σ acts trivially on the factor group $N_{\overline{G}}(\overline{T})/\overline{T}$ (see Lemma 4.2.5), then [17, Proposition 6] implies that σ normalizes each $\overline{G}_i$. Therefore, none of G_i is isomorphic to ${}^3D_4(q^3)$. If $\Phi(\overline{G})$ coincides with A_n or D_n, then [18, Propositions 7, 8, 10] imply that none of G_i is isomorphic to ${}^3D_4(q^3)$. Therefore in any case none of G_i is isomorphic to ${}^3D_4(q^3)$. Moreover Lemma 3.2.1 implies that $|K_G:(K_G\cap C)|$ divides $|C_{\overline{G}}(x)/C_{\overline{G}}(x)^0|$ and $C_{\overline{G}}(x)/C_{\overline{G}}(x)^0$ is a 2-group. In [18] it is proven that if a root system Φ has type D_n and Ψ is its subsystem of type D_4, then none element from $N_{W(\Phi)}(W(\Psi))$ induces a symmetry of order 3 of the Dynkin diagram of Ψ. Since ζ^2 is a field automorphism, lack of a symmetry of order 3 together with [17, Proposition 6] implies that for each G_i the automorphism $\zeta_{2'}$ is field (but not graph o graph-field). Therefore the group of induced automorphisms $\langle \mathrm{Aut}_{\widetilde{K}}(\mathbf{P}G_i),\mathbf{P}G_i\rangle$ satisfies to the conditions of Theorem 5.4.1 for all i.

Now consider $\widetilde{G}=\widetilde{K}(\mathbf{P}G_1\times\ldots\times\mathbf{P}G_m)\leq\langle K,C\rangle/Z(C)$ (probably, $m=0$), where $\widetilde{K}=KZ(C)/Z(C)$ is a Carter subgroup of $\widetilde{G}$ (see Lemma 2.4.1) and, for all i, $Z(\mathbf{P}G_i)=\{e\}$. By Lemma 2.2.3 we have that $\mathrm{Aut}_{\widetilde{K}}(\mathbf{P}G_1)$ is a Carter subgroup of $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_1)$. Since $\mathbf{P}G_1$ is a finite group of Lie type satisfying Theorem 5.4.1, by induction we obtain that $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_1)$ satisfies **(ESyl2)**. Similarly we have that $\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_i)$ satisfies **(ESyl2)** for all i. Since

$$\mathrm{Aut}_{\langle K,C\rangle/Z(C)}(\mathbf{P}G_i)\geq\mathrm{Aut}_{\widetilde{G}}(\mathbf{P}G_i),$$

Lemmas 4.3.1 and 4.3.3 imply that $\mathrm{Aut}_{\langle K,C\rangle/Z(C)}(\mathbf{P}G_i)$ satisfies **(ESyl2)**. Since $C_{\langle K,C\rangle/Z(C)}(\mathbf{P}G_1\times\ldots\times\mathbf{P}G_m)=\{e\}$, Lemma 2.4.5 with normal subgroups $C_{\langle K,C\rangle/Z(C)}(\mathbf{P}G_1)\cap N_{\langle K,C\rangle/Z(C)}(\mathbf{P}G_1),\ldots,$ $C_{\langle K,C\rangle/Z(C)}(\mathbf{P}G_m)\cap N_{\langle K,C\rangle/Z(C)}(\mathbf{P}G_1)$ implies that $N_{\langle K,C\rangle/Z(C)}(\mathbf{P}G_1)$ satisfies **(ESyl2)**. Now

$$|\langle K,C\rangle/Z(C):N_{\langle K,C\rangle/Z(C)}(\mathbf{P}G_1)|=2^t,$$

and each element of odd order of $\langle K,C\rangle/Z(C)$ normalizes $\mathbf{P}G_1$, thus, by Lemma 2.4.6, we obtain that the factor group $\langle K,C\rangle/Z(C)$ satisfies **(ESyl2)** and, by Lemma 2.4.5 $\langle K,C\rangle$ satisfies **(ESyl2)**. Since $|\mathbf{P}G_i|<\mathbf{Cmin}$, then $\langle K,C\rangle$ satisfy **(C)**. By Lemma 2.4.3 we obtain that there exists a Carter subgroup F of $\langle K,C\rangle$ containing a Sylow 2-subgroup of $\langle K,C\rangle$. By Theorem 2.1.4 subgroups F and K are conjugate in $\langle K,C\rangle$, thus K contains a Sylow 2-subgroup Q of $\langle K,C\rangle$. Since $|C_G(x):C|$ is a power of 2 and $\langle K,C\rangle$ normalizes $C_G(x)$, we obtain that $|\langle K,C_G(x)\rangle:\langle K,C\rangle|$ is a power of 2. Moreover by construction each element of odd order of $\langle K,C_G(x)\rangle$ is in $\langle K,C\rangle$. Thus by Lemma 2.4.6 $\langle K,C_G(x)\rangle$ satisfies **(ESyl2)** and K contains a Sylow 2-subgroup Q of $\langle K,C_G(x)\rangle$.

Let ΓQ be a Sylow 2-subgroup of $\langle G,\zeta g\rangle$ containing Q and $t\in Z(\Gamma Q)\cap G$. Then $t\in C_G(x)$, hence, $t\in Z(Q)$ and $t\in Z(K)$. Thus we may substitute x by t in arguments above and obtain that $Q=\Gamma Q$, i. e., K contains a Sylow 2-subgroup of $\langle G,\zeta g\rangle$, than completes the proof of Theorem 5.4.1.

5.7 Carter subgroups of finite groups are conjugate

Before we formulate the main theorem, note a corollary of Theorem 5.4.1.

Corollary 5.7.1. **Cmin** $= \infty$, *i. e.* $\mathcal{A} = \varnothing$.

Proof. Indeed, let $\mathcal{A} \neq \varnothing$ and $A \in \mathcal{A}$ is such that the equality $|F^*(A)| =$ **Cmin** holds. Since $F^*(A) = O^{p'}(\overline{G}_\sigma)$ for an adjoint simple connected linear algebraic group $\overline{G}$ and a Frobenius map σ, denote the intersection $A \cap \overline{G}_\sigma$ by G. As we noted in the beginning of subsection 6.1, we may assume that $A = KF^*(A) = KG$. Therefore A satisfies either to the conditions of Theorem 5.3.1, or to the conditions of Theorem 5.4.1. In both cases we have proved that Carter subgroups of A are conjugate, that contradicts to the choice of A. ☐

In order to state the main theorem without using of the classification of finite simple groups, we give the following definition. A finite group is said to be a *K-group* if all its non-Abelian composition factors are known simple groups.

Theorem 5.7.2. (Main Theorem) *Let G be a finite K-group. Then Carter subgroups of G are conjugate.*

Proof. By Theorems 3.3.5, 1.5.6, 5.2.3, 5.3.1, and 5.4.1 of the present paper, and also by [26] we obtain that for each known simple group S and each nilpotent subgroup N of a group of its automorphisms, Carter subgroups of $\langle N, S\rangle$ are conjugate. So G satisfies **(C)**. Hence by Theorem 2.1.4, Carter subgroups of G are conjugate. ☐

From Lemma 2.4.1 and Main Theorem 5.7.2 it follows that a homomorphic image of a Carter subgroup is a Carter subgroup.

Theorem 5.7.3. *Let G be a finite K-group, H a Carter subgroup of G, and N a normal subgroup of G. Then HN/N is a Carter subgroup of G/N.*

6 Existency criterion

6.1 Brief review Краткий обзор результатов параграфа

In this section we shall obtain a criterion of existence of Carter subgroups in a finite group in terms of its normal series. Note that there exist finite groups without Carter subgroups, a minimal counter example is Alt_5. We shall construct an example showing that an essential improvement of the criterion is impossible. At the end of the section, for convenience of the reader, we assemble the classification of Carter subgroups in finite almost simple groups, that is obtained in the present paper.

Recall that in view of Theorem 5.7.2 in every almost simple group with known simple socle Carter subgroups are conjugate. Thus, modulo the classification of finite simple groups, in every finite group Carter subgroups are conjugate. In this section by a finite group we always mean a finite group satisfying **(C)**, thus the results of the section do not depend on the classification of finite simple groups.

Definition 6.1.1. Let $G = G_0 \geq G_1 \geq \ldots \geq G_n = \{e\}$ be a chief series of G (recall that G is assumed to satisfy **(C)**). Then $G_i/G_{i+1} = T_{i,1} \times \ldots \times T_{i,k_i}$, where $T_{i,1} \simeq \ldots \simeq T_{i,k_i} \simeq T_i$ and T_i is a simple group. If $i \geqslant 1$, then denote by $\overline{K}_i$ a Carter subgroup of G/G_i (if it exists) and by

K_i its complete preimage in G/G_{i+1}. If $i=0$, then $\overline{K}_0=\{e\}$ and $K_0=G/G_1$ (note that $\overline{K}_0$ always exists). A finite group G is said to satisfy **(E)**, if for each i,j, either $\overline{K}_i$ does not exists, or $\mathrm{Aut}_{K_i}(T_{i,j})$ contains a Carter subgroup.

By Theorem 6.2.2 and Theorem 5.7.3 it follows that if a finite group satisfies **(E)**, then, for every i, subgroup $\overline{K}_i$ exists, so the first part of condition **(E)** is never satisfied. Recall that by Theorem 5.7.3 a homomorphic image of a Carter subgroup is a Carter subgroup. We shall constantly use this fact.

6.2 Criterion

Below we shall need an additional information on the structure of Carter subgroups in groups of special type. Let A' be a group with a normal subgroup T'. Consider the direct product $A_1\times\ldots\times A_k$, where $A_1\simeq\ldots\simeq A_k\simeq A'$, and its normal subgroup $T=T_1\times\ldots\times T_k$, where $T_1\simeq\ldots\simeq T_k\simeq T'$. Consider the symmetric group Sym_k, acting on $A_1\times\ldots\times A_k$ by $A_i^s=A_{i^s}$, for all $s\in S$ and define X to be equal to a semidirect product $(A_1\times\ldots\times A_k)\leftthreetimes\mathrm{Sym}_k$ (permutation wreath product of A' and Sym_k). Denote by A the direct product $A_1\times\ldots\times A_k$ and by π_i the projection $\pi_i:A\rightarrow A_i$. In these notations the following lemma holds.

Lemma 6.2.1. *Let G be a subgroup of X such that $T\leq G$, $G/(G\cap T)$ is nilpotent, and $(G\cap A)^{\pi_i}=A_i$. Assume also that A is solvable. Let K be a Carter subgroup of G.*
Then $(K\cap A)^{\pi_i}$ is a Carter subgroup of A_i.

Proof. Assume that the statement is false and let G be a counter example of minimal order with k minimal. Then $S=G/(G\cap A)$ is transitive and primitive. Indeed, if S is not transitive, then $S\leq\mathrm{Sym}_{k_1}\times\mathrm{Sym}_{k-k_1}$, hence $G\leq G_1\times G_2$. If we denote by $\psi_i:G\rightarrow G_i$ the natural homomorphism, then $G^{\psi_i}=G_i$ satisfies conditions of the lemma and $K^{\psi_i}=K_i$ is a Carter subgroup of G_i. Clearly $(G\cap A)^{\pi_j}=(G_i\cap A^{\psi_i})^{\pi_j}$, where $i=1$ if $j\in\{1,\ldots,k_1\}$ and $i=2$ if $j\in\{k_1+1,\ldots,k\}$, i. e., the following diagrams are commutative:

$$\begin{tikzcd}
G\cap A \arrow[rr, "\pi_j"] \arrow[dr, "\psi_1"'] & & A_j, \\
& G_1\cap A^{\psi_1} \arrow[ur, "\pi_j"'] &
\end{tikzcd}
\qquad
\begin{tikzcd}
G\cap A \arrow[rr, "\pi_j"] \arrow[dr, "\psi_2"'] & & A_j. \\
& G_2\cap A^{\psi_2} \arrow[ur, "\pi_j"'] &
\end{tikzcd}$$

Thus we obtain the statement by induction. If S is transitive, but is not primitive, let

$$\Omega_1=\{T_1,\ldots,T_m\},\Omega_2=\{T_{m+1},\ldots,T_{2m}\},\ldots,\Omega_l=\{T_{(l-1)m+1},\ldots,T_{lm}\}$$

be a system of imprimitivity. Then it contains a nontrivial nontransitive normal subgroup

$$F'\leq\underbrace{\mathrm{Sym}_m\times\ldots\times\mathrm{Sym}_m}_{l\text{ times}},$$

where $k=m\cdot l$. Consider a complete preimage F of F' in X. Then $G\cap F\leq F_1\times\ldots\times F_l$. Denote by $\psi_i:F\rightarrow F_i$ a natural projection, then $(G\cap F)^{\psi_i}=F_i$. Note that all F_i-s satisfy conditions of the lemma and, if we define $T_i'=T_{(i-1)m+1}\times\ldots\times T_{im}$, then G satisfies conditions of the lemma with $T'=T_1'\times\ldots\times T_l'$ and $A'=F$. By induction we have that $(K\cap F)^{\psi_i}$ is a Carter subgroup of F_i and, if $j\in\{m\cdot(i-1)+1,\ldots m\cdot i\}$, then $\big((K\cap F)^{\psi_i}\cap A^{\psi_i}\big)^{\pi_j}$ is a Carter subgroup of A_j. Since $(G \cap A)^{\pi_j} = \big((K \cap F)^{\psi_i} \cap A^{\psi_i}\big)^{\pi_j}$ (for suitable i), we get the statement by induction.

Let Y' be a minimal normal subgroup of G, contained in T (if Y' is trivial, then T is trivial and we have nothing to prove, since G is nilpotent in this case). Thus Y' is a normal elementary Abelian p-group. Let $Y_i = (Y')^{\pi_i}$, then $Y = Y_1 \times \ldots \times Y_k$ is a nontrivial normal subgroup of G (Y is a subgroup of G since $T \leq G$). Let $\bar{\pi}_i : (G \cap A) \rightarrow A_i/Y_i = \overline{A}_i$ be a projection, corresponding to π_i. Denote by $\overline{K} = KY/Y$ a corresponding Carter subgroup of $\overline{G} = G/Y$. Then $\overline{G}$ satisfies conditions of the Lemma. By induction, $(\overline{K} \cap \overline{A})^{\bar{\pi}_i}$ is a Carter subgroup of $\overline{A}_i$. Let K_1 be a complete preimage of $\overline{K}$ in G, and let Q be a Hall p'-subgroup of K_1. Then $(Q \cap A)^{\pi_i}$ is a Hall p'-subgroup of $(K_1 \cap A)^{\pi_i}$. In view of the proof of [8, Theorem 20.1.4], we obtain that $K = N_{K_1}(Q)$ is a Carter subgroup of G and $(N_{K_1 \cap A}(Q \cap A))^{\pi_i}$ is a Carter subgroup of A_i. Thus we need to show that $(N_{K_1 \cap A}(Q \cap A))^{\pi_i} = (N_{K_1 \cap S}(Q))^{\pi_i}$. By induction the equality $(N_{\overline{K} \cap \overline{A}}(\overline{A} \cap \overline{Q}))^{\bar{\pi}_i} = (N_{\overline{K} \cap \overline{G}}(\overline{Q}))^{\bar{\pi}_i}$ holds. Thus we need to prove that $(N_Y(Q \cap A))^{\pi_i} = (N_Y(Q))^{\pi_i}$. Note also that $(N_Y(Q \cap A))^{\pi_i} \leq N_{Y_i}((Q \cap A)^{\pi_i})$.

Since S is transitive and primitive subgroup of Sym_k, then $k = r$ is a prime and $S = \langle s \rangle$ is cyclic. If $r = p$, then $Q \cap A = Q$ and we have nothing to prove. Otherwise let h be an r-element of K, generating S modulo $K \cap A$. Clearly $Q = (Q \cap A)\langle h \rangle$. Let $t \in Y_i$ be an element of $N_{Y_i}((Q \cap A)^{\pi_i})$. Then $(t \cdot t^h \cdot \ldots \cdot t^{h^{r-1}}) \in N_Y(Q)$ and $t^{\pi_i} = (t \cdot t^h \cdot \ldots \cdot t^{h^{r-1}})^{\pi_i}$, hence $(N_Y(Q \cap A))^{\pi_i} \leq N_{Y_i}((Q \cap A)^{\pi_i}) \leq (N_Y(Q))^{\pi_i} \leq (N_Y(Q \cap A))^{\pi_i}$. □

Theorem 6.2.2. *Let G be a finite group. Then G contains a Carter subgroup if and only if G satisfies* **(E)**.

Proof. We prove the part "only if" first. Let H be a minimal normal subgroup of G. Then $H = T_1 \times \ldots \times T_k$, where $T_1 \simeq \ldots \simeq T_k \simeq T$ are simple groups.

If H is elementary Abelian (i. e., T is cyclic of prime order), then $\mathrm{Aut}(T)$ is solvable and contains a Carter subgroup. Assume that T is a non-Abelian simple group. Clearly K is a Carter subgroup of KH. By Lemma 2.2.3 we obtain that $\mathrm{Aut}_{KH}(T_i)$ contains a Carter subgroup for all i. Induction by the order of the group completes the proof of necessity.

Now we prove the "if" part. Again assume by contradiction that G is a counter example of minimal order, i. e., that G does not contain a Carter subgroup, but G satisfies **(E)**. Let H be a minimal normal subgroup of G. Then $H = T_1 \times \ldots \times T_k$, where $T_1 \simeq \ldots \simeq T_k \simeq T$, and T is a finite simple group.

By definition, G/H satisfies **(E)**, thus, by induction, there exists a Carter subgroup $\overline{K}$ of $\overline{G} = G/H$. Let K be a complete preimage of $\overline{K}$, then K satisfies **(E)**. If $K \neq G$, then by induction K contains a Carter subgroup K'. Note that K' is a Carter subgroup of G. Indeed, assume that $x \in N_G(K') \setminus K'$. Since $K'H/H = \overline{K}$ is a Carter subgroup of $\overline{G}$, we obtain that $x \in K$. But K' is a Carter subgroup of K, thus $x \in K'$. Hence $G = K$, i. e., G/H is nilpotent.

If H is Abelian, then G is solvable, therefore, G contains a Carter subgroup. So assume that T is a non-Abelian finite simple group. We shall show first that $C_G(H)$ is trivial. Assume that $C_G(H) = M$ is nontrivial. Since T is a non-Abelian simple group, it follows that $M \cap H = \{e\}$, so M is nilpotent. By Lemma 2.1.2 we obtain that G/M satisfies **(E)**. By induction we obtain that G/M contains a Carter subgroup $\overline{K}$. Let K' be a complete preimage of $\overline{K}$ in G. Then K' is solvable, therefore contains a Carter subgroup K. Like above we obtain that K is a Carter subgroup of G, a contradiction. Hence $C_G(H) = \{e\}$.

Since H is a minimal normal subgroup of G, we obtain that $\mathrm{Aut}_G(T_1) \simeq \mathrm{Aut}_G(T_2) \simeq \ldots \simeq \mathrm{Aut}_G(T_k)$. Thus there exists a monomorphism

$$\varphi : G \rightarrow (\mathrm{Aut}_G(T_1) \times \ldots \times \mathrm{Aut}_G(T_k)) \leftthreetimes \mathrm{Sym}_k = G_1$$

and we identify G with G^φ. Denote by K_i a Carter subgroup of $\mathrm{Aut}_G(T_i)$ and by A the subgroup $\mathrm{Aut}_G(T_1) \times \ldots \times \mathrm{Aut}_G(T_k)$. Since G/H is nilpotent then $K_iT_i = \mathrm{Aut}_G(T_i)$ and $G_1 = (K_1T_1 \times \ldots \times K_kT_k) \leftthreetimes \mathrm{Sym}_k$. Let $\pi_i : G \cap A \to (G \cap A)/C_{(G\cap A)}(T_i)$ be canonical projections. Since $G/(G \cap A)$ is transitive, we obtain that $(G \cap A)^{\pi_i} = K_iT_i$.

Since $\mathrm{Aut}_{G\cap A}(T_i) = K_iT_i$, then $G \cap A$ satisfies **(E)**. By induction it contains a Carter subgroup M. By Lemma 2.2.3 we obtain that M^{π_i} is a Carter subgroup of K_iT_i, therefore we may assume that $M^{\pi_i} = K_i$. In particular, if $R = (K_1 \cap T_1) \times \ldots \times (K_k \cap T_k)$, then $M \leq N_G(R)$. In view of Theorems 2.1.4 and 5.7.2, Carter subgroups in each finite group are conjugate. Since $(G\cap A)/H$ is nilpotent, we get that $G\cap A = MH$, so $G = N_G(M)H$. More over $N_G(M) \cap A = M$, hence $N_G(M)$ is solvable. Since M normalizes R, and $M^{\pi_i} = K_i$, we obtain that $N_G(M)$ normalizes R, so $N_G(M)R$ is solvable. Therefore it contains a Carter subgroup K. By Lemma 6.2.1, $(K \cap A)^{\pi_i}$ is a Carter subgroup of $(N_G(M)R \cap A)^{\pi_i}$ (R play the role of subgroup T from Lemma 6.2.1 in this case), so $(K \cap A)^{\pi_i} = K_i$. Assume that $x \in N_G(K) \setminus K$. Since $G/H = N_G(M)H/H = KH/H$, it follows that $x \in H$. Therefore $x^{\pi_i} \in (N_G(K) \cap A)^{\pi_i} \leq N_{T_i}((K \cap A)^{\pi_i}) = K_i$. Since $\bigcap_i \mathrm{Ker}(\pi_i) = \{e\}$, it follows that $x \in R \leq N_G(M)R$. But K is a Carter subgroup of $N_G(M)R$, hence $x \in K$. This contradiction completes the proof. ☐

6.3 Example

In this subsection we shall construct an example, showing that we cannot substitute condition **(E)** by a weaker condition: for each composition factor S of G, $\mathrm{Aut}_G(S)$ contains a Carter subgroup. This example also shows that an extension of a group containing a Carter subgroup, by a group, containing a Carter subgroup, may fail to contain a Carter subgroup.

Consider $L = \mathbf{P}\mathrm{SL}_2(3^3) \leftthreetimes \langle \varphi \rangle$, where φ is a field automorphism of $\mathbf{P}\mathrm{SL}_2(3^3)$. Let $X = (L_1 \times L_2) \leftthreetimes \mathrm{Sym}_2$, where $L_1 \simeq L_2 \simeq L$ and if $\sigma = (1, 2) \in \mathrm{Sym}_2 \setminus \{e\}$, $(x, y) \in L_1 \times L_2$, then $\sigma(x, y)\sigma = (y, x)$ (permutation wreath product of L and Sym_2). Denote by $H = \mathbf{P}\mathrm{SL}_2(3^3) \times \mathbf{P}\mathrm{SL}_2(3^3)$ a minimal normal subgroup of X and by $M = L_1 \times L_2$. Let $G = (H \leftthreetimes \langle (\varphi, \varphi^{-1}) \rangle) \leftthreetimes \mathrm{Sym}_2$ be a subgroup of X. Then the following statements hold:

1. For every composition factor S of G, $\mathrm{Aut}_G(S)$ contains a Carter subgroup.
2. $G \cap M \trianglelefteq G$ contains a Carter subgroup.
3. $G/(G \cap L)$ is nilpotent.
4. G does not contain a Carter subgroup.

1. Clearly we need to check the statement only for non-Abelian composition factors. Every non-Abelian composition factor S of G is isomorphic to $\mathbf{P}\mathrm{SL}_2(3^3)$ and $\mathrm{Aut}_G(S) = L$. By Theorem 5.4.1, L contains a Carter subgroup (that is equal to a Sylow 3-subgroup).

2. Since $(G \cap M)/H$ is nilpotent, and from the previous statement we obtain that $G \cap M$ satisfies **(E)**, so it contains a Carter subgroup (it is easy to see that a Sylow 3-subgroup of $G \cap M$ is a Carter subgroup of $G \cap M$).

3. Evident.

4. Assume that K is a Carter subgroup of G. Then KH/H is a Carter subgroup of G/H. But G/H is a non-Abelian group of order 6, hence $G/H \simeq \mathrm{Sym}_3$ and KH/H is a Sylow 2-subgroup of G/H. By Lemma 2.1.2 $\mathrm{Aut}_K(\mathbf{P}\mathrm{SL}_2(3^3))$ is a Carter subgroup of $\mathrm{Aut}_{KH}(\mathbf{P}\mathrm{SL}_2(3^3)) = \mathbf{P}\mathrm{SL}_2(3^3)$. But $\mathbf{P}\mathrm{SL}_2(3^3)$ does not contain Carter subgroups in view of Theorem 5.4.1.

6.4 Classification of Carter subgroups

In view of condition **(E)** and Theorem 6.2.2, the description of Carter subgroups in finite groups is reduced to the classification of Carter subgroups in almost simple groups A with $A/F^*(A)$ nilpotent. The classification of Carter subgroups in groups with this condition is obtained in previous sections and we give it here for convenient usage.

We prove first the following theorem, showing that if, for a subgroup S of $\mathrm{Aut}(G)$, there exists a Carter subgroup, then it exists in every larger group $S \leq A \leq \mathrm{Aut}(G)$ (here G is a known simple group).

Theorem 6.4.1. *Let G be a finite simple group and $G \leq A \leq \mathrm{Aut}(G)$ an almost simple group with simple socle G. Assume that A contains a subgroup S such that $G \leq S$ and S contains a Carter subgroup.*
Then A contains a Carter subgroup.

Proof. Let K be a Carter subgroup of S. Clearly we may assume that $S = KG$.

Assume that either $G \simeq \mathrm{Alt}_n$ for some $n \geqslant 5$, or G is sporadic. Since by Lemma 3.2.11 each element of odd prime order of G is conjugate to its inverse, and since $|\mathrm{Aut}(G) : G|$ is a 2-power, Lemmas 2.4.2 and 2.4.6 imply that if some group $G \leq S \leq \mathrm{Aut}(G)$ contains a Carter subgroup K, then K, is a Sylow 2-subgroup of S. Since $|A : S|$ is a 2-power, the statement of the theorem in this case follows from Lemma 2.4.6.

Assume that $G = {}^3D_4(q)$. By [44, Theorem 1.2(vi)] each element of G is conjugate to its inverse. If q is odd, then Lemma 4.2.5 implies that K is a Sylow 2-subgroup of S. So by Lemmas 2.4.6 and 4.2.5 it follows that A satisfies **(ESyl2)**, i. e., contains a Carter subgroup. If $q = 2^t$ is even, then by Theorems 5.3.1 and 5.4.1 it follows that $S = \mathrm{Aut}(G)$ and we have nothing to prove.

Assume that G is a group of Lie type, $G \not\simeq {}^3D_4(q)$ and, if $G \simeq D_4(q)$, then $S \leq A_1$, where $A_1 \leq \mathrm{Aut}(D_4(q))$ is defined in Theorem 5.3.1. Then S satisfies one of conditions (a)–(d) of Theorem 5.4.1. Consider all these cases separately.

Assume that S satisfies (a). In this case we have $|\mathrm{Aut}(G) : S| \leqslant 2$ and so, for each A such that $S \leq A \leq \mathrm{Aut}(G)$, either $A = S$, or $A = \mathrm{Aut}(G)$. In any case A satisfies to statement (a) of Theorem 5.4.1 and contains a Carter subgroup.

Assume that S satisfies to statement (b). Then $|\mathrm{Aut}(G) : S| = 2$ and either $A = S$, or $A = \mathrm{Aut}(G)$. In the first case we have nothing to prove. In the second case $\widehat{G} = \mathbf{P}GL_2(3^t)$ satisfies **(ESyl2)**, hence by Lemma 4.3.3 group A also satisfies **(ESyl2)** and by Lemma 2.4.3 contains a Carter subgroup.

Assume that S satisfies statement (c) of Theorem 5.4.1. Then $S = \mathrm{Aut}(G)$ and we have nothing to prove.

Assume that S satisfies condition (d) of Theorem 5.4.1. By Lemma 4.3.1, $S \cap \widehat{G}$ satisfies **(ESyl2)**. By Lemma 4.3.3 every subgroup A of $\mathrm{Aut}G$, containing $S \cap \widehat{G}$ also satisfies **(ESyl2)**, hence by Lemma 2.4.3, it contains a Carter subgroup.

Now assume that $G = D_4(q)$ and S satisfies conditions of Theorem 5.3.1. Since graph automorphisms of orders 2 and 3 do not commute, only one of them can be contained in a nilpotent subgroup. Thus we may assume that only one of them is contained in A. Then every subgroup A containing S, either satisfies to Theorem 5.3.1, or satisfies to Theorem 5.4.1, condition (a), if q is even and condition (d), if q is odd, i. e., it contains a Carter subgroup. ☐

Note that from Theorem 6.4.1 and [9] an interesting corollary follows.

Lemma 6.4.2. *Let S be a known finite simple group, $S \not\simeq J_1$ and $G = \mathrm{Aut}(S)$. Then G contains a Carter subgroup.*

Proof. By [9, Theorems 2 and 3], if S is not of Lie type and is not equal to J_1, then group of its automorphisms $\mathrm{Aut}(S)$ satisfies **(ESyl2)** and, by Lemma 2.4.3, contains a Carter subgroup. Now, if S is of Lie type in even characteristic, then $\mathrm{Aut}(S)$ contains a Carter subgroup in view of Theorem 5.4.1(a). If S is of Lie type in odd characteristic and $S \not\simeq {}^2G_2(3^{2n+1})$, then $\widehat{S}$ satisfies **(ESyl2)**, so contains a Carter subgroup by Lemma 2.4.3. By Theorem 6.4.1, $\mathrm{Aut}(S)$ contains a Carter subgroup. Finally, if $S \simeq {}^2G_2(3^{2n+1})$, then $\mathrm{Aut}(S)$ contains a Carter subgroup in view of Theorem 5.4.1(c). ☐

Tables given below are arranged in the following way. In the first column is given a simple group S such that Carter subgroups of $\mathrm{Aut}(S)$ are classified. In the second column we give conditions for a subgroup A of its group of automorphisms for A to contain a Carter subgroup. In the third column we give the structure of a Carter subgroup K. In every subgroup of $\mathrm{Aut}(S)$ lying between S and A Carter subgroups does non exist. By $P_r(G)$ a Sylow r-subgroup of G is denoted. By φ we denote a field automorphism of a group of Lie type S, by τ we denote a graph automorphism of a group of Lie type S contained in K (since graph automorphisms of order 2 and 3 of $D_4(q)$ does not commute, only one of them can be in K). If A does not contains a graph automorphism, then we suppose $\tau = e$. By ψ we denote a field automorphism of S of maximal order contained in A (it is a power of φ, but $\langle\psi\rangle$ can be different from $\langle\varphi\rangle$). By $K(U_3(2))$ a Carter subgroup of order $2 \cdot 3$ of $\widehat{{}^2A_2(2)}$ is denoted. If G is solvable, then by $K(G)$ we denote a Carter subgroup of G. In Table 10 by ζ is denoted a graph-field automorphism of order $2t$ of $A_2(2^{2t})$.

Table 7: Groups of automorphisms of alternating groups, containing Carter subgroups

Group S	Conditions on A	Structure of K
Alt_5	$A = \mathrm{Sym}_5$	$K = P_2(\mathrm{Sym}_5)$
Alt_n, $n \geqslant 6$	none	$K = P_2(S)$

Table 8: Groups of automorphisms of sporadic groups, containing Carter subgroups

Group S	Conditions on A	Structure of K
J_2, J_3, Suz, HN	$A = \mathrm{Aut}(S)$	$K = P_2(A)$
$\not\simeq J_1, J_2, J_3, Suz, HN$	none	$K = P_2(A)$

Table 9: Groups of automorphisms of exceptional groups of Lie type, containing Carter subgroups

Group S	Conditions on A	Structure of K
${}^2B_2(2^{2n+1})$, $n \geqslant 1$	$A = \mathrm{Aut}(S)$	$K = \langle\varphi\rangle \times P_2({}^2B_2(2))$
$({}^2F_4(2))'$	none	$K = P_2(A)$
${}^2F_4(2^{2n+1})$, $n \geqslant 1$	$A = \mathrm{Aut}(S)$	$K = \langle\varphi\rangle \times P_2({}^2F_4(2))$
${}^2G_3(3^{2n+1})$	$A = \mathrm{Aut}(G)$	$\langle\varphi\rangle \leftthreetimes (2 \times P)$, where $\lvert P\rvert = 3^{\lvert\varphi\rvert_3}$
remaining, q odd	none	$K = P_2(A) \times K(O(N_A(P_2(A))))$
remaining, $q = 2^t$	$\varphi g \in A$, $g \in \widehat{S}$	$\langle\tau, \varphi\rangle \leftthreetimes P_2(S_{\varphi_{2'}})$

Table 10: Groups of automorphisms of classical groups, containing Carter subgroups

Group S	Conditions on A	Structure of K
$A_1(q)$, $q \equiv \pm 1 \pmod 8$	none	$K = N_A(P_2(S))$
$A_1(q)$, $q \equiv \pm 3 \pmod 8$	$\widehat{S} \leq A$	$K = N_A(P_2(\widehat{S}))$
$A_n(2^t)$, $t \geqslant 2$, if $n = 1$	$\varphi g \in A$, $g \in \widehat{S}$	$K = \langle \varphi, \tau \rangle \rightthreetimes S_{\varphi_{2'}}$
$A_2(2^{2t})$, $3 \nmid t$	$\langle S, \zeta g \rangle \leq A \leq S \leftthreetimes \langle \zeta \rangle$, $C_{A \cap \widehat{S}}(\varphi_{2'}) \simeq \mathbf{P}\mathrm{GU}_3(2)$	$K = \langle \zeta g \rangle \times K(\mathbf{P}\mathrm{GU}_3(2))$
$A_n(q)$, q odd, $n \geqslant 2$	none	$K = P_2(A) \times K(O(N_A(P_2(A))))$
${}^2A_2(2^t)$, t odd, $3 \nmid t$	$\langle S, \varphi_{2'} g \rangle \leq A \leq \widehat{S} \leftthreetimes \langle \varphi_{2'} \rangle$ $C_{A \cap \widehat{S}}(\varphi_{2'}) \simeq \mathbf{P}\mathrm{GU}_3(2)$ $C_{A \cap \widehat{S}}(\varphi_{2'}) \simeq \mathbf{P}\mathrm{SU}_3(2)$	$K = \langle \varphi_{2'} \rangle \times K(\mathbf{P}\mathrm{GU}_3(2))$ $K = \langle \varphi_{2'} \rangle \times P_2(\mathbf{P}\mathrm{SU}_3(2))$
${}^2A_2(2^t)$	$A = \mathrm{Aut}(S)$	$K = \langle \varphi \rangle \rightthreetimes P_2(S_{\varphi_{2'}})$
${}^2A_n(q)$, q odd	none	$K = P_2(A) \times K(O(N_A(P_2(A))))$
${}^2A_n(2^t)$, $n \geqslant 3$	$A = \mathrm{Aut}(S)$	$K = \langle \varphi \rangle \rightthreetimes P_2(S_{\varphi_{2'}})$
$B_2(q)$, $q \equiv \pm 1 \pmod 8$	none	$K = P_2(A) \times K(O(N_A(P_2(A))))$
$B_2(2^t)$, $t \geqslant 2$	$\varphi \in A$	$K = \langle \varphi, \tau \rangle \rightthreetimes P_2((S_\tau)_\varphi)$
$B_2(q)$, $q \equiv \pm 3 \pmod 8$	$\widehat{S} \leq A$	$K = P_2(A) \times K(O(N_A(P_2(A))))$
$B_n(q)$, q odd, $n \geqslant 3$	none	$K = P_2(A) \times K(O(N_A(P_2(A))))$
$C_n(q)$, $q \equiv \pm 1 \pmod 8$	none	$K = P_2(A) \times K(O(N_A(P_2(A))))$
$C_n(q)$, $q \equiv \pm 3 \pmod 8$	$\widehat{S} \leq A$	$K = P_2(A) \times K(O(N_A(P_2(A))))$
$C_n(2^t)$, $n \geqslant 3$	$A = \mathrm{Aut}(S)$	$K = \langle \varphi \rangle \times P_2(S_{\varphi_{2'}})$

$D_4(q)$, q odd	none	if $\lvert\tau\rvert \leqslant 2$, then
		$K = P_2(A) \times K(O(N_A(P_2(A))))$;
		if $\lvert\tau\rvert = 3$, then
		$K = \langle \tau, \psi \rangle \rightthreetimes P_2(S_\tau)$

$D_4(2^t)$	$\varphi \in A$	if $\lvert\tau\rvert \leqslant 2$, then
		$K = \langle \tau, \varphi \rangle \rightthreetimes P_2(S_{\varphi_{2'}})$;
		if $\lvert\tau\rvert = 3$, then
		$K = \langle \tau, \varphi \rangle \rightthreetimes P_2((S_\tau)_{\varphi_{2'}})$

$D_n(q)$, q odd, $n \geqslant 5$	none	$K = P_2(A) \times K(O(N_A(P_2(A))))$
$D_n(2^t)$, $n \geqslant 5$	$\varphi \in A$	$K = \langle \tau, \varphi \rangle \rightthreetimes P_2(S_{\varphi_{2'}})$
${}^2D_n(q)$, q odd	none	$K = P_2(A) \times K(O(N_A(P_2(A))))$
${}^2D_n(2^t)$	$A = \mathrm{Aut}(S)$	$K = \langle \varphi \rangle \rightthreetimes P_2(S_{\varphi_{2'}})$

List of Tables

1 Root systems and extended Dynkin diagrams 6

2 Groups of Lie type which are not simple 7

3 Finite almost simple groups, which are not minimal counter examples 16

4 Unipotent classes in $G_2(q)$, $q = 3^t$. 19

5 Unipotent classes $F_4(q)$, $q = 3^t$ 20

6 Unipotent classes in $E_6(\overline{\mathbb{F}}_3)$ 21

7 Groups of automorphisms of alternating groups, containing Carter subgroups . 55

8 Groups of automorphisms of sporadic groups, containing Carter subgroups . . . 55

9 Groups of automorphisms of exceptional groups of Lie type, containing Carter subgroups 55

10 Groups of automorphisms of classical groups, containing Carter subgroups . . . 56

References

[1] A. Borel et al., *Seminar on linear algebraic groups*, Lecture notes in Mathematics, **131** (1970), Springer.

[2] N.A. Vavilov, Nilpotent self-normalizing subgroups of the general linear group over a finite field, *Zap. nauchn. Sem. Leningrad Otdel Mat. Inst. Steklov* LOMI, **86**, (1979), 34–39.

[3] E.P. Vdovin, On the conjugacy problem for Carter subgroups, *SMJ*, **47**, N 4 (2006), 597–600.

[4] E.P. Vdovin, Carter subgroups of almost simple groups, *Algebra and logic*, **46**, N 2 (2007) , 90–119.

[5] E.P. Vdovin, On the existence of Carter subgroups, *Proc. Stekl. Inst. Math. Suppl. 1*, **13**, N 1 (2007), 79–88.

[6] E.P. Vdovin, Conjugacy of Carter subgroups in finite groups, *Doklady RAN*, **415**, N 3 (2007), 300–303 (In Russian).

[7] E.B. Dynkin , Semisimple subalgebras of semisimple Lie algebras, *Mat. sbornik*, **30**, N 2 (1952), 349–462 (In Russian).

[8] M.I. Kargapolov, Yu.I. Meralyzkov, *Foundations of group theory*, Moscow, «Nauka», Fizmatlit, 1996 (In Russian).

[9] A.S. Kondratiev, Normalizers of the Sylow 2-subgroups in finite simple groups, *Math.notes*, **78**, N 3 (2005), 338–346.

[10] A.S. Kondrat'ev, V.D. Mazurov, 2-signalizers of finite simple groups, *Algebra and logic*, **42**, N 5 (2003), 333–348.

[11] V.M. Levchuk, Ya.N. Nuzhin, Structure of Ree groups, *Algebra and logic*, **24**, N 1 (1985), 16–26.

[12] J. E. Humphreys, *Linear algebraic groups*, Springer-Verlag, New York, 1972.

[13] A. Borel and J. de Siebental, Les-sous-groupes fermés de rang maximum des groupes de Lie clos, *Comment. Math. Helv.*, **23**, (1949), 200–221.

[14] R.W. Carter, Nilpotent self-normalizing subgroups of soluble groups, *Math. Z.*, **75**, (1961), 136–139.

[15] R.W. Carter, *Simple groups of Lie type*, John Wiley and Sons, 1972.

[16] R.W. Carter, Conjugacy classes in the Weyl group, *Compositio Mathematica*, **25**, N 1 (1972), 1–59.

[17] R.W. Carter, Centralizers of semisimple elements in finite groups of Lie type, *Proc. London Math. Soc.* (3), **37**, N 3 (1978), 491–507.

[18] R.W. Carter, Centralizers of semisimple elements in the finite classical groups, *Proc. London Math. Soc.* (3), **42**, N 1 (1981), 1–41.

[19] R.W. Carter, *Finite groups of Lie type, conjugacy classes and complex characters*, John Wiley and Sons, 1985.

[20] R.W. Carter, P. Fong, The Sylow 2−subgroups of the finite classical groups, *J. Algebra*, **1**, (1964), 139–151.

[21] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker, and R.A. Wilson, *Atlas of finite groups*, Clarendon Press, Oxford (1985).

[22] A. D'aniello, Sull' esistenza di sottogruppi nilpotenti autonormalizzanti in alcuni gruppi semplici, II, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.*, **74**, (1983), 1–6.

[23] F. Dalla Volta, A. Lucchini, and M.C. Tamburini, On the conjugacy problem for Carter subgroups, *Comm. Algebra*, **26**, N 2 (1998), 395–401.

[24] D. Deriziotis, *Conjugacy classes and centralizers of semisimple elements in finite groups of Lie type*, Vorlesungen aus dem Fachbereich Mathematic der Universität Essen, **11** 1984.

[25] L. Di Martino and M.C. Tamburini, I sottogruppi nilpotenti autonormalizzanti di S_n e di A_n, *Istit. Lombardo Accad. Sci. Lett. Rend. A*, **110**, (1976), 235–241.

[26] L. Di Martino and M.C. Tamburini, Carter subgroups of projective linear groups, *Boll. Un. Mat. Ital. B*, **7**, (1987), 905–915.

[27] L. Di Martino, M.C. Tamburini and A.E. Zalesskii, Carter subgroups in classical groups, *J. London Math. Soc.* (2), **55**, (1997), 264–276.

[28] H. Enomoto, The conjugacy classes of Chevalley groups of type (G_2) over finite fields of characteristic 2 or 3, *Journal of the Faculty of Science University of Tokio, section I*, **16**, N 3 (1970), 497–512.

[29] W. Feit, J. Thompson, Solvability of groups of odd order, *Pacif. J. Math.*, **13**, N 3 (1963), 775–1029.

[30] W. Feit and G. Zukerman, Reality properties of conjugacy classes in spin groups and symplectic groups, *Algebraists' homage:papers in ring theory and related topics*, New Haven, 1981, 239–253.

[31] D. Gorenstein, R. Lyons, R. Solomon, *The classification of the finite simple groups. Number 3. Part I. Chapter A. Almost simple K-groups.* Mathematical Surveys and Monographs, **40**, N 3, American Mathematical Society, Providence, RI, 1998.

[32] D. Gorenstein, R. Lyons, *The local structure of finite groups of characteristic* 2 *type*, Mem. AMS, **42**, (1983), ISSN 0065-9266.

[33] B. Hartley, G. Shute, Monomorphisms and direct limits of finite groups of Lie type, *Quart.J.Math.Oxford Ser. (2)*, **35**, N 137 (1984), 49–71.

[34] J.E. Humphreys, *Conjugacy classes in semisimple algebraic groups*, American Mathematical Society, Providence, Rhode Island, Mathematical Survey and Monographs, **43**, 1995.

[35] F. Menegazzo, M.C. Tamburini, A property of the normalizers of Sylow p-subgroups in simple groups, *Quad.Sem.Mat.Brescia*, 45/02 (2002).

[36] K. Mizuno, The conjugate classes of Chevalley groups of type E_6, *Journal of the Faculty of Science University of Tokio, section IA* **24**, N 3 (1977), 525–565.

[37] A. Previtali, M.C. Tamburini, E.P. Vdovin, The Carter subgroups of some classical groups, *Bull. London Math. Soc.*, **36**, N 1 (2004), 145–155.

[38] D.J.S. Robinson, *A Course in the Theory of Groups*, Springer, 1996.

[39] T. Shoji, The conjugacy classes of Chevalley groups of type (F_4) over finite fields of characteristic $p \neq 2$, *Journal of the Faculty of Science University of Tokio, section IA* **21**, N 1 (1974), 1–19.

[40] G.M. Seitz, Unipotent elements, tilting modules, and saturation, *Inv. Math*, **141**, N 3 (2000), 467–502.

[41] R. Steinberg, Automorphisms of finite linear groups, *Canad. J. Math.*, **12**, N 4 (1960), 606–615.

[42] R. Steinberg, *Endomorphisms of algebraic groups*, Mem.AMS, **80**, 1968.

[43] M.C. Tamburini, E.P. Vdovin, Carter subgroups of finite groups, *J. Algebra*, **255**, N 1 (2002), 148–163.

[44] P.H. Tiep, A.E. Zalesski, Real conjugacy classes in algebraic groups and finite groups of Lie type, *J. Group Theory*, **8**, N 3 (2005), 291–315.

[45] H.N. Ward, On Ree's series of simple groups, *Trans. Amer. Math. Soc.*, **121**, N 1 (1966), 62–80.

[46] M.J. Wonenburger, Transformations which are products of involutions, *J. Math. Mech.*, **16**, N 1 (1966), 32–38.